

**Acetylsalicylic acid**

Section	Indications
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	<div>Acute rheumatic fever without mention of heart involvement</div> <div>Mucocutaneous lymph node syndrome</div> <div>Juvenile idiopathic arthritis</div>

**Adalimumab**

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	<div>Juvenile idiopathic arthritis</div>

**Allopurinol**

Section	Indications
Medicines used to treat gout Oral > Solid: 100 mg	<div>Gout</div>

**Anakinra**

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	<div>Juvenile systemic arthritis</div>

**Azathioprine**

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid > tablet: 50 mg (scored)	<div>Rheumatoid arthritis, serology unspecified</div> <div>Psoriatic arthritis</div>

**Certolizumab pegol**

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for <div>Juvenile idiopathic arthritis</div>

## Chloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)

### Indications

Rheumatoid arthritis

## Colchicine

### Section

Medicines used to treat gout

Oral > Solid: 0.5 mg

### Indications

Gout

## Etanercept

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to adalimumab for

Juvenile idiopathic arthritis

## Golimumab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to adalimumab for

Juvenile idiopathic arthritis

## Hydroxychloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 200 mg (as sulfate)

### Indications

Systemic lupus erythematosus

Cutaneous lupus erythematosus of unspecified type

## Infliximab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to adalimumab for

Juvenile idiopathic arthritis

## Methotrexate

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Medicines for juvenile joint diseases

Oral > Solid > tablet: 2.5 mg (as sodium salt)

### Indications

Rheumatoid arthritis, serology unspecified

Juvenile idiopathic arthritis

## Penicillamine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 250 mg

### Indications

Rheumatoid arthritis, serology unspecified

## Probenecid

### Section

Medicines used to treat gout

Oral > Solid: 500 mg

### Indications

Gout

## Sulfasalazine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 500 mg

### Indications

Rheumatoid arthritis, serology unspecified

## Tocilizumab

### Section

Medicines for juvenile joint diseases

Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial

Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe

### Indications

Juvenile systemic arthritis

## Triamcinolone hexacetonide

### Section

Medicines for juvenile joint diseases

Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

### Indications

Juvenile idiopathic arthritis