

Caffeine citrate

Section

Medicines administered to the neonate [c]
 Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL)
 Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)

Indications

Apnoea of newborn

Chlorhexidine

Section

Medicines administered to the neonate [c]
 Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)
 Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine)

Indications

Omphalitis of newborn

Ibuprofen

Section

Medicines administered to the neonate [c]
 Parenteral > General injections > IV: 5 mg per mL

Indications

Patent arterial duct

Indometacin

Section

Medicines administered to the neonate [c]

Indications

Therapeutic equivalent to ibuprofen for

Patent arterial duct

Prostaglandin E1

Section

Medicines administered to the neonate [c]
 Parenteral > General injections > IV: 0.5 mg per mL in alcohol

Indications

Other specified congenital anomaly of great arteries including arterial duct

Prostaglandin E2

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 1 mg per mL

Indications

Therapeutic equivalent to prostaglandin E1 for

Other specified congenital anomaly of great arteries including arterial duct

Other specified congenital anomaly of great arteries including arterial duct

Surfactant

Section

Medicines administered to the neonate [c]

Respiratory > Suspension: 80 mg per mL for intratracheal instillation; 25 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn