



# Model List of Essential Medicines

## Acetylsalicylic acid

Section	Indications
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Acute rheumatic fever without mention of heart involvement Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis

## Adalimumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

## Anakinra

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	Juvenile systemic arthritis

## Certolizumab pegol

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

## Etanercept

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

## Golimumab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

## Infliximab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to <a href="#">adalimumab</a> for <b>Juvenile idiopathic arthritis</b>

## Methotrexate

Section	Indications
Medicines for juvenile joint diseases Oral > Solid > tablet: 2.5 mg (as sodium salt)	<b>Juvenile idiopathic arthritis</b>

## Tocilizumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe	<b>Juvenile systemic arthritis</b>

## Triamcinolone hexacetonide

Section	Indications
Medicines for juvenile joint diseases Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial	<b>Juvenile idiopathic arthritis</b>