

Acetylsalicylic acid

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs) Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Pain

Amitriptyline

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 25 mg; 10 mg; 75 mg	Palliative care

Codeine

Section	Indications
Opioid analgesics Oral > Solid: 30 mg tablet (codeine phosphate)	Pain

Cyclizine

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 50 mg tablet Parenteral > General injections > unspecified: 50 mg per mL	Palliative care

Dexamethasone

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per 5 mL Oral > Solid: 2 mg; 4 mg Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)	Palliative care

Diazepam

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per 5 mL Oral > Solid: 5 mg; 10 mg Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube Parenteral > General injections > unspecified: 5 mg per mL	Palliative care

Docusate sodium

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 50 mg per 5 mL oral liquid Oral > Solid: 100 mg capsule	Palliative care

Dolasetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron for Palliative care

Fentanyl

Section	Indications
Opioid analgesics Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour Oral > Solid: 100 to 800 µg sub-lingual tablet (as citrate); 200 to 1600 µg lozenge (as citrate) Oral > Other: 200 to 1200 µg buccal film (as citrate)	Chronic cancer pain Other specified chronic cancer related pain

Gabapentin

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs) Oral > Solid: 100 mg; 200 mg; 300 mg; 400 mg; 600 mg; 800 mg	Neuropathic pain

Granisetron

Section

Medicines for other common symptoms in palliative care

Indications

Therapeutic equivalent to [ondansetron](#) for

Palliative care

Haloperidol

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per mL

Oral > Solid: 0.5 mg; 2 mg; 5 mg

Parenteral > General injections > unspecified: 5 mg per mL in 1-mL ampoule

Indications

Palliative care

Hydromorphone

Section

Opioid analgesics

Indications

Therapeutic equivalent to [morphine](#) for

Pain

Hyoscine butylbromide

Section

Medicines for other common symptoms in palliative care

Parenteral > General injections > unspecified: 20 mg per mL injection

Indications

Palliative care

Hyoscine hydrobromide

Section

Medicines for other common symptoms in palliative care

Local > Topical > Transdermal patch: 1 mg per 72 hours transdermal patch

Parenteral > General injections > unspecified: 400 µg per mL injection; 600 µg per mL injection

Indications

Palliative care

Ibuprofen

Section

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral > Liquid: 200 mg per 5 mL; 100 mg per 5 mL

Oral > Solid > tablet: 200 mg; 400 mg; 600 mg

Indications

Pain

Lactulose

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 3.1 to 3.7 g per 5 mL oral liquid

Indications

Palliative care

Loperamide

Section

Medicines for other common symptoms in palliative care

Oral > Solid: 2 mg solid oral dosage form

Indications

Palliative care

Methadone

Section

Opioid analgesics

Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

Oral > Solid: 5 mg (methadone hydrochloride); 10 mg (methadone hydrochloride)

Indications

Chronic cancer pain

Metoclopramide

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 5 mg per 5 mL

Oral > Solid: 10 mg (hydrochloride)

Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)

Indications

Palliative care

Midazolam

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per mL oral liquid

Oral > Solid: 7.5 mg solid oral dosage form; 15 mg solid oral dosage form

Parenteral > General injections > unspecified: 1 mg per mL injection; 5 mg per mL injection

Indications

Palliative care

Morphine

Section	Indications
Opioid analgesics Oral > Liquid: 10 mg per 5 mL (morphine hydrochloride or morphine sulfate) Oral > Solid: 10 mg immediate release (morphine sulfate); 10 to 200 mg slow release (morphine hydrochloride or morphine sulfate) Oral > Other: 20 to 200 mg slow release granules to mix with water (morphine sulfate) Parenteral > General injections > unspecified: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)	Pain

Ondansetron

Section	Indications
Medicines for other common symptoms in palliative care Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride) Oral > Liquid: 4 mg per 5 mL (ondansetron base) Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base)	Palliative care

Oxycodone

Section	Indications
Opioid analgesics	Therapeutic equivalent to morphine for Pain

Palonosetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron for Palliative care

Paracetamol (acetaminophen)

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs) Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 100 mg; 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg	Pain

Senna

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 7.5 mg per 5 mL oral liquid

Indications

Palliative care

Tramadol

Section

Opioid analgesics

Oral > Liquid: 100 mg per mL (hydrochloride)

Oral > Solid: 50 mg (hydrochloride) immediate release; 50 mg (hydrochloride) controlled release; 100 mg (hydrochloride) controlled release; 150 mg (hydrochloride) controlled release; 200 mg (hydrochloride) controlled release; 300 mg (hydrochloride) controlled release; 400 mg (hydrochloride) controlled release

Parenteral > General injections > unspecified: 50 mg per mL in 2 mL ampoule (hydrochloride)

Indications

Chronic cancer pain

Tropisetron

Section

Medicines for other common symptoms in palliative care

Indications

Therapeutic equivalent to ondansetron for

Palliative care