

## Acetylcysteine

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule  
Oral > Liquid: 10%; 20%

### Indications

Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Acute or subacute hepatic failure

## Activated charcoal

### Section

Antidotes and other substances used in poisonings > Non-specific  
Oral > Other: powder

### Indications

Unspecified injury, poisoning or certain other consequences of external causes

## Atropine

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)

### Indications

Exposure to or harmful effects of undetermined intent of pesticides

## Calcium gluconate

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule

### Indications

Hypermagnesaemia

## Deferoxamine

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)

### Indications

Harmful effects of drugs, medicaments or biological substances, not elsewhere classified

## Dimercaprol

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Fomepizole

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule (sulfate); 1 g per mL in 1.5 mL ampoule (base)

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Ipecacuanha

### Section

Antidotes and other substances used in poisonings > Non-specific  
Oral > Liquid: 0.14% ipecacuanha alkaloids calculated as emetine

### Indications

Unspecified injury, poisoning or certain other consequences of external causes

## Methionine

### Section

Antidotes and other substances used in poisonings > Specific  
Oral > Solid: 250 mg (DL)

### Indications

Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

## Methylthioninium chloride

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule

### Indications

Acquired methaemoglobinaemia

## Naloxone

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)

### Indications

Unintentional exposure to or harmful effects of opioids or related analgesics

## Penicillamine

### Section

Antidotes and other substances used in poisonings > Specific  
Oral > Solid: 250 mg

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Potassium ferric hexacyanoferrate

### Section

Antidotes and other substances used in poisonings > Specific  
Oral > Solid: Powder for oral administration

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Pralidoxime

### Section

Antidotes and other substances used in poisonings > Specific

.

### Indications

Exposure to or harmful effects of undetermined intent of pesticides

## Sodium calcium edetate

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Sodium nitrite

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Sodium thiosulfate

### Section

Antidotes and other substances used in poisonings > Specific  
Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

## Succimer

### Section

Antidotes and other substances used in poisonings > Specific  
Oral > Solid: 100 mg

### Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified