

Acetylcysteine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule Oral > Liquid: 10%; 20%	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs Acute or subacute hepatic failure

Atropine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	Exposure to or harmful effects of undetermined intent of pesticides

Calcium gluconate

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Hypermagnesaemia

Deferoxamine

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)	Harmful effects of drugs, medicaments or biological substances, not elsewhere classified

Dimercaprol

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Fomepizole

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule (sulfate); 1 g per mL in 1.5 mL ampoule (base)	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Methionine

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg (DL)	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Methylthioninium chloride

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule	Acquired methaemoglobinaemia

Naloxone

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)	Unintentional exposure to or harmful effects of opioids or related analgesics

Penicillamine

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: 250 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Potassium ferric hexacyanoferrate

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: Powder for oral administration	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Pralidoxime

Section	Indications
Antidotes and other substances used in poisonings > Specific .	Exposure to or harmful effects of undetermined intent of pesticides

Sodium calcium edetate

Section	Indications
Antidotes and other substances used in poisonings > Specific Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium nitrite

Section

Antidotes and other substances used in poisonings > Specific
Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium thiosulfate

Section

Antidotes and other substances used in poisonings > Specific
Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Succimer

Section

Antidotes and other substances used in poisonings > Specific
Oral > Solid: 100 mg

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified