

Albendazole

| Section | Indications |
|---|--|
| Antifilarials Oral > Solid: 400 mg tablet (chewable) | <input type="text" value="Filariasis"/> |
| Intestinal anthelmintics Oral > Solid: 400 mg tablet (chewable) | <input type="text" value="Echinococcosis"/> <input type="text" value="Ascariasis"/> <input type="text" value="Enterobiasis"/> <input type="text" value="Trichuriasis"/> <input type="text" value="Hookworm diseases"/> <input type="text" value="Cysticercosis"/> |
| Cysticidal medicines Oral > Solid: 400 mg tablet (chewable) | <input type="text" value="Other specified echinococcosis"/> <input type="text" value="Cysticercosis of central nervous system"/> |

Diethylcarbamazine

| Section | Indications |
|---|---|
| Antifilarials Oral > Solid: 50 mg tablet (dihydrogen citrate); 100 mg tablet (dihydrogen citrate) | <input type="text" value="Lymphatic filariasis"/> |

Ivermectin

| Section | Indications |
|---|---|
| Intestinal anthelmintics Oral > Solid: 3 mg tablet (scored) | <input type="text" value="Strongyloidiasis"/> <input type="text" value="Ascariasis"/> <input type="text" value="Trichuriasis"/> <input type="text" value="Hookworm diseases"/> <input type="text" value="Ancylostomiasis"/> |
| Antifilarials Oral > Solid: 3 mg tablet (scored) | <input type="text" value="Onchocerciasis"/> <input type="text" value="Lymphatic filariasis"/> |

Levamisole

Section

Intestinal anthelmintics

Oral > Solid: 50 mg tablet; 150 mg tablet (as hydrochloride)

Indications

Ascariasis

Mebendazole

Section

Intestinal anthelmintics

Oral > Solid: 100 mg tablet (chewable); 500 mg tablet (chewable)

Indications

Ascariasis

Enterobiasis

Trichuriasis

Hookworm diseases

Cysticidal medicines

Oral > Solid: 500 mg tablet (chewable)

Other specified echinococcosis

Niclosamide

Section

Intestinal anthelmintics

Oral > Solid: 500 mg tablet (chewable)

Indications

Hymenolepiasis

Diphyllobothriasis

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

Oxamniquine

Section

Antischistosomes and other antitrematode medicines

Oral > Liquid: 250 mg per 5 mL

Oral > Solid: 250 mg capsule

Indications

Schistosomiasis due to *Schistosoma mansoni*

Praziquantel

Section

Intestinal anthelmintics

Oral > Solid: 600 mg tablet

Oral > Solid: 600 mg tablet; 150 mg tablet

Antischistosomes and other antitrematode medicines

Oral > Solid: 600 mg tablet

Cysticidal medicines

Oral > Solid: 600 mg tablet; 500 mg tablet

Indications

Diphyllobothriasis

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

Hymenolepiasis

Paragonimiasis

Clonorchiasis

Opisthorchiasis

Schistosomiasis

Cysticercosis of central nervous system

Pyrantel

Section

Intestinal anthelmintics

Oral > Liquid: 50 mg per mL (as embonate or pamoate)

Oral > Solid: 250 mg chewable tablet (as embonate or pamoate)

Indications

Ascariasis

Trichuriasis

Hookworm diseases

Enterobiasis

Suramin sodium

Section

Antifilarials

Parenteral > General injections > IV: 1 g in vial

Indications

Filariasis

Triclabendazole

Section

Antischistosomes and other antitrematode medicines

Oral > Solid: 250 mg tablet

Indications

Paragonimiasis

Fascioliasis