

## Albendazole

## Section

Intestinal anthelmintics

Oral &gt; Solid: 400 mg tablet (chewable)

## Indications

Echinococcosis

Enterobiasis

Trichuriasis

Ascariasis

Hookworm diseases

Cysticercosis

## Ivermectin

## Section

Intestinal anthelmintics

Oral &gt; Solid: 3 mg tablet (scored)

## Indications

Strongyloidiasis

Trichuriasis

Ascariasis

Hookworm diseases

Ancylostomiasis

## Levamisole

## Section

Intestinal anthelmintics

Oral &gt; Solid: 50 mg tablet; 150 mg tablet (as hydrochloride)

## Indications

Ascariasis

## Mebendazole

## Section

Intestinal anthelmintics

Oral &gt; Solid: 100 mg tablet (chewable); 500 mg tablet (chewable)

## Indications

Enterobiasis

Trichuriasis

Ascariasis

Hookworm diseases

## Nicosamide

### Section

Intestinal anthelmintics

Oral > Solid: 500 mg tablet (chewable)

### Indications

Hymenolepiasis

Diphyllobothriasis

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

## Praziquantel

### Section

Intestinal anthelmintics

Oral > Solid: 600 mg tablet; 150 mg tablet

Oral > Solid: 600 mg tablet

### Indications

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

Diphyllobothriasis

Hymenolepiasis

## Pyrantel

### Section

Intestinal anthelmintics

Oral > Liquid: 50 mg per mL (as embonate or pamoate)

Oral > Solid: 250 mg chewable tablet (as embonate or pamoate)

### Indications

Enterobiasis

Trichuriasis

Ascariasis

Hookworm diseases