

**Amikacin**

## Section

Antituberculosis medicines  
 Parenteral > General injections > unspecified: 100 mg per 2 mL in 2 mL vial (as sulfate); 250 mg per mL in 2 mL vial (as sulfate)

## Indications

Multi-drug resistant Mycobacterium tuberculosis

**Amoxicillin + clavulanic acid**

## Section

Antituberculosis medicines  
 Oral > Liquid: 250 mg (as trihydrate) + 62.5 mg per 5 mL (as potassium salt) powder for oral liquid  
 Oral > Solid: 500 mg (as trihydrate) + 125 mg (as potassium salt)

## Indications

Multi-drug resistant Mycobacterium tuberculosis

**Bedaquiline**

## Section

Antituberculosis medicines  
 Oral > Solid: 100 mg tablet; 20 mg tablet

## Indications

Multi-drug resistant Mycobacterium tuberculosis

**Capreomycin**

## Section

Antituberculosis medicines  
 Parenteral > General injections > IV: 1000 mg powder for injection (vial)

## Indications

Multi-drug resistant Mycobacterium tuberculosis

**Clofazimine**

## Section

Antituberculosis medicines  
 Oral > Solid: 100 mg; 50 mg

## Indications

Multi-drug resistant Mycobacterium tuberculosis

**Cycloserine**

## Section

Antituberculosis medicines  
 Oral > Solid: 250 mg; 125 mg

## Indications

Multi-drug resistant Mycobacterium tuberculosis

## Delamanid

### Section

Antituberculosis medicines

Oral > Solid: 50 mg tablet; 25 mg tablet (dispersible)

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Ethambutol

### Section

Antituberculosis medicines

Oral > Liquid: 25 mg per mL

Oral > Solid: 100 mg (hydrochloride); 400 mg (hydrochloride); 100 mg tablet (dispersible)

Parenteral > General injections > IV: 1000 mg per 10 mL; 2000 mg per 20 mL

### Indications

Tuberculosis

Other specified tuberculosis

## Ethambutol + isoniazid

### Section

Antituberculosis medicines

Oral > Solid: 400 mg + 150 mg tablet

### Indications

Tuberculosis

## Ethambutol + isoniazid + pyrazinamide + rifampicin

### Section

Antituberculosis medicines

Oral > Solid: 275 mg + 75 mg + 400 mg + 150 mg tablet

### Indications

Tuberculosis

## Ethambutol + isoniazid + rifampicin

### Section

Antituberculosis medicines

Oral > Solid: 275 mg + 75 mg + 150 mg tablet

### Indications

Tuberculosis

## Ethionamide

### Section

Antituberculosis medicines

Oral > Solid: 125 mg tablet; 250 mg tablet; 125 mg tablet (dispersible)

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Gatifloxacin

### Section

Antituberculosis medicines  
Oral > Solid: 200 mg; 400 mg

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Imipenem + cilastatin

### Section

Antituberculosis medicines

### Indications

Therapeutic equivalent to meropenem for

Multi-drug resistant Mycobacterium tuberculosis

## Isoniazid

### Section

Antituberculosis medicines  
Oral > Liquid: 50 mg per 5 mL  
Oral > Solid: 100 mg tablet; 100 mg tablet (dispersible); 300 mg tablet  
Parenteral > General injections > IV: 100 mg per mL

### Indications

Tuberculosis

Other specified tuberculosis

## Isoniazid + pyrazinamide + rifampicin

### Section

Antituberculosis medicines  
Oral > Solid: 50 mg + 150 mg + 75 mg tablet (dispersible)

### Indications

Tuberculosis

## Isoniazid + rifampicin

### Section

Antituberculosis medicines  
Oral > Solid: 50 mg + 75 mg tablet (dispersible); 75 mg + 150 mg tablet;  
150 mg + 300 mg tablet

### Indications

Tuberculosis

## Isoniazid + rifapentine

### Section

Antituberculosis medicines  
Oral > Solid: 300 mg + 300 mg tablet (scored)

### Indications

Latent tuberculosis

## Kanamycin (injection)

### Section

Antituberculosis medicines

Parenteral > General injections > IV: 1000 mg powder for injection (vial)

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Levofloxacin

### Section

Antituberculosis medicines

Oral > Solid: 250 mg tablet; 500 mg tablet; 750 mg tablet; 100 mg tablet (dispersible)

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Linezolid

### Section

Antituberculosis medicines

Oral > Liquid: 100 mg per 5 mL powder for oral liquid

Oral > Solid: 600 mg tablet; 150 mg tablet (dispersible)

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Meropenem

### Section

Antituberculosis medicines

Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Moxifloxacin

### Section

Antituberculosis medicines

Oral > Solid: 400 mg tablet; 100 mg tablet (dispersible)

Oral > Solid: 400 mg tablet

### Indications

Multi-drug resistant Mycobacterium tuberculosis

Tuberculosis

## Ofloxacin

### Section

Antituberculosis medicines

Oral > Solid: 200 mg; 400 mg

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## P-aminosalicylic acid

### Section

Antituberculosis medicines

Oral > Solid: 4 g granules in sachet

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Protionamide

### Section

Antituberculosis medicines

### Indications

Therapeutic equivalent to ethionamide for

Multi-drug resistant Mycobacterium tuberculosis

## Pyrazinamide

### Section

Antituberculosis medicines

Oral > Liquid: 30 mg per mL

Oral > Solid: 400 mg; 150 mg tablet (dispersible); 500 mg

### Indications

Tuberculosis

## Rifabutin

### Section

Antituberculosis medicines

Oral > Solid: 150 mg

### Indications

Tuberculosis

## Rifampicin

### Section

Antituberculosis medicines

Oral > Liquid: 20 mg per mL

Oral > Solid: 150 mg; 300 mg

Parenteral > General injections > IV: 600 mg in vial powder for injection

### Indications

Tuberculosis

Other specified tuberculosis

## Rifapentine

### Section

Antituberculosis medicines

Oral > Solid: 150 mg tablet; 300 mg tablet

### Indications

Tuberculosis

Latent tuberculosis

## Streptomycin (injection)

### Section

Antituberculosis medicines

Parenteral > General injections > IV: 1 g powder for injection (as sulfate)  
as vial

### Indications

Multi-drug resistant Mycobacterium tuberculosis

## Terizidone

### Section

Antituberculosis medicines

### Indications

Therapeutic equivalent to cycloserine for

Multi-drug resistant Mycobacterium tuberculosis