

Amphotericin b

Section	Indications
<p>Antifungal medicines</p> <p>Parenteral > General injections > IV: 50 mg powder for injection in vial (as deoxycholate or liposomal complex)</p>	<p>Blastomycosis</p> <p>Aspergillosis</p> <p>Coccidioidomycosis</p> <p>Candidosis</p> <p>Cryptococcosis</p> <p>Histoplasmosis</p> <p>Sporotrichosis</p>

Clotrimazole

Section	Indications
<p>Antifungal medicines</p> <p>Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet</p>	<p>Vulvovaginal candidosis</p>

Fluconazole

Section	Indications
<p>Antifungal medicines</p> <p>Parenteral > General injections > IV: 2 mg per mL in vial</p> <p>Oral > Liquid: 50 mg per 5 mL</p> <p>Oral > Solid: 50 mg capsule</p>	<p>Candidosis</p> <p>Cryptococcosis</p>

Flucytosine

Section	Indications
<p>Antifungal medicines</p> <p>Parenteral > General injections > IV: 2.5 g per 250 mL infusion</p> <p>Oral > Solid: 250 mg capsule</p>	<p>Cryptococcosis</p>

Griseofulvin

Section	Indications
<p>Antifungal medicines</p> <p>Oral > Liquid: 125 mg per 5 mL</p> <p>Oral > Solid: 125 mg; 250 mg</p>	<p>Dermatophytosis</p>

Itraconazole

Section

Antifungal medicines

Oral > Liquid: 10 mg per mL

Oral > Solid: 100 mg

Indications

Chronic pulmonary aspergillosis

Chromoblastomycosis

Penicilliosis

Sporotrichosis

Histoplasmosis

Paracoccidioidomycosis

Nystatin

Section

Antifungal medicines

Oral > Liquid: 100000 IU per mL; 50 mg per 5 mL

Oral > Solid: 500000 IU tablet; 100000 IU lozenge; 100000 IU tablet

Local > Topical > Other: 100000 IU pessary

Indications

Candidosis

Potassium iodide

Section

Antifungal medicines

Local > Topical > Solution: Saturated solution

Indications

Sporotrichosis

Voriconazole

Section

Antifungal medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection

Oral > Liquid: 40 mg per mL powder for oral liquid

Oral > Solid: 50 mg tablet; 200 mg tablet

Indications

Chronic pulmonary aspergillosis

Invasive aspergillosis