

## Amphotericin B

| Section   | Indications   |
|---|---|
| Antifungal medicines<br>Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate) | <input type="text" value="Blastomycosis"/><br><input type="text" value="Cryptococcosis"/><br><input type="text" value="Candidosis"/><br><input type="text" value="Coccidioidomycosis"/><br><input type="text" value="Histoplasmosis"/><br><input type="text" value="Sporotrichosis"/><br><input type="text" value="Aspergillosis"/> |

## Anidulafungin

| Section              | Indications  |
|----------------------|--|
| Antifungal medicines | Therapeutic equivalent to micafungin for<br><input type="text" value="Systemic or invasive candidosis"/> |

## Caspofungin

| Section              | Indications  |
|----------------------|--|
| Antifungal medicines | Therapeutic equivalent to micafungin for<br><input type="text" value="Systemic or invasive candidosis"/> |

## Clotrimazole

| Section  | Indications  |
|--|--|
| Antifungal medicines<br>Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet | <input type="text" value="Vulvovaginal candidosis"/> |

## Fluconazole

| Section  | Indications  |
|--|--|
| Antifungal medicines<br>Parenteral > General injections > IV: 2 mg per mL in vial<br>Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid<br>Oral > Solid > capsule: 50 mg | <input type="text" value="Cryptococcosis"/><br><input type="text" value="Candidosis"/> |

## Flucytosine

### Section

Antifungal medicines

Parenteral > General injections > IV: 2.5 g per 250 mL infusion

Oral > Solid: 250 mg capsule

### Indications

Cryptococcosis

## Griseofulvin

### Section

Antifungal medicines

Oral > Liquid: 125 mg per 5 mL

Oral > Solid: 125 mg; 250 mg

### Indications

Dermatophytosis

## Itraconazole

### Section

Antifungal medicines

Oral > Liquid: 10 mg per mL

Oral > Solid: 100 mg

### Indications

Chronic pulmonary aspergillosis

Chromoblastomycosis

Histoplasmosis

Talaromycosis

Sporotrichosis

Paracoccidioidomycosis

## Micafungin

### Section

Antifungal medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection

### Indications

Systemic or invasive candidosis

## Nystatin

### Section

Antifungal medicines

Oral > Liquid: 100000 IU per mL

Local > Topical > Other: 100000 IU pessary

Oral > Solid > dosage form: 500000 IU

Oral > Solid > lozenge: 100000 IU

### Indications

Candidosis

## Potassium iodide

### Section

Antifungal medicines

Local > Topical > Solution: Saturated solution

### Indications

Sporotrichosis

## Voriconazole

### Section

Antifungal medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection

Oral > Liquid: 40 mg per mL powder for oral liquid

Oral > Solid: 50 mg tablet; 200 mg tablet

### Indications

Chronic pulmonary aspergillosis

Invasive aspergillosis