

Amphotericin B

Section	Indications
<p>Antifungal medicines</p> <p>Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)</p>	<p>Blastomycosis</p> <p>Aspergillosis</p> <p>Cryptococcosis</p> <p>Candidosis</p> <p>Coccidioidomycosis</p> <p>Histoplasmosis</p> <p>Sporotrichosis</p>

Anidulafungin

Section	Indications
<p>Antifungal medicines</p>	<p>Therapeutic equivalent to micafungin for</p> <p>Systemic or invasive candidosis</p>

Caspofungin

Section	Indications
<p>Antifungal medicines</p>	<p>Therapeutic equivalent to micafungin for</p> <p>Systemic or invasive candidosis</p>

Clotrimazole

Section	Indications
<p>Antifungal medicines</p> <p>Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet</p>	<p>Vulvovaginal candidosis</p>

Fluconazole

Section	Indications
<p>Antifungal medicines</p> <p>Parenteral > General injections > IV: 2 mg per mL in vial</p> <p>Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid</p> <p>Oral > Solid > capsule: 50 mg</p>	<p>Cryptococcosis</p> <p>Candidosis</p>

Flucytosine

Section

Antifungal medicines

Parenteral > General injections > IV: 2.5 g per 250 mL infusion

Oral > Solid: 250 mg capsule

Indications

Cryptococcosis

Griseofulvin

Section

Antifungal medicines

Oral > Liquid: 125 mg per 5 mL

Oral > Solid: 125 mg; 250 mg

Indications

Dermatophytosis

Itraconazole

Section

Antifungal medicines

Oral > Liquid: 10 mg per mL

Oral > Solid: 100 mg

Indications

Chronic pulmonary aspergillosis

Chromoblastomycosis

Histoplasmosis

Talaromycosis

Sporotrichosis

Paracoccidioidomycosis

Micafungin

Section

Antifungal medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection

Indications

Systemic or invasive candidosis

Nystatin

Section

Antifungal medicines

Oral > Liquid: 100000 IU per mL

Local > Topical > Other: 100000 IU pessary

Oral > Solid > dosage form: 500000 IU

Oral > Solid > lozenge: 100000 IU

Indications

Candidosis

Potassium iodide

Section

Antifungal medicines

Local > Topical > Solution: Saturated solution

Indications

Sporotrichosis

Voriconazole

Section

Antifungal medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection

Oral > Liquid: 40 mg per mL powder for oral liquid

Oral > Solid: 50 mg tablet; 200 mg tablet

Indications

Chronic pulmonary aspergillosis

Invasive aspergillosis