

**Amodiaquine**

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)	<div>Malaria due to Plasmodium falciparum</div> <div>Malaria due to Plasmodium vivax</div>

**Amodiaquine + sulfadoxine + pyrimethamine**

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] dispersible tablets in co-package	<div>Malaria</div>

**Amphotericin B**

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	<div>Visceral leishmaniasis</div>

**Artemether**

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)	<div>Malaria due to Plasmodium falciparum</div> <div>Malaria due to Plasmodium vivax</div>

**Artemether + lumefantrine**

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet	<div>Malaria due to Plasmodium falciparum</div> <div>Malaria due to Plasmodium vivax</div>

## Artesunate

### Section

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution

Oral > Solid: 50 mg tablet

Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Artesunate + amodiaquine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Artesunate + mefloquine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet

### Indications

Malaria due to Plasmodium vivax

Malaria due to Plasmodium falciparum

## Artesunate + pyronaridine tetraphosphate

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Benznidazole

### Section

American trypanosomiasis

Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg

### Indications

Chagas disease

## Chloroquine

### Section

Antimalarial medicines > For chemoprevention

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate)

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium ovale

Malaria due to Plasmodium malariae

Malaria due to Plasmodium vivax

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Dihydroartemisinin + piperazine phosphate

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

## Diloxanide

### Section

Antiamoebic and anti giardiasis medicines

Oral > Solid: 500 mg (furoate)

### Indications

Amoebiasis

## Doxycycline

### Section

Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg (hydrochloride or hyclate)

Antimalarial medicines > For curative treatment

Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

## Eflornithine

### Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)

### Indications

African trypanosomiasis

## Fexinidazole

### Section

Medicines for the treatment of 1st stage African trypanosomiasis

Oral > Solid: 600 mg

Medicines for the treatment of 2nd stage African trypanosomiasis

Oral > Solid: 600 mg

### Indications

African trypanosomiasis

African trypanosomiasis

## Hydroxychloroquine

### Section

Antimalarial medicines > For chemoprevention

### Indications

Therapeutic equivalent to chloroquine for

Malaria due to Plasmodium falciparum

## Mefloquine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 250 mg tablet (hydrochloride)

Antimalarial medicines > For chemoprevention

Oral > Solid: 250 mg tablet (hydrochloride)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

Malaria due to Plasmodium falciparum

## Meglumine antimoniate

### Section

Antileishmaniasis medicines

Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Melarsoprol

### Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)

### Indications

African trypanosomiasis

## Metronidazole

### Section

#### Antiamoebic and anti giardiasis medicines

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Parenteral > General injections > unspecified: 500 mg in 100 mL vial

Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg

### Indications

Amoebiasis

## Miltefosine

### Section

#### Antileishmaniasis medicines

Oral > Solid: 10 mg; 50 mg

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Nifurtimox

### Section

#### Medicines for the treatment of 2nd stage African trypanosomiasis

Oral > Solid > tablet: 120 mg (scored); 30 mg (scored)

#### American trypanosomiasis

Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

### Indications

African trypanosomiasis

Chagas disease

## Paromomycin

### Section

#### Antileishmaniasis medicines

Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)

#### Antiamoebic and anti giardiasis medicines

Oral > Liquid: 125 mg per 5 mL as sulfate

Oral > Solid: 250 mg as sulfate

### Indications

Visceral leishmaniasis

Amoebiasis

## Pentamidine

### Section

#### Antileishmaniasis medicines

Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)

#### Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 200 mg tablet; 300 mg tablet

#### Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection

### Indications

Visceral leishmaniasis

Pneumocystosis

African trypanosomiasis

## Primaquine

### Section

#### Antimalarial medicines > For curative treatment

Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)

### Indications

Malaria due to Plasmodium ovale

Malaria due to Plasmodium vivax

## Proguanil

### Section

#### Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg tablet (hydrochloride)

### Indications

Malaria due to Plasmodium falciparum

## Pyrimethamine

### Section

#### Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 25 mg

### Indications

Toxoplasmosis

## Quinine

### Section

#### Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride)

Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)

### Indications

Malaria due to Plasmodium falciparum

## Sodium stibogluconate

### Section

Antileishmaniasis medicines

Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Sulfadiazine

### Section

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 500 mg

### Indications

Toxoplasmosis

## Sulfadoxine + pyrimethamine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 500 mg + 25 mg tablet

Antimalarial medicines > For chemoprevention

Oral > Solid: 500 mg + 25 mg tablet

Oral > Solid: 250 mg + 12.5 mg tablet

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

Malaria due to Plasmodium falciparum

Malaria

## Sulfamethoxazole + trimethoprim

### Section

Antipneumocystosis and antitoxoplasmosis medicines

Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule

Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid

Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet

Oral > Solid > dispersible tablet: 100 mg + 20 mg

### Indications

Pneumocystosis

## Suramin sodium

### Section

Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IV: 1 g in vial

### Indications

African trypanosomiasis

## Tinidazole

### Section

Antiamoebic and anti giardiasis medicines

### Indications

Therapeutic equivalent to metronidazole for

Amoebiasis