

Amodiaquine

Section

Antimalarial medicines > For curative treatment
 Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)

Indications

Malaria due to Plasmodium falciparum

Amodiaquine + sulfadoxine + pyrimethamine

Section

Antimalarial medicines > For chemoprevention
 Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1]
 dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg
 + 25 mg [1] dispersible tablets in co-package

Indications

Malaria

Amphotericin B

Section

Antileishmaniasis medicines
 Parenteral > General injections > IV: 50 mg powder for injection in vial (as
 deoxycholate or liposomal complex)

Indications

Visceral leishmaniasis

Artemether

Section

Antimalarial medicines > For curative treatment
 Parenteral > General injections > IV: 80 mg per mL in 1 mL ampoule (oily
 injection)

Indications

Malaria due to Plasmodium falciparum

Artemether + lumefantrine

Section

Antimalarial medicines > For curative treatment
 Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet

Indications

Malaria due to Plasmodium falciparum

Artesunate

Section

Antimalarial medicines > For curative treatment
 Parenteral > General injections > IV: 60 mg in ampoule containing
 anhydrous artesunic acid with a separate ampoule of 5% sodium
 bicarbonate solution
 Oral > Solid: 50 mg tablet
 Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules

Indications

Malaria due to Plasmodium falciparum

Artesunate + amodiaquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet

Indications

Malaria due to Plasmodium falciparum

Artesunate + mefloquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet

Indications

Malaria due to Plasmodium falciparum

Artesunate + pyronaridine tetraphosphate

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules

Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

Benznidazole

Section

American trypanosomiasis

Oral > Solid: 100 mg tablet; 50 mg tablet (scored); 12.5 mg tablet

Indications

Chagas disease

Chloroquine

Section

Antimalarial medicines > For chemoprevention

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate)

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)

Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium ovale

Malaria due to Plasmodium vivax

Malaria due to Plasmodium malariae

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

Dihydroartemisinin + piperazine phosphate

Section

Antimalarial medicines > For curative treatment
Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet

Indications

Malaria due to Plasmodium falciparum

Diloxanide

Section

Antiamoebic and anti giardiasis medicines
Oral > Solid: 500 mg (furoate)

Indications

Amoebiasis

Doxycycline

Section

Antimalarial medicines > For curative treatment
Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Antimalarial medicines > For chemoprevention
Oral > Solid: 100 mg (hydrochloride or hyclate)

Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

Eflornithine

Section

Medicines for the treatment of 2nd stage African trypanosomiasis
Parenteral > General injections > IV: 200 mg per mL in 100 mL bottle (hydrochloride)

Indications

African trypanosomiasis

Fexinidazole

Section

Medicines for the treatment of 1st stage African trypanosomiasis
Oral > Solid: 600 mg

Medicines for the treatment of 2nd stage African trypanosomiasis
Oral > Solid: 600 mg

Indications

African trypanosomiasis

African trypanosomiasis

Hydroxychloroquine

Section

Antimalarial medicines > For chemoprevention

Indications

Therapeutic equivalent to chloroquine for

Malaria due to Plasmodium falciparum

Mefloquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 250 mg tablet (hydrochloride)

Antimalarial medicines > For chemoprevention

Oral > Solid: 250 mg tablet (hydrochloride)

Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

Meglumine antimoniate

Section

Antileishmaniasis medicines

Parenteral > General injections > IM: 30% in 5 mL ampoule equivalent to approximately 8.1% antimony

Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

Melarsoprol

Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)

Indications

African trypanosomiasis

Metronidazole

Section

Antiamoebic and anti giardiasis medicines

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Oral > Solid: 200 to 500 mg tablet

Parenteral > General injections > unspecified: 500 mg in 100 mL vial

Indications

Amoebiasis

Miltefosine

Section

Antileishmaniasis medicines

Oral > Solid: 10 mg; 50 mg

Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

Nifurtimox

Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Oral > Solid: 120 mg

American trypanosomiasis

Oral > Solid: 30 mg tablet; 120 mg tablet; 250 mg tablet

Indications

African trypanosomiasis

Chagas disease

Paromomycin

Section

Antileishmaniasis medicines

Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)

Antiamoebic and anti giardiasis medicines

Oral > Liquid: 125 mg per 5 mL as sulfate

Oral > Solid: 250 mg as sulfate

Indications

Visceral leishmaniasis

Amoebiasis

Pentamidine

Section

Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IM: 200 mg (as isetionate) powder for injection

Antileishmaniasis medicines

Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 200 mg tablet; 300 mg tablet

Indications

African trypanosomiasis

Visceral leishmaniasis

Pneumocystosis

Primaquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)

Indications

Malaria due to Plasmodium ovale

Malaria due to Plasmodium vivax

Proguanil

Section

Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg tablet (hydrochloride)

Indications

Malaria due to Plasmodium falciparum

Pyrimethamine

Section

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 25 mg

Indications

Toxoplasmosis

Quinine

Section

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule
(quinine hydrochloride)

Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)

Indications

Malaria due to Plasmodium falciparum

Sodium stibogluconate

Section

Antileishmaniasis medicines

Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial

Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

Sulfadiazine

Section

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 500 mg

Indications

Toxoplasmosis

Sulfadoxine + pyrimethamine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 500 mg + 25 mg tablet

Antimalarial medicines > For chemoprevention

Oral > Solid: 500 mg + 25 mg tablet

Oral > Solid: 250 mg + 12.5 mg tablet

Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

Malaria

Sulfamethoxazole + trimethoprim

Section

Antipneumocystosis and antitoxoplasmosis medicines

Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule

Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid

Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet

Indications

Pneumocystosis

Suramin sodium

Section

Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IV: 1 g in vial

Indications

African trypanosomiasis

Tinidazole

Section

Antiamoebic and anti giardiasis medicines

Indications

Therapeutic equivalent to [metronidazole](#) for

Amoebiasis