

**Amodiaquine**

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)	<div>Malaria due to Plasmodium falciparum</div> <div>Malaria due to Plasmodium vivax</div>

**Amodiaquine + sulfadoxine + pyrimethamine**

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] dispersible tablets in co-package	<div>Malaria</div>

**Amphotericin B**

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	<div>Visceral leishmaniasis</div>

**Artemether**

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)	<div>Malaria due to Plasmodium falciparum</div> <div>Malaria due to Plasmodium vivax</div>

**Artemether + lumefantrine**

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet	<div>Malaria due to Plasmodium falciparum</div> <div>Malaria due to Plasmodium vivax</div>

## Artesunate

### Section

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution

Oral > Solid: 50 mg tablet

Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Artesunate + amodiaquine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Artesunate + mefloquine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Artesunate + pyronaridine tetraphosphate

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Benznidazole

### Section

American trypanosomiasis

Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg

### Indications

Chagas disease

## Chloroquine

### Section

Antimalarial medicines > For chemoprevention

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate)

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Malaria due to *Plasmodium ovale*

Malaria due to *Plasmodium malariae*

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Dihydroartemisinin + piperazine phosphate

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Diloxanide

### Section

Antiamoebic and anti giardiasis medicines

Oral > Solid: 500 mg (furoate)

### Indications

Amoebiasis

## Doxycycline

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg (hydrochloride or hyclate)

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium falciparum*

## Eflornithine

### Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)

### Indications

African trypanosomiasis

## Fexinidazole

### Section

Medicines for the treatment of 1st stage African trypanosomiasis  
Oral > Solid: 600 mg

Medicines for the treatment of 2nd stage African trypanosomiasis  
Oral > Solid: 600 mg

### Indications

African trypanosomiasis

African trypanosomiasis

## Hydroxychloroquine

### Section

Antimalarial medicines > For chemoprevention

### Indications

Therapeutic equivalent to chloroquine for

Malaria due to Plasmodium falciparum

## Mefloquine

### Section

Antimalarial medicines > For curative treatment  
Oral > Solid: 250 mg tablet (hydrochloride)

Antimalarial medicines > For chemoprevention  
Oral > Solid: 250 mg tablet (hydrochloride)

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

Malaria due to Plasmodium falciparum

## Meglumine antimoniate

### Section

Antileishmaniasis medicines  
Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule

### Indications

Visceral leishmaniasis

Cutaneous leishmaniasis

Mucocutaneous leishmaniasis

## Melarsoprol

### Section

Medicines for the treatment of 2nd stage African trypanosomiasis  
Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)

### Indications

African trypanosomiasis

## Metronidazole

### Section

#### Antiamoebic and anti giardiasis medicines

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Parenteral > General injections > unspecified: 500 mg in 100 mL vial

Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg

### Indications

Amoebiasis

## Miltefosine

### Section

#### Antileishmaniasis medicines

Oral > Solid: 10 mg; 50 mg

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Nifurtimox

### Section

#### American trypanosomiasis

Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

#### Medicines for the treatment of 2nd stage African trypanosomiasis

Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

### Indications

Chagas disease

African trypanosomiasis

## Paromomycin

### Section

#### Antileishmaniasis medicines

Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)

#### Antiamoebic and anti giardiasis medicines

Oral > Liquid: 125 mg per 5 mL as sulfate

Oral > Solid: 250 mg as sulfate

### Indications

Visceral leishmaniasis

Amoebiasis

## Pentamidine

### Section

#### Antileishmaniasis medicines

Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)

#### Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection

#### Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid > tablet: 200 mg (as isethionate); 300 mg (as isethionate)

### Indications

Visceral leishmaniasis

African trypanosomiasis

Pneumocystosis

## Primaquine

### Section

#### Antimalarial medicines > For curative treatment

Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)

### Indications

Malaria due to Plasmodium ovale

Malaria due to Plasmodium vivax

## Proguanil

### Section

#### Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg tablet (hydrochloride)

### Indications

Malaria due to Plasmodium falciparum

## Pyrimethamine

### Section

#### Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 25 mg

### Indications

Toxoplasmosis

## Quinine

### Section

#### Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride)

Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)

### Indications

Malaria due to Plasmodium falciparum

## Sodium stibogluconate

### Section

Antileishmaniasis medicines

Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial

### Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

## Sulfadiazine

### Section

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 500 mg

### Indications

Toxoplasmosis

## Sulfadoxine + pyrimethamine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 500 mg + 25 mg tablet

Antimalarial medicines > For chemoprevention

Oral > Solid: 500 mg + 25 mg tablet

Oral > Solid: 250 mg + 12.5 mg tablet

### Indications

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium vivax

Malaria due to Plasmodium falciparum

Malaria

## Sulfamethoxazole + trimethoprim

### Section

Antipneumocystosis and antitoxoplasmosis medicines

Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule

Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid

Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet

Oral > Solid > dispersible tablet: 100 mg + 20 mg

### Indications

Pneumocystosis

## Suramin sodium

### Section

Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IV: 1 g in vial

### Indications

African trypanosomiasis

## Tinidazole

### Section

Antiamoebic and anti giardiasis medicines

### Indications

Therapeutic equivalent to metronidazole for

Amoebiasis