

Amodiaquine

Section	Indications
Antimalarial medicines > For curative treatment	Malaria due to <i>Plasmodium falciparum</i>
Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)	Malaria due to <i>Plasmodium vivax</i>

Amodiaquine + sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For chemoprevention	Malaria
Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] dispersible tablets in co-package	

Artemether

Section	Indications
Antimalarial medicines > For curative treatment	Malaria due to <i>Plasmodium falciparum</i>
Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)	Malaria due to <i>Plasmodium vivax</i>

Artemether + lumefantrine

Section	Indications
Antimalarial medicines > For curative treatment	Malaria due to <i>Plasmodium falciparum</i>
Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet	Malaria due to <i>Plasmodium vivax</i>

Artesunate

Section	Indications
Antimalarial medicines > For curative treatment	Malaria due to <i>Plasmodium falciparum</i>
Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution	Malaria due to <i>Plasmodium vivax</i>
Oral > Solid: 50 mg tablet	
Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules	

Artesunate + amodiaquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + mefloquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + pyronaridine tetraphosphate

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Chloroquine

Section

Antimalarial medicines > For chemoprevention

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Malaria due to *Plasmodium ovale*

Malaria due to *Plasmodium malariae*

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)

Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)

Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Dihydroartemisinin + piperazine phosphate

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Doxycycline

Section

[Antimalarial medicines > For curative treatment](#)

Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

[Antimalarial medicines > For chemoprevention](#)

Oral > Solid: 100 mg (hydrochloride or hyclate)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium falciparum*

Hydroxychloroquine

Section

[Antimalarial medicines > For chemoprevention](#)

Indications

Therapeutic equivalent to [chloroquine](#) for

Malaria due to *Plasmodium falciparum*

Mefloquine

Section

[Antimalarial medicines > For curative treatment](#)

Oral > Solid: 250 mg tablet (hydrochloride)

[Antimalarial medicines > For chemoprevention](#)

Oral > Solid: 250 mg tablet (hydrochloride)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Malaria due to *Plasmodium falciparum*

Primaquine

Section

[Antimalarial medicines > For curative treatment](#)

Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)

Indications

Malaria due to *Plasmodium ovale*

Malaria due to *Plasmodium vivax*

Proguanil

Section

[Antimalarial medicines > For chemoprevention](#)

Oral > Solid: 100 mg tablet (hydrochloride)

Indications

Malaria due to *Plasmodium falciparum*

Quinine

Section

[Antimalarial medicines](#) > [For curative treatment](#)

Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride)

Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)

Indications

Malaria due to *Plasmodium falciparum*

Sulfadoxine + pyrimethamine

Section

[Antimalarial medicines](#) > [For curative treatment](#)

Oral > Solid: 500 mg + 25 mg tablet

[Antimalarial medicines](#) > [For chemoprevention](#)

Oral > Solid: 500 mg + 25 mg tablet

Oral > Solid: 250 mg + 12.5 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Malaria due to *Plasmodium falciparum*

Malaria