

**Amodiaquine**

## Section

Antimalarial medicines > For curative treatment  
 Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)

## Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

**Artemether**

## Section

Antimalarial medicines > For curative treatment  
 Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)

## Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

**Artemether + lumefantrine**

## Section

Antimalarial medicines > For curative treatment  
 Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet

## Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

**Artesunate**

## Section

Antimalarial medicines > For curative treatment  
 Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution  
 Oral > Solid: 50 mg tablet  
 Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules

## Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

**Artesunate + amodiaquine**

## Section

Antimalarial medicines > For curative treatment  
 Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet

## Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Artesunate + mefloquine

### Section

Antimalarial medicines > For curative treatment  
Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Artesunate + pyronaridine tetraphosphate

### Section

Antimalarial medicines > For curative treatment  
Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Chloroquine

### Section

Antimalarial medicines > For curative treatment  
Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate)  
Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate)  
Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Dihydroartemisinin + piperaquine phosphate

### Section

Antimalarial medicines > For curative treatment  
Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Doxycycline

### Section

Antimalarial medicines > For curative treatment  
Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

### Indications

Malaria due to *Plasmodium falciparum*

## Mefloquine

### Section

Antimalarial medicines > For curative treatment  
Oral > Solid: 250 mg tablet (hydrochloride)

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

## Primaquine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)

### Indications

Malaria due to *Plasmodium ovale*

Malaria due to *Plasmodium vivax*

## Quinine

### Section

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride)

Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)

### Indications

Malaria due to *Plasmodium falciparum*

## Sulfadoxine + pyrimethamine

### Section

Antimalarial medicines > For curative treatment

Oral > Solid: 500 mg + 25 mg tablet

### Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*