

Amodiaquine + sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] dispersible tablets in co-package	Malaria

Chloroquine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) Oral > Solid: 150 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax Malaria due to Plasmodium ovale Malaria due to Plasmodium malariae

Doxycycline

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 100 mg (hydrochloride or hyclate)	Malaria due to Plasmodium falciparum

Hydroxychloroquine

Section	Indications
Antimalarial medicines > For chemoprevention	Therapeutic equivalent to chloroquine for Malaria due to Plasmodium falciparum

Mefloquine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 250 mg tablet (hydrochloride)	Malaria due to Plasmodium falciparum

Proguanil

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 100 mg tablet (hydrochloride)	Malaria due to Plasmodium falciparum

Sulfadoxine + pyrimethamine

Section

[Antimalarial medicines](#) > For chemoprevention

Oral > Solid: 500 mg + 25 mg tablet

Oral > Solid: 250 mg + 12.5 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria