

Abiraterone

Section	Indications
Hormones and antihormones Oral > Solid: 250 mg; 500 mg	Malignant neoplasms of prostate

Adalimumab

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Axial spondyloarthritis Crohn disease site Rheumatoid arthritis

Afatinib

Section	Indications
Targeted therapies	Therapeutic equivalent to erlotinib for Other specified malignant neoplasms of bronchus or lung

All-trans retinoic acid

Section	Indications
Targeted therapies Oral > Solid: 10 mg capsule	Acute myeloid leukaemia with recurrent genetic abnormalities

Allopurinol

Section	Indications
Supportive medicines Oral > Solid: 100 mg; 300 mg	Tumour lysis syndrome

Anastrozole

Section	Indications
Hormones and antihormones Oral > Solid: 1 mg tablet	Other specified malignant neoplasms of breast Malignant neoplasms of breast

Arsenic trioxide

Section

Cytotoxic medicines

Parenteral > General injections > IV: 1 mg per mL concentrate for solution for infusion; 2 mg per mL concentrate for solution for infusion

Indications

Acute myeloid leukaemia with recurrent genetic abnormalities

Asparaginase

Section

Cytotoxic medicines

Powder for injection: 10 000 IU in vial.

Parenteral > General injections > unspecified: 10000 IU in vial powder for injection

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Lymphoid leukaemia, not elsewhere classified

Atezolizumab

Section

Immunomodulators

Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion

Indications

Other specified malignant neoplasms of bronchus or lung

Axicabtagene ciloleucel

Section

Antineoplastics and supportive medicines

Cell suspension for infusion: 2×10^6 anti-CD19 CAR-positive viable T-cells per kg of body weight.

Indications

Diffuse large B-cell lymphomas

Azacitidine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 100 mg in vial powder for injection

Indications

Myeloid leukaemia

Azathioprine

Section

Immunomodulators for non-malignant disease

Oral > Solid: 50 mg

Parenteral > General injections > IV: 100 mg in vial powder for injection (as sodium salt); 50 mg in vial powder for injection (as sodium salt)

Oral > Liquid: 10 mg per mL

Oral > Solid > tablet: 50 mg (scored); 25 mg

Indications

Relapsing-remitting multiple sclerosis

Failure or rejection of transplanted organs or tissues

Bendamustine

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL

Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Follicular lymphoma

Bicalutamide

Section

Hormones and antihormones

Oral > Solid: 50 mg

Indications

Malignant neoplasms of prostate

Binimetinib

Section

Targeted therapies

Indications

Therapeutic equivalent to trametinib for

Other specified melanoma of skin

Bleomycin

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 15 mg in vial (as sulfate)

Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Bortezomib

Section

Targeted therapies

Parenteral > General injections > unspecified: 3.5 mg in vial powder for injection

Indications

Plasma cell myeloma

Calcium folinate (leucovorin calcium)

Section	Indications
Cytotoxic medicines Oral > Solid: 15 mg tablet Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule Oral > Solid > tablet: 15 mg; 5 mg; 25 mg	Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasm metastasis in large intestine Malignant trophoblastic neoplasms of placenta Burkitt lymphoma including Burkitt leukaemia Osteosarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of rectum Malignant neoplasms of colon

Capecitabine

Section	Indications
Cytotoxic medicines Oral > Solid: 150 mg tablet; 500 mg tablet	Malignant neoplasms of colon Other specified malignant neoplasms of breast Malignant neoplasm metastasis in large intestine Malignant neoplasms of rectum

Carboplatin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL	Other specified gliomas of brain Other specified carcinomas of ovary Retinoblastoma Other specified malignant neoplasms of bronchus or lung Malignant neoplasms of nasopharynx Malignant neoplasms of kidney, except renal pelvis Malignant neoplasms of breast Osteosarcoma of bone and articular cartilage of unspecified sites Malignant neoplasms of lip, oral cavity or pharynx Malignant neoplasms of cervix uteri Other specified malignant neoplasms of the ovary Germ cell tumour of testis

Certolizumab pegol

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) for

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) for

Crohn disease site

Therapeutic equivalent to [adalimumab](#) for

Rheumatoid arthritis

Chlorambucil

Section

Cytotoxic medicines

Oral > Solid: 2 mg tablet

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Ciclosporin

Section

Immunomodulators for non-malignant disease

Parenteral > General injections > IV: 50 mg per mL in 1 mL ampoule concentrate for injection

Oral > Liquid: 100 mg per mL

Oral > Solid > capsule: 25 mg

Indications

Failure or rejection of transplanted organs or tissues

Cisplatin

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection

Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Germ cell tumour of testis

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

Cladribine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 1 mg per mL in 10 mL vial; 2 mg per mL in 5 mL vial

Indications

Langerhans cell histiocytosis

Cobimetinib

Section

Targeted therapies

Indications

Therapeutic equivalent to trametinib for

Other specified melanoma of skin

Crizotinib

Section

Targeted therapies

Oral > Solid: 200 mg; 250 mg

Oral > Solid > capsule: 200 mg; 250 mg

Indications

Other specified malignant neoplasms of bronchus or lung

Anaplastic large cell lymphoma, ALK-positive

Cyclophosphamide

Section	Indications
Cytotoxic medicines	Unspecified malignant neoplasms of ill-defined or unspecified sites
Oral > Solid: 25 mg tablet	Burkitt lymphoma including Burkitt leukaemia
Parenteral > General injections > unspecified: 500 mg in vial powder for injection	Malignant neoplasms of breast
Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection	Ewing sarcoma of bone and articular cartilage of unspecified sites
Oral > Solid: 25 mg; 50 mg	Lymphoid leukaemia, not elsewhere classified
Oral > Solid: 25 mg; 50 mg tablet	Chronic lymphocytic leukaemia or small lymphocytic lymphoma
	Diffuse large B-cell lymphomas
	Plasma cell myeloma
	Other specified gliomas of brain
	Other specified malignant neoplasms of breast
	Malignant trophoblastic neoplasms of placenta
	Follicular lymphoma
	Malignant neoplasms of kidney, except renal pelvis
	Rhabdomyosarcoma primary site
	Hodgkin lymphoma
	Anaplastic large cell lymphoma, ALK-negative
	Anaplastic large cell lymphoma, ALK-positive

Cytarabine

Section	Indications
Cytotoxic medicines	Unspecified malignant neoplasms of ill-defined or unspecified sites
Parenteral > General injections > unspecified: 100 mg in vial powder for injection	Lymphoid leukaemia, not elsewhere classified
Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial	Myeloid leukaemia
	Acute myeloid leukaemia with recurrent genetic abnormalities
	Burkitt lymphoma including Burkitt leukaemia
	Anaplastic large cell lymphoma, ALK-negative
	Anaplastic large cell lymphoma, ALK-positive
	Langerhans cell histiocytosis

Dabrafenib

Section	Indications
Targeted therapies Oral > Solid: 50 mg; 75 mg	Other specified melanoma of skin

Dacarbazine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection	Hodgkin lymphoma

Dactinomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 500 µg in vial powder for injection	Malignant trophoblastic neoplasms of placenta Unspecified malignant neoplasms of ill-defined or unspecified sites Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site Ewing sarcoma of bone and articular cartilage of unspecified sites

Daratumumab

Section	Indications
Immunomodulators Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL	Plasma cell myeloma

Dasatinib

Section	Indications
Targeted therapies Oral > Solid: 20 mg tablet; 50 mg tablet; 70 mg tablet; 80 mg tablet; 100 mg tablet; 140 mg tablet	Chronic myeloid leukaemia, not elsewhere classified

Daunorubicin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)

Indications

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Acute myeloid leukaemia with recurrent genetic abnormalities

Dexamethasone

Section

Hormones and antihormones

Oral > Solid: 4 mg

Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

Oral > Liquid: 2 mg per 5 mL

Oral > Solid: 2 mg; 4 mg

Indications

Plasma cell myeloma

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Docetaxel

Section

Cytotoxic medicines

Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL

Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

Malignant neoplasms of prostate

Unspecified malignant neoplasms of ill-defined or unspecified sites

Doxorubicin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride)

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride); 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Parenteral > General injections > IV: 50 mg in vial (hydrochloride) +; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride); 10 mg in vial (hydrochloride)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Rhabdomyosarcoma primary site

Follicular lymphoma

Burkitt lymphoma including Burkitt leukaemia

Ewing sarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of breast

Lymphoid leukaemia, not elsewhere classified

Other specified malignant neoplasms of breast

Diffuse large B-cell lymphomas

Plasma cell myeloma

Malignant neoplasms of kidney, except renal pelvis

Osteosarcoma of bone and articular cartilage of unspecified sites

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Encorafenib

Section

Targeted therapies

Indications

Therapeutic equivalent to dabrafenib for

Other specified melanoma of skin

Enzalutamide

Section

Hormones and antihormones

Oral > Solid: 40 mg

Indications

Therapeutic equivalent to abiraterone for

Malignant neoplasms of prostate

Malignant neoplasms of prostate

Erlotinib

Section

Targeted therapies

Oral > Solid: 100 mg; 150 mg

Indications

Other specified malignant neoplasms of bronchus or lung

Etanercept

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) for

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) for

Crohn disease site

Therapeutic equivalent to [adalimumab](#) for

Rheumatoid arthritis

Etoposide

Section

Cytotoxic medicines

Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule

Oral > Solid: 100 mg capsule

Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate)

Oral > Solid: 100 mg capsule; 50 mg capsule

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Ewing sarcoma of bone and articular cartilage of unspecified sites

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Germ cell tumour of testis

Malignant trophoblastic neoplasms of placenta

Malignant neoplasms of kidney, except renal pelvis

Osteosarcoma of bone and articular cartilage of unspecified sites

Hodgkin lymphoma

Retinoblastoma

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Everolimus

Section

Targeted therapies

Oral > Solid: 2.5 mg; 5 mg; 7.5 mg; 10 mg; 2 mg tablet (dispersible); 3 mg tablet (dispersible); 5 mg tablet (dispersible)

Indications

Other specified gliomas of brain

Filgrastim

Section

Immunomodulators

Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial

Indications

Acquired neutropaenia

Fingolimod

Section

Immunomodulators for non-malignant disease

Oral > Solid: 0.25 mg; 0.5 mg

Indications

Relapsing-remitting multiple sclerosis

Fludarabine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection

Oral > Solid: 10 mg tablet (fludarabine phosphate)

Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Fluorouracil

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule

Parenteral > General injections > IV: 50 mg per mL in vial

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Malignant neoplasms of nasopharynx

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of breast

Malignant neoplasms of rectum

Malignant neoplasms of colon

Flutamide

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [bicalutamide](#) for

Malignant neoplasms of prostate

Fulvestrant

Section

Hormones and antihormones

Parenteral > General injections > IV: 250 mg per 5 mL

Indications

Malignant neoplasms of breast

Gefitinib

Section

Targeted therapies

Indications

Therapeutic equivalent to [erlotinib](#) for

Other specified malignant neoplasms of bronchus or lung

Gemcitabine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection

Indications

Other specified carcinomas of ovary

Other specified malignant neoplasms of bronchus or lung

Golimumab

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) for

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) for

Crohn disease site

Therapeutic equivalent to [adalimumab](#) for

Rheumatoid arthritis

Goserelin

Section

Hormones and antihormones

Indications

Therapeutic equivalent to leuporelin for

Malignant neoplasms of breast

Therapeutic equivalent to leuporelin for

Malignant neoplasms of prostate

Hydrocortisone

Section

Hormones and antihormones

Parenteral > General injections > unspecified: 100 mg in vial (as sodium succinate)

Indications

Lymphoid leukaemia, not elsewhere classified

Unspecified malignant neoplasms of ill-defined or unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Hydroxycarbamide (hydroxyurea)

Section

Cytotoxic medicines

Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g

Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg

Indications

Squamous cell carcinoma of oropharynx

Chronic myeloid leukaemia, not elsewhere classified

Ibrutinib

Section

Targeted therapies

Oral > Solid: 140 mg

Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Ifosfamide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection	Other specified malignant neoplasms of the ovary Germ cell tumour of testis Ewing sarcoma of bone and articular cartilage of unspecified sites Osteosarcoma of bone and articular cartilage of unspecified sites Rhabdomyosarcoma primary site Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive

Imatinib

Section	Indications
Targeted therapies Oral > Solid: 100 mg; 400 mg	B lymphoblastic leukaemia or lymphoma with t(9:22) (q34;q11.2); BCR-ABL1 Gastrointestinal stromal tumour of unspecified gastrointestinal sites Chronic myeloid leukaemia, not elsewhere classified

Infliximab

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab for Axial spondyloarthritis Therapeutic equivalent to adalimumab for Crohn disease site Therapeutic equivalent to adalimumab for Rheumatoid arthritis

Irinotecan

Section

Cytotoxic medicines

Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial

Indications

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of kidney, except renal pelvis

Rhabdomyosarcoma primary site

Lenalidomide

Section

Immunomodulators

Oral > Solid: 25 mg

Indications

Plasma cell myeloma

Leuprorelin

Section

Hormones and antihormones

Parenteral > General injections > IM: 7.5 mg in prefilled syringe; 22.5 mg in prefilled syringe

Indications

Malignant neoplasms of breast

Malignant neoplasms of prostate

Lisocabtagene maraleucel

Section

Antineoplastics and supportive medicines

Indications

Therapeutic equivalent to axicabtagene ciloleucel for

Diffuse large B-cell lymphomas

Melphalan

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection

Oral > Solid: 2 mg

Indications

Plasma cell myeloma

Mercaptopurine

Section

Cytotoxic medicines

Oral > Solid: 50 mg tablet

Oral > Liquid: 20 mg per mL

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Acute myeloid leukaemia with recurrent genetic abnormalities

Lymphoid leukaemia, not elsewhere classified

Langerhans cell histiocytosis

Mesna

Section

Supportive medicines

Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule

Oral > Solid: 400 mg; 600 mg

Indications

Osteosarcoma of bone and articular cartilage of other specified sites

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Ewing sarcoma of bone and articular cartilage of unspecified sites

Rhabdomyosarcoma primary site

Unspecified malignant neoplasms of ill-defined or unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of kidney, except renal pelvis

Methotrexate

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium salt)

Oral > Solid: 2.5 mg (as sodium salt)

Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection

Immunomodulators for non-malignant disease

Parenteral > General injections > SC: pre-filled syringe in multiple strengths

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Lymphoid leukaemia, not elsewhere classified

Acute myeloid leukaemia with recurrent genetic abnormalities

Malignant trophoblastic neoplasms of placenta

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of breast

Osteosarcoma of bone and articular cartilage of unspecified sites

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Crohn disease site

Psoriatic arthritis

Methylprednisolone

Section

Hormones and antihormones

Parenteral > General injections > IV: 40 mg per mL in 1 mL single-dose vial (as sodium succinate); 40 mg per mL in 5 mL multi-dose vial (as sodium succinate); 80 mg per mL in 1 mL single-dose vial (as sodium succinate)

Indications

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

Nilotinib

Section

Targeted therapies

Oral > Solid: 150 mg capsule; 200 mg capsule

Indications

Chronic myeloid leukaemia, not elsewhere classified

Nilutamide

Section

Hormones and antihormones

Indications

Therapeutic equivalent to bicalutamide for

Malignant neoplasms of prostate

Nivolumab

Section

Immunomodulators

Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion

Indications

Other specified malignant neoplasms of bronchus or lung

Melanoma of skin

Oxaliplatin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial

Indications

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of colon

Paclitaxel

Section

Cytotoxic medicines

Parenteral > General injections > IV: 6 mg per mL

Indications

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Kaposi sarcoma of unspecified primary site

Other specified malignant neoplasms of breast

Other specified carcinomas of ovary

Malignant neoplasms of breast

Malignant neoplasms of cervix uteri

Pegaspargase

Section

Cytotoxic medicines

Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection

Indications

Lymphoid leukaemia, not elsewhere classified

Pembrolizumab

Section

Immunomodulators

Indications

Therapeutic equivalent to nivolumab for

Melanoma of skin

Pertuzumab

Section

Targeted therapies

Parenteral > General injections > IV: 240 mg per 14 mL in vial
concentrated solution

Indications

Carcinoma of breast, specialised type

Prednisolone

Section

Hormones and antihormones

Oral > Liquid: 5 mg per mL

Oral > Solid: 5 mg; 25 mg

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Plasma cell myeloma

Follicular lymphoma

Lymphoid leukaemia, not elsewhere classified

Malignant neoplasms of prostate

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Hodgkin lymphoma

Diffuse large B-cell lymphomas

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Prednisone

Section

Hormones and antihormones

Indications

Therapeutic equivalent to prednisolone for

Plasma cell myeloma

Therapeutic equivalent to prednisolone for

Follicular lymphoma

Therapeutic equivalent to prednisolone for

Lymphoid leukaemia, not elsewhere classified

Therapeutic equivalent to prednisolone for

Malignant neoplasms of prostate

Therapeutic equivalent to prednisolone for

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Therapeutic equivalent to prednisolone for

Hodgkin lymphoma

Therapeutic equivalent to prednisolone for

Diffuse large B-cell lymphomas

Therapeutic equivalent to prednisolone for

Anaplastic large cell lymphoma, ALK-negative

Therapeutic equivalent to prednisolone for

Anaplastic large cell lymphoma, ALK-positive

Therapeutic equivalent to prednisolone for

Langerhans cell histiocytosis

Procarbazine

Section

Cytotoxic medicines

Oral > Solid: 50 mg (as hydrochloride)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Hodgkin lymphoma

Rasburicase

Section

Supportive medicines

Parenteral > General injections > IV: 1.5 mg in vial powder and solvent for solution; 7.5 mg in vial powder and solvent for solution

Indications

Tumour lysis syndrome

Realgar-indigo naturalis formulation

Section

Cytotoxic medicines

Oral > Solid: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg)

Indications

Acute myeloid leukaemia with recurrent genetic abnormalities

Rituximab

Section

Targeted therapies

Parenteral > General injections > IV: 100 mg per 10 mL in 10 mL vial; 500 mg per 50 mL in 50 mL vial

Indications

Follicular lymphoma

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Diffuse large B-cell lymphomas

Tacrolimus

Section

Immunomodulators for non-malignant disease

Parenteral > General injections > IV: 5 mg per mL in 1 mL vial

Oral > Liquid: 0.2 mg granules for oral suspension; 1 mg granules for oral suspension

Oral > Solid: 0.5 mg (immediate-release); 0.75 mg (immediate-release); 1 mg (immediate-release); 2 mg (immediate-release); 5 mg (immediate-release)

Indications

Failure or rejection of transplanted organs or tissues

Tamoxifen

Section

Hormones and antihormones

Oral > Solid: 10 mg (as citrate); 20 mg (as citrate)

Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

Thalidomide

Section

Immunomodulators

Oral > Solid: 50 mg

Indications

Plasma cell myeloma

Tioguanine

Section

Cytotoxic medicines

Oral > Solid: 40 mg

Indications

Lymphoid leukaemia, not elsewhere classified

Tisagenlecleucel

Section

Antineoplastics and supportive medicines

Indications

Therapeutic equivalent to axicabtagene ciloleucel for

Diffuse large B-cell lymphomas

Tislelizumab

Section

Immunomodulators

Parenteral > General injections > IV: 100 mg per 10 mL

Indications

Malignant neoplasms of urinary tract

Hodgkin lymphoma

Trametinib

Section

Targeted therapies

Oral > Solid: 0.5 mg; 2 mg

Indications

Other specified melanoma of skin

Trastuzumab

Section

Targeted therapies

Parenteral > General injections > IV: 60 mg in vial powder for injection;
150 mg in vial powder for injection; 440 mg in vial powder for injection

Indications

Carcinoma of breast, specialised type

Trastuzumab emtansine

Section

Targeted therapies

Parenteral > General injections > IV: 100 mg in vial powder for injection;
160 mg in vial powder for injection

Indications

Carcinoma of breast, specialised type

Triptorelin

Section

Hormones and antihormones

Indications

Therapeutic equivalent to leuporelin for

Malignant neoplasms of breast

Therapeutic equivalent to leuporelin for

Malignant neoplasms of prostate

Vemurafenib

Section

Targeted therapies

Indications

Therapeutic equivalent to dabrafenib for

Other specified melanoma of skin

Vinblastine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate)

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Vincristine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate) Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Malignant trophoblastic neoplasms of placenta Follicular lymphoma Kaposi sarcoma of unspecified primary site Rhabdomyosarcoma primary site Hodgkin lymphoma Diffuse large B-cell lymphomas Retinoblastoma Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Langerhans cell histiocytosis

Vinorelbine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 50 mg per 5 mL in 5 mL vial Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial Oral > Solid: 20 mg; 30 mg; 80 mg	Other specified malignant neoplasms of bronchus or lung Other specified malignant neoplasms of breast Rhabdomyosarcoma primary site

Zanubrutinib

Section	Indications
Targeted therapies Oral > Solid: 80 mg	Mantle cell lymphoma

Zoledronic acid

Section	Indications
Supportive medicines Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle	Malignant neoplasm metastasis in bone or bone marrow

