

Apixaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran for Venous thromboembolism Therapeutic equivalent to dabigatran for Other specified atrial fibrillation

Cyanocobalamin

Section	Indications
Antianaemia medicines .	Anaemias or other erythrocyte disorders

Dabigatran

Section	Indications
Medicines affecting coagulation Oral > Solid: 110 mg; 150 mg	Venous thromboembolism Other specified atrial fibrillation

Dalteparin

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to enoxaparin for Acute ischaemic heart disease Therapeutic equivalent to enoxaparin for Venous thromboembolism

Darbepoetin alfa

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents for Anaemia due to chronic disease

Desmopressin

Section	Indications
Medicines affecting coagulation	Haemophilia A
Parenteral > General injections > IV: 4 µg per mL in 1 mL ampoule (acetate)	Von Willebrand disease
Local > Nasal > Spray: 10 µg per dose (acetate)	

Edoxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran for
	Venous thromboembolism
	Therapeutic equivalent to dabigatran for
	Other specified atrial fibrillation

Enoxaparin

Section	Indications
Medicines affecting coagulation	Acute ischaemic heart disease
Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 120 mg per 0.8 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule	Venous thromboembolism
Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 120 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule	

Epoetin alfa

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents for
	Anaemia due to chronic disease

Epoetin beta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents for Anaemia due to chronic disease

Epoetin theta

Section	Indications
Antianaemia medicines	Therapeutic equivalent to erythropoiesis-stimulating agents for Anaemia due to chronic disease

Erythropoiesis-stimulating agents

Section	Indications
Antianaemia medicines Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe	Anaemia due to chronic disease

Ferrous salt

Section	Indications
Antianaemia medicines Oral > Liquid: 25 mg per mL iron (as sulfate) Oral > Solid: 60 mg iron	Iron deficiency anaemia

Ferrous salt + folic acid

Section	Indications
Antianaemia medicines Oral > Solid: 60 mg iron + 2.8 mg	Iron deficiency anaemia

Folic acid

Section

Antianaemia medicines

Oral > Solid: 1 mg tablet; 5 mg tablet

Oral > Solid: 400 µg

Indications

Folate deficiency anaemia

Anencephaly or similar anomalies

Spina bifida

Cephalocele

Heparin sodium

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 1000 IU per mL in 5 mL ampoule;
20000 IU per mL in 5 mL ampoule; 5000 IU per mL in 5 mL ampoule

Indications

Acute ischaemic heart disease

Unstable angina

Venous thromboembolism

Hydroxocobalamin

Section

Antianaemia medicines

Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate)

Indications

Megaloblastic anaemia due to vitamin B12 deficiency

Hydroxycarbamide (hydroxyurea)

Section

Other medicines for haemoglobinopathies

Oral > Solid > dosage form: 200 mg; 500 mg; 1 g; 100 mg

Indications

Other specified sickle cell disorders or other haemoglobinopathies

Methoxy polyethylene glycol-epoetin beta

Section

Antianaemia medicines

Indications

Therapeutic equivalent to erythropoiesis-stimulating agents for

Anaemia due to chronic disease

Nadroparin

Section

Medicines affecting coagulation

Indications

Therapeutic equivalent to [enoxaparin](#) for

[Acute ischaemic heart disease](#)

Therapeutic equivalent to [enoxaparin](#) for

[Venous thromboembolism](#)

Phytomenadione

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule

Oral > Solid: 10 mg tablet

Indications

[Haemorrhagic disorder due to other specified circulating anticoagulants](#)

[Haemorrhagic or haematological disorders of fetus or newborn](#)

Protamine sulfate

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 10 mg per mL in 5 mL ampoule

Indications

[Haemorrhagic disorder due to other specified circulating anticoagulants](#)

Rivaroxaban

Section

Medicines affecting coagulation

Indications

Therapeutic equivalent to [dabigatran](#) for

[Venous thromboembolism](#)

Therapeutic equivalent to [dabigatran](#) for

[Other specified atrial fibrillation](#)

Tranexamic acid

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule

Indications

[Haemorrhage, not elsewhere classified](#)

Warfarin

Section

Medicines affecting coagulation

Oral > Solid: 5 mg (sodium salt); 1 mg (sodium salt); 2 mg (sodium salt);
0.5 mg (sodium salt)

Indications

Other specified diseases of arteries or arterioles

Cerebral ischaemic stroke due to embolic occlusion

Venous thromboembolism

Atrial fibrillation