

**Cyanocobalamin**

## Section

Antianaemia medicines

## Indications

Anaemias and other erythrocyte disorders

**Darbepoetin alfa**

## Section

Antianaemia medicines

## Indications

Therapeutic equivalent to erythropoiesis-stimulating agents for

Anaemia of chronic disease

**Epoetin alfa**

## Section

Antianaemia medicines

## Indications

Therapeutic equivalent to erythropoiesis-stimulating agents for

Anaemia of chronic disease

**Epoetin beta**

## Section

Antianaemia medicines

## Indications

Therapeutic equivalent to erythropoiesis-stimulating agents for

Anaemia of chronic disease

**Epoetin theta**

## Section

Antianaemia medicines

## Indications

Therapeutic equivalent to erythropoiesis-stimulating agents for

Anaemia of chronic disease

## Erythropoiesis-stimulating agents

### Section

#### Antianaemia medicines

Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe

### Indications

Anaemia of chronic disease

## Ferrous salt

### Section

#### Antianaemia medicines

Oral > Liquid: 25 mg per mL iron (as sulfate)

Oral > Solid: 60 mg iron

### Indications

Iron deficiency anaemia

## Ferrous salt + folic acid

### Section

#### Antianaemia medicines

Oral > Solid: 60 mg iron + 400 µg tablet

Oral > Solid: 60 mg iron + 2.8 mg

### Indications

Nutritional anaemias complicating pregnancy, childbirth and the puerperium

Iron deficiency anaemia

## Folic acid

### Section

#### Antianaemia medicines

Oral > Solid: 1 mg tablet; 5 mg tablet

Oral > Solid: 400 µg

### Indications

Folate deficiency anaemia

Anencephaly or similar anomalies

Spina bifida

Cephalocele

## Hydroxocobalamin

### Section

#### Antianaemia medicines

Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate)

### Indications

Megaloblastic anaemia due to vitamin B12 deficiency

## Methoxy polyethylene glycol-epoetin beta

### Section

Antianaemia medicines

### Indications

Therapeutic equivalent to erythropoiesis-stimulating agents for

Anaemia of chronic disease