



Model List of Essential Medicines

Anti-d immunoglobulin

| Section | Indications |
|--|---------------------------------------|
| Human immunoglobulins | Maternal care for red cell antibodies |
| Parenteral > General injections > IV: 250 µg in single-dose vial | Rh incompatibility reaction |
| Parenteral > General injections > IM: 250 µg in single-dose vial | |

Anti-rabies immunoglobulin

| Section | Indications |
|---|-------------|
| Human immunoglobulins | Rabies |
| Parenteral > General injections > IM: 150 IU per mL in vial | |
| Parenteral > Locoregional injections > Other: 150 IU per mL in vial | |

Anti-tetanus immunoglobulin

| Section | Indications |
|--|-------------|
| Human immunoglobulins | Tetanus |
| Parenteral > General injections > IM: 500 IU in vial | |

C1 esterase inhibitor

| Section | Indications |
|---|-----------------------|
| Plasma-derived medicines | Hereditary angioedema |
| Parenteral > General injections > IV: 500 IU in vial powder for injection | |

Dextran 40

| Section | Indications |
|--------------------|--------------|
| Plasma substitutes | Hypovolaemia |
| . | |

Dextran 70

| Section | Indications |
|--|--------------|
| Plasma substitutes | Hypovolaemia |
| Parenteral > General injections > IV: 6% injectable solution | |

Fresh-frozen plasma

| Section | Indications |
|---------------------------------------|--|
| Blood and blood components | Blood transfusion without reported diagnosis |
| Parenteral > General injections > IV: | |

Normal immunoglobulin

| Section | Indications |
|--|-----------------------------------|
| Human immunoglobulins | Primary immunodeficiencies |
| Parenteral > General injections > IV: 5% protein solution; 10% protein solution | Mucocutaneous lymph node syndrome |
| Parenteral > General injections > IM: 16% protein solution | Langerhans cell histiocytosis |
| Parenteral > General injections > SC: 15% protein solution; 16% protein solution | |

Platelets

| Section | Indications |
|---------------------------------------|--|
| Blood and blood components | Blood transfusion without reported diagnosis |
| Parenteral > General injections > IV: | |

Polygeline

| Section | Indications |
|--|--|
| Plasma substitutes | Therapeutic equivalent to dextran 70 for |
| Parenteral > General injections > IV: 3.5% injectable solution | Hypovolaemia |
| Parenteral > General injections > IV: 3.5% in solution | Hypovolaemia |
| | |

Red blood cells

| Section | Indications |
|---------------------------------------|--|
| Blood and blood components | Blood transfusion without reported diagnosis |
| Parenteral > General injections > IV: | |

Whole blood

| Section | Indications |
|---------------------------------------|--|
| Blood and blood components | Blood transfusion without reported diagnosis |
| Parenteral > General injections > IV: | |