



Model List of Essential Medicines

Anti-d immunoglobulin

Section	Indications
Human immunoglobulins	Maternal care for red cell antibodies
PARENTERAL > General injections > IV: 250 µg in single-dose vial	Rh incompatibility reaction
PARENTERAL > General injections > IM: 250 µg in single-dose vial	

Anti-rabies immunoglobulin

Section	Indications
Human immunoglobulins	Rabies
PARENTERAL > General injections > IM: 150 IU per mL in vial	
PARENTERAL > Locoregional injections > Other: 150 IU per mL in vial	

Anti-tetanus immunoglobulin

Section	Indications
Human immunoglobulins	Tetanus
PARENTERAL > General injections > IM: 500 IU in vial	

C1 esterase inhibitor

Section	Indications
Plasma-derived medicines	Hereditary angioedema
PARENTERAL > General injections > IV: 500 IU in vial powder for injection	

Coagulation factor IX

Section	Indications
Blood coagulation factors	Coagulation defects, purpura or other haemorrhagic or related conditions
PARENTERAL > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection	Haemophilia B

Coagulation factor IX complex

Section	Indications
Blood coagulation factors	Therapeutic equivalent to coagulation factor IX for Coagulation defects, purpura or other haemorrhagic or related conditions
	Therapeutic equivalent to coagulation factor IX for Haemophilia B

Coagulation factor VIII

Section	Indications
Blood coagulation factors Parenteral > General injections > IV: 500 IU in vial powder for injection; 250 IU in vial powder for injection; 1000 IU in vial powder for injection	Haemophilia A

Normal immunoglobulin

Section	Indications
Human immunoglobulins Parenteral > General injections > IV: 5% protein solution; 10% protein solution Parenteral > General injections > IM: 16% protein solution Parenteral > General injections > SC: 15% protein solution; 16% protein solution	Primary immunodeficiencies Mucocutaneous lymph node syndrome Langerhans cell histiocytosis