

## Anti-d immunoglobulin

| Section  | Indications                           |
|--|---------------------------------------|
| Human immunoglobulins  | Maternal care for red cell antibodies |
| PARENTERAL > General injections > IV: 250 µg in single-dose vial | Rh incompatibility reaction           |
| PARENTERAL > General injections > IM: 250 µg in single-dose vial |                                       |

## Anti-rabies immunoglobulin

| Section   | Indications |
|---|-------------|
| Human immunoglobulins   | Rabies      |
| PARENTERAL > General injections > IM: 150 IU per mL in vial         |             |
| PARENTERAL > Locoregional injections > Other: 150 IU per mL in vial |             |

## Anti-tetanus immunoglobulin

| Section  | Indications |
|--|-------------|
| Human immunoglobulins                                | Tetanus     |
| PARENTERAL > General injections > IM: 500 IU in vial |             |

## C1 esterase inhibitor

| Section   | Indications           |
|---|-----------------------|
| Plasma-derived medicines  | Hereditary angioedema |
| PARENTERAL > General injections > IV: 500 IU in vial powder for injection |                       |

## Coagulation factor IX

| Section   | Indications  |
|---|--|
| Blood coagulation factors   | Coagulation defects, purpura or other haemorrhagic or related conditions |
| PARENTERAL > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection | Haemophilia B  |

## Coagulation factor IX complex

| Section                   | Indications   |
|---------------------------|---|
| Blood coagulation factors | Therapeutic equivalent to coagulation factor IX pour Coagulation defects, purpura or other haemorrhagic or related conditions |
|                           | Therapeutic equivalent to coagulation factor IX pour Haemophilia B  |

## Coagulation factor VIII

| Section   | Indications   |
|---|---------------|
| Blood coagulation factors<br>Parenteral > General injections > IV: 500 IU in vial powder for injection; 250 IU in vial powder for injection; 1000 IU in vial powder for injection | Haemophilia A |

## Normal immunoglobulin

| Section  | Indications  |
|--|--|
| Human immunoglobulins<br>Parenteral > General injections > IV: 5% protein solution; 10% protein solution<br>Parenteral > General injections > IM: 16% protein solution<br>Parenteral > General injections > SC: 15% protein solution; 16% protein solution | Primary immunodeficiencies<br>Mucocutaneous lymph node syndrome<br>Langerhans cell histiocytosis |