

## Anti-d immunoglobulin

| Section                                                                                                                                                       | Indications                                                          |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|
| Human immunoglobulins<br>Parenteral > General injections > IV: 250 µg in single-dose vial<br>Parenteral > General injections > IM: 250 µg in single-dose vial | Maternal care for red cell antibodies<br>Rh incompatibility reaction |

## Anti-rabies immunoglobulin

| Section                                                                                                                                                     | Indications |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| Human immunoglobulins<br>Parenteral > General injections > IM: 150 IU per mL in vial<br>Parenteral > Locoregional injections > Other: 150 IU per mL in vial | Rabies      |

## Anti-tetanus immunoglobulin

| Section                                                                       | Indications |
|-------------------------------------------------------------------------------|-------------|
| Human immunoglobulins<br>Parenteral > General injections > IM: 500 IU in vial | Tetanus     |

## C1 esterase inhibitor

| Section                                                                                               | Indications           |
|-------------------------------------------------------------------------------------------------------|-----------------------|
| Plasma-derived medicines<br>Parenteral > General injections > IV: 500 IU in vial powder for injection | Hereditary angioedema |

## Normal immunoglobulin

| Section                                                                                                                                                                                                                                                    | Indications                                                                                      |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------|
| Human immunoglobulins<br>Parenteral > General injections > IV: 5% protein solution; 10% protein solution<br>Parenteral > General injections > IM: 16% protein solution<br>Parenteral > General injections > SC: 15% protein solution; 16% protein solution | Primary immunodeficiencies<br>Mucocutaneous lymph node syndrome<br>Langerhans cell histiocytosis |