

Acetylsalicylic acid

Section

Anti-platelet medicines
Oral > Solid: 100 mg

Indications

Acute myocardial infarction

Cerebral ischaemic stroke

Acetylsalicylic acid + atorvastatin + ramipril

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease
Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg

Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease
Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg

Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

Alteplase

Section

Thrombolytic medicines
Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection

Indications

Cerebral ischaemic stroke

Amiodarone

Section

Antiarrhythmic medicines
Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride)
Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)

Indications

Cardiac arrhythmia

Amlodipine

Section

Antihypertensive medicines

Oral > Solid: 5 mg (as maleate, mesylate or besylate)

Indications

Essential hypertension

Atenolol

Section

Antianginal medicines

Oral > Solid: 50 mg; 100 mg

Antihypertensive medicines

Oral > Solid: 50 mg; 100 mg

Indications

Angina pectoris

Essential hypertension

Therapeutic equivalent to *bisoprolol* pour

Essential hypertension

Atorvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to *simvastatin* pour

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* pour

Coronary atherosclerosis

Atorvastatin + perindopril + amlodipine

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease

Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg

Indications

Chronic ischaemic heart disease

Atherosclerotic chronic arterial occlusive disease

Cerebral ischaemic stroke

Bisoprolol

Section

Antianginal medicines

Oral > Solid: 1.25 mg; 5 mg

Medicines used in heart failure

Oral > Solid: 1.25 mg; 5 mg

Antihypertensive medicines

Oral > Solid: 1.25 mg; 5 mg

Antiarrhythmic medicines

Oral > Solid: 1.25 mg; 5 mg

Indications

Angina pectoris

Heart failure

Essential hypertension

Cardiac arrhythmia

Bumetanide

Section

Medicines used in heart failure

Indications

Therapeutic equivalent to [furosemide](#) pour

Heart failure

Captopril

Section

Medicines used in heart failure

Oral > Solid: 25 mg

Indications

Heart failure

Carvedilol

Section

Antianginal medicines

Medicines used in heart failure

Antihypertensive medicines

Antiarrhythmic medicines

Indications

Therapeutic equivalent to [bisoprolol](#) pour

Angina pectoris

Therapeutic equivalent to [bisoprolol](#) pour

Heart failure

Therapeutic equivalent to [bisoprolol](#) pour

Essential hypertension

Therapeutic equivalent to [bisoprolol](#) pour

Cardiac arrhythmia

Chlorothiazide

Section

Antihypertensive medicines

Medicines used in heart failure

Indications

Therapeutic equivalent to hydrochlorothiazide pour

Essential hypertension

Therapeutic equivalent to hydrochlorothiazide pour

Heart failure

Chlortalidone

Section

Antihypertensive medicines

Medicines used in heart failure

Indications

Therapeutic equivalent to hydrochlorothiazide pour

Essential hypertension

Therapeutic equivalent to hydrochlorothiazide pour

Heart failure

Clopidogrel

Section

Anti-platelet medicines

Oral > Solid: 75 mg; 300 mg

Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

Digitoxin

Section

Medicines used in heart failure

Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule

Oral > Solid: 50 µg; 100 µg

Indications

Heart failure

Digoxin

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid: 62.5 µg; 250 µg

Medicines used in heart failure

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg

Indications

Cardiac arrhythmia

Heart failure

Dopamine

Section

Medicines used in heart failure

Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)

Indications

Heart failure

Cardiogenic shock

Enalapril

Section

Antihypertensive medicines

Oral > Liquid: 1 mg per mL (as hydrogen maleate)

Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)

Medicines used in heart failure

Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)

Indications

Essential hypertension

Heart failure

Epinephrine

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)

Indications

Cardiac arrest

Fluvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to *simvastatin* pour

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* pour

Coronary atherosclerosis

Furosemide

Section

Medicines used in heart failure

Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL

Oral > Solid > tablet: 40 mg; 20 mg

Indications

Heart failure

Glyceryl trinitrate

Section

Antianginal medicines

Oral > Solid: 500 µg sublingual tablet

Indications

Angina pectoris

Hydralazine

Section

Antihypertensive medicines

Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride)

Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride)

Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)

Indications

Essential hypertension

Gestational hypertension

Hydrochlorothiazide

Section

Antihypertensive medicines

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg; 12.5 mg

Medicines used in heart failure

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg

Indications

Essential hypertension

Heart failure

Indapamide

Section

Antihypertensive medicines

Medicines used in heart failure

Indications

Therapeutic equivalent to hydrochlorothiazide pour

Essential hypertension

Therapeutic equivalent to hydrochlorothiazide pour

Heart failure

Isoprenaline

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride)

Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)

Indications

Bradycardia

Cardiogenic shock

Isosorbide dinitrate

Section

Antianginal medicines

Oral > Solid: 5 mg sublingual tablet

Indications

Angina pectoris

Lidocaine

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)

Indications

Ventricular tachyarrhythmia

Lisinopril + amlodipine

Section

Antihypertensive medicines

Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg

Indications

Essential hypertension

Lisinopril + hydrochlorothiazide

Section

Antihypertensive medicines

Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg

Indications

Essential hypertension

Losartan

Section

Antihypertensive medicines

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Medicines used in heart failure

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Indications

Essential hypertension

Heart failure

Lovastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to *simvastatin* pour

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* pour

Coronary atherosclerosis

Methyldopa

Section

Antihypertensive medicines

Oral > Solid: 250 mg

Indications

Essential hypertension

Gestational hypertension

Metoprolol

Section

Antianginal medicines

Therapeutic equivalent to *bisoprolol* pour

Angina pectoris

Medicines used in heart failure

Therapeutic equivalent to *bisoprolol* pour

Heart failure

Antihypertensive medicines

Therapeutic equivalent to *bisoprolol* pour

Essential hypertension

Antiarrhythmic medicines

Therapeutic equivalent to *bisoprolol* pour

Cardiac arrhythmia

Nifedipine

Section

Antianginal medicines

Oral > Solid: 10 mg

Antihypertensive medicines

Oral > Solid: 10 mg tablet; sustained-release formulations

Indications

Angina pectoris

Essential hypertension

Pravastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to *simvastatin* pour

Mixed hyperlipidaemia

Therapeutic equivalent to *simvastatin* pour

Coronary atherosclerosis

Procainamide

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride)

Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)

Indications

Cardiac arrhythmia

Propranolol

Section

Antihypertensive medicines

Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

Antianginal medicines

Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)

Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

Antiarrhythmic medicines

Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride)

Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)

Indications

Essential hypertension

Angina pectoris

Cardiac arrhythmia

Quinidine

Section

Antiarrhythmic medicines
Oral > Solid: 200 mg (sulfate)

Indications

Cardiac arrhythmia

Simvastatin

Section

Lipid-lowering agents
Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg

Indications

Mixed hyperlipidaemia

Coronary atherosclerosis

Sodium nitroprusside

Section

Antihypertensive medicines
Parenteral > General injections > IV: 50 mg in ampoule powder for infusion

Indications

Hypertensive crisis

Spironolactone

Section

Medicines used in heart failure
Oral > Solid > tablet: 25 mg

Indications

Heart failure

Streptokinase

Section

Thrombolytic medicines
Parenteral > General injections > IV: 1.5 million IU in vial powder for injection

Indications

Acute myocardial infarction

Telmisartan + amlodipine

Section

Antihypertensive medicines
Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg

Indications

Essential hypertension

Telmisartan + hydrochlorothiazide

Section

Antihypertensive medicines
Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg

Indications

Essential hypertension

Ticagrelor

Section

Anti-platelet medicines

Oral > Solid > tablet: 60 mg; 90 mg

Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

Torsemide

Section

Medicines used in heart failure

Indications

Therapeutic equivalent to furosemide pour

Heart failure

Verapamil

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Antianginal medicines

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Indications

Supraventricular tachyarrhythmia

Angina pectoris