

Acetylsalicylic acid

Section

Anti-platelet medicines

Oral > Solid: 100 mg

Indications

Acute myocardial infarction

Cerebral ischaemic stroke

Acetylsalicylic acid + atorvastatin + ramipril

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease

Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg

Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease

Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg

Indications

Atherosclerotic chronic arterial occlusive disease

Chronic ischaemic heart disease

Cerebral ischaemic stroke

Alteplase

Section

Thrombolytic medicines

Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection

Indications

Cerebral ischaemic stroke

Amiodarone

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride)

Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)

Indications

Cardiac arrhythmia

Amlodipine

Section

Antihypertensive medicines

Oral > Solid: 5 mg (as maleate, mesylate or besylate)

Indications

Essential hypertension

Atenolol

Section

Antianginal medicines

Oral > Solid: 50 mg; 100 mg

Antihypertensive medicines

Oral > Solid: 50 mg; 100 mg

Indications

Angina pectoris

Essential hypertension

Therapeutic equivalent to [bisoprolol](#) pour

Essential hypertension

Atorvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Atorvastatin + perindopril + amlodipine

Section

Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease

Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg

Indications

Chronic ischaemic heart disease

Cerebral ischaemic stroke

Atherosclerotic chronic arterial occlusive disease

Bisoprolol

Section	Indications
Antianginal medicines Oral > Solid: 1.25 mg; 5 mg	Angina pectoris
Medicines used in heart failure Oral > Solid: 1.25 mg; 5 mg	Heart failure
Antihypertensive medicines Oral > Solid: 1.25 mg; 5 mg	Essential hypertension
Antiarrhythmic medicines Oral > Solid: 1.25 mg; 5 mg	Cardiac arrhythmia

Bumetanide

Section	Indications
Medicines used in heart failure	Therapeutic equivalent to furosemide pour Heart failure

Captopril

Section	Indications
Medicines used in heart failure Oral > Solid: 25 mg	Heart failure

Carvedilol

Section	Indications
Antianginal medicines	Therapeutic equivalent to bisoprolol pour Angina pectoris
Medicines used in heart failure	Therapeutic equivalent to bisoprolol pour Heart failure
Antihypertensive medicines	Therapeutic equivalent to bisoprolol pour Essential hypertension
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol pour Cardiac arrhythmia

Chlorothiazide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chlortalidone

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Clopidogrel

Section	Indications
Anti-platelet medicines Oral > Solid: 75 mg; 300 mg	Acute ischaemic heart disease Presence of coronary angioplasty implant or graft

Digitoxin

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule Oral > Solid: 50 µg; 100 µg	Heart failure

Digoxin

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid: 62.5 µg; 250 µg	Cardiac arrhythmia
Medicines used in heart failure Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule Oral > Liquid: 50 µg per mL Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg	Heart failure

Dopamine

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)	Heart failure Cardiogenic shock

Enalapril

Section	Indications
Antihypertensive medicines Oral > Liquid: 1 mg per mL (as hydrogen maleate) Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Essential hypertension
Medicines used in heart failure Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)	Heart failure

Epinephrine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)	Cardiac arrest

Fluvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Furosemide

Section

Medicines used in heart failure

Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL

Oral > Solid > tablet: 40 mg; 20 mg

Indications

Heart failure

Glyceryl trinitrate

Section

Antianginal medicines

Oral > Solid: 500 µg sublingual tablet

Indications

Angina pectoris

Hydralazine

Section

Antihypertensive medicines

Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride)

Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride)

Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)

Indications

Essential hypertension

Gestational hypertension

Hydrochlorothiazide

Section

Antihypertensive medicines

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg; 12.5 mg

Medicines used in heart failure

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg

Indications

Essential hypertension

Heart failure

Indapamide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Isoprenaline

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride) Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)	Bradycardia Cardiogenic shock

Isosorbide dinitrate

Section	Indications
Antianginal medicines Oral > Solid: 5 mg sublingual tablet	Angina pectoris

Lidocaine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)	Ventricular tachyarrhythmia

Lisinopril + amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg	Essential hypertension

Lisinopril + hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg	Essential hypertension

Losartan

Section

Antihypertensive medicines

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Medicines used in heart failure

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Indications

Essential hypertension

Heart failure

Lovastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Methyldopa

Section

Antihypertensive medicines

Oral > Solid: 250 mg

Indications

Essential hypertension

Gestational hypertension

Metoprolol

Section

Antianginal medicines

Therapeutic equivalent to [bisoprolol](#) pour

Angina pectoris

Medicines used in heart failure

Therapeutic equivalent to [bisoprolol](#) pour

Heart failure

Antihypertensive medicines

Therapeutic equivalent to [bisoprolol](#) pour

Essential hypertension

Antiarrhythmic medicines

Therapeutic equivalent to [bisoprolol](#) pour

Cardiac arrhythmia

Nifedipine

Section	Indications
Antianginal medicines Oral > Solid: 10 mg	Angina pectoris
Antihypertensive medicines Oral > Solid: 10 mg tablet; sustained-release formulations	Essential hypertension

Pravastatin

Section	Indications
Lipid-lowering agents	Therapeutic equivalent to simvastatin pour Mixed hyperlipidaemia Therapeutic equivalent to simvastatin pour Coronary atherosclerosis

Procainamide

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride) Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)	Cardiac arrhythmia

Propranolol

Section	Indications
Antihypertensive medicines Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Essential hypertension
Antianginal medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride) Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Angina pectoris
Antiarrhythmic medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride) Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Cardiac arrhythmia

Quinidine

Section

Antiarrhythmic medicines

Oral > Solid: 200 mg (sulfate)

Indications

Cardiac arrhythmia

Simvastatin

Section

Lipid-lowering agents

Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg

Indications

Mixed hyperlipidaemia

Coronary atherosclerosis

Sodium nitroprusside

Section

Antihypertensive medicines

Parenteral > General injections > IV: 50 mg in ampoule powder for infusion

Indications

Hypertensive crisis

Spirolactone

Section

Medicines used in heart failure

Oral > Solid: 25 mg

Indications

Heart failure

Streptokinase

Section

Thrombolytic medicines

Parenteral > General injections > IV: 1.5 million IU in vial powder for injection

Indications

Acute myocardial infarction

Telmisartan + amlodipine

Section

Antihypertensive medicines

Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg

Indications

Essential hypertension

Telmisartan + hydrochlorothiazide

Section

Antihypertensive medicines

Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg

Indications

Essential hypertension

Ticagrelor

Section

Anti-platelet medicines

Oral > Solid > tablet: 60 mg; 90 mg

Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

Torsemide

Section

Medicines used in heart failure

Indications

Therapeutic equivalent to furosemide pour

Heart failure

Verapamil

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Antianginal medicines

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Indications

Supraventricular tachyarrhythmia

Angina pectoris