

**Acamprosate calcium**

## Section

Medicines for alcohol use disorders  
Oral > Solid > tablet: 333 mg

## Indications

Disorders due to use of alcohol

**Buprenorphine**

## Section

Medicines for opioid use disorders

## Indications

Therapeutic equivalent to **methadone** for

Opioid dependence

**Bupropion**

## Section

Medicines for nicotine use disorders  
Oral > Solid: 150 mg sustained-release (hydrochloride)

## Indications

Nicotine dependence

**Methadone**

## Section

Medicines for opioid use disorders  
Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

## Indications

Opioid dependence

**Naltrexone**

## Section

Medicines for alcohol use disorders  
Parenteral > General injections > IM: 380 mg in vial (extended-release)  
Oral > Solid > tablet: 50 mg

## Indications

Disorders due to use of alcohol

**Nicotine replacement therapy**

## Section

Medicines for nicotine use disorders  
Oral > Other: 2 mg chewing gum; 4 mg chewing gum; 2 mg lozenge; 4 mg lozenge; 1 mg per actuation oral spray  
Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour

## Indications

Nicotine dependence

## Varenicline

### Section

Medicines for nicotine use disorders

Oral > Solid: 0.5 mg; 1 mg

### Indications

Nicotine dependence