

**Acetylsalicylic acid**

## Section

Juvenile joint diseases

Oral &gt; Solid: 100 to 500 mg

Local &gt; Rectal &gt; Suppository: 50 to 150 mg

## Indications

Acute rheumatic fever without mention of heart involvement

Mucocutaneous lymph node syndrome

Juvenile idiopathic arthritis

**Allopurinol**

## Section

Medicines used to treat gout

Oral &gt; Solid: 100 mg

## Indications

Gout

**Azathioprine**

## Section

Disease-modifying agents used in rheumatoid disorders (DMARDs)

Oral &gt; Solid: 50 mg

## Indications

Psoriatic arthritis

Rheumatoid arthritis

**Chloroquine**

## Section

Disease-modifying agents used in rheumatoid disorders (DMARDs)

Oral &gt; Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)

## Indications

Rheumatoid arthritis

**Colchicine**

## Section

Medicines used to treat gout

Oral &gt; Solid: 0.5 mg

## Indications

Gout

**Hydroxychloroquine**

## Section

Disease-modifying agents used in rheumatoid disorders (DMARDs)

Oral &gt; Solid: 200 mg (as sulfate)

## Indications

Systemic lupus erythematosus

## Methotrexate

### Section

Disease-modifying agents used in rheumatoid disorders (DMARDs)  
Oral > Solid: 2.5 mg (as sodium salt)

### Indications

Rheumatoid arthritis

## Penicillamine

### Section

Disease-modifying agents used in rheumatoid disorders (DMARDs)  
Oral > Solid: 250 mg

### Indications

Rheumatoid arthritis

## Probenecid

### Section

Medicines used to treat gout  
Oral > Solid: 500 mg

### Indications

Gout

## Sulfasalazine

### Section

Disease-modifying agents used in rheumatoid disorders (DMARDs)  
Oral > Solid: 500 mg

### Indications

Rheumatoid arthritis