

**Acetylsalicylic acid**

| Section                                                                                                            | Indications                                                                                                                                                 |
|--------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Medicines for juvenile joint diseases<br>Oral > Solid: 100 to 500 mg<br>Local > Rectal > Suppository: 50 to 150 mg | <div>Acute rheumatic fever without mention of heart involvement</div> <div>Mucocutaneous lymph node syndrome</div> <div>Juvenile idiopathic arthritis</div> |

**Adalimumab**

| Section                                                                                                                                               | Indications                              |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------|
| Medicines for juvenile joint diseases<br>Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL | <div>Juvenile idiopathic arthritis</div> |

**Allopurinol**

| Section                                              | Indications     |
|------------------------------------------------------|-----------------|
| Medicines used to treat gout<br>Oral > Solid: 100 mg | <div>Gout</div> |

**Azathioprine**

| Section                                                                | Indications                                                                          |
|------------------------------------------------------------------------|--------------------------------------------------------------------------------------|
| Disease-modifying anti-rheumatic drugs (DMARDs)<br>Oral > Solid: 50 mg | <div>Rheumatoid arthritis, serology unspecified</div> <div>Psoriatic arthritis</div> |

**Certolizumab pegol**

| Section                               | Indications                                                                                          |
|---------------------------------------|------------------------------------------------------------------------------------------------------|
| Medicines for juvenile joint diseases | Therapeutic equivalent to <a href="#">adalimumab</a> for<br><div>Juvenile idiopathic arthritis</div> |

**Chloroquine**

| Section                                                                                                                                           | Indications                     |
|---------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------|
| Disease-modifying anti-rheumatic drugs (DMARDs)<br>Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate) | <div>Rheumatoid arthritis</div> |

## Colchicine

### Section

Medicines used to treat gout

Oral > Solid: 0.5 mg

### Indications

Gout

## Etanercept

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

## Golimumab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

## Hydroxychloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 200 mg (as sulfate)

### Indications

Systemic lupus erythematosus

Cutaneous lupus erythematosus of unspecified type

## Infliximab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

## Methotrexate

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Medicines for juvenile joint diseases

Oral > Solid > tablet: 2.5 mg (as sodium salt)

### Indications

Rheumatoid arthritis, serology unspecified

Juvenile idiopathic arthritis

## Penicillamine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 250 mg

### Indications

Rheumatoid arthritis, serology unspecified

## Probenecid

### Section

Medicines used to treat gout

Oral > Solid: 500 mg

### Indications

Gout

## Sulfasalazine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 500 mg

### Indications

Rheumatoid arthritis, serology unspecified