

Acetylsalicylic acid

Section	Indications
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Acute rheumatic fever without mention of heart involvement Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis

Adalimumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

Allopurinol

Section	Indications
Medicines used to treat gout Oral > Solid: 100 mg	Gout

Anakinra

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	Juvenile systemic arthritis

Azathioprine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 50 mg	Rheumatoid arthritis, serology unspecified Psoriatic arthritis

Certolizumab pegol

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Chloroquine

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)

Indications

Rheumatoid arthritis

Colchicine

Section

Medicines used to treat gout

Oral > Solid: 0.5 mg

Indications

Gout

Etanercept

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

Golimumab

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

Hydroxychloroquine

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 200 mg (as sulfate)

Indications

Systemic lupus erythematosus

Cutaneous lupus erythematosus of unspecified type

Infliximab

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for

Juvenile idiopathic arthritis

Methotrexate

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid > tablet: 2.5 mg (as sodium salt)	Rheumatoid arthritis, serology unspecified
Medicines for juvenile joint diseases Oral > Solid > tablet: 2.5 mg (as sodium salt)	Juvenile idiopathic arthritis

Penicillamine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 250 mg	Rheumatoid arthritis, serology unspecified

Probenecid

Section	Indications
Medicines used to treat gout Oral > Solid: 500 mg	Gout

Sulfasalazine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 500 mg	Rheumatoid arthritis, serology unspecified

Tocilizumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe	Juvenile systemic arthritis

Triamcinolone hexacetonide

Section	Indications
Medicines for juvenile joint diseases Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial	Juvenile idiopathic arthritis