

## Acetylsalicylic acid

| Section                                                                                                            | Indications                                                                                                                                                 |
|--------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Medicines for juvenile joint diseases<br>Oral > Solid: 100 to 500 mg<br>Local > Rectal > Suppository: 50 to 150 mg | <div>Acute rheumatic fever without mention of heart involvement</div> <div>Mucocutaneous lymph node syndrome</div> <div>Juvenile idiopathic arthritis</div> |

## Adalimumab

| Section                                                                                                                                               | Indications                              |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------|
| Medicines for juvenile joint diseases<br>Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL | <div>Juvenile idiopathic arthritis</div> |

## Allopurinol

| Section                                              | Indications     |
|------------------------------------------------------|-----------------|
| Medicines used to treat gout<br>Oral > Solid: 100 mg | <div>Gout</div> |

## Anakinra

| Section                                                                                                                 | Indications                            |
|-------------------------------------------------------------------------------------------------------------------------|----------------------------------------|
| Medicines for juvenile joint diseases<br>Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe | <div>Juvenile systemic arthritis</div> |

## Azathioprine

| Section                                                                | Indications                                                                          |
|------------------------------------------------------------------------|--------------------------------------------------------------------------------------|
| Disease-modifying anti-rheumatic drugs (DMARDs)<br>Oral > Solid: 50 mg | <div>Rheumatoid arthritis, serology unspecified</div> <div>Psoriatic arthritis</div> |

## Certolizumab pegol

| Section                               | Indications                                                                           |
|---------------------------------------|---------------------------------------------------------------------------------------|
| Medicines for juvenile joint diseases | Therapeutic equivalent to adalimumab pour<br><div>Juvenile idiopathic arthritis</div> |

## Chloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)

### Indications

Rheumatoid arthritis

## Colchicine

### Section

Medicines used to treat gout

Oral > Solid: 0.5 mg

### Indications

Gout

## Etanercept

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

## Golimumab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

## Hydroxychloroquine

### Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 200 mg (as sulfate)

### Indications

Systemic lupus erythematosus

Cutaneous lupus erythematosus of unspecified type

## Infliximab

### Section

Medicines for juvenile joint diseases

### Indications

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

## Methotrexate

| Section                                                                                           | Indications                                |
|---------------------------------------------------------------------------------------------------|--------------------------------------------|
| Disease-modifying anti-rheumatic drugs (DMARDs)<br>Oral > Solid > tablet: 2.5 mg (as sodium salt) | Rheumatoid arthritis, serology unspecified |
| Medicines for juvenile joint diseases<br>Oral > Solid > tablet: 2.5 mg (as sodium salt)           | Juvenile idiopathic arthritis              |

## Penicillamine

| Section                                                                 | Indications                                |
|-------------------------------------------------------------------------|--------------------------------------------|
| Disease-modifying anti-rheumatic drugs (DMARDs)<br>Oral > Solid: 250 mg | Rheumatoid arthritis, serology unspecified |

## Probenecid

| Section                                              | Indications |
|------------------------------------------------------|-------------|
| Medicines used to treat gout<br>Oral > Solid: 500 mg | Gout        |

## Sulfasalazine

| Section                                                                 | Indications                                |
|-------------------------------------------------------------------------|--------------------------------------------|
| Disease-modifying anti-rheumatic drugs (DMARDs)<br>Oral > Solid: 500 mg | Rheumatoid arthritis, serology unspecified |

## Tocilizumab

| Section                                                                                                                                                                                                                                    | Indications                 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|
| Medicines for juvenile joint diseases<br>Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial<br>Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe | Juvenile systemic arthritis |

## Triamcinolone hexacetonide

| Section                                                                                                               | Indications                   |
|-----------------------------------------------------------------------------------------------------------------------|-------------------------------|
| Medicines for juvenile joint diseases<br>Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial | Juvenile idiopathic arthritis |