



# Liste modèle des médicaments essentiels

## Aztreonam

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 1 g in vial powder for injection; 2 g in vial powder for injection	Other specified bacterial diseases

## Ceftaroline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 400 mg in vial (as fosamil) powder for injection; 600 mg in vial (as fosamil) powder for injection	Other specified bacterial diseases

## Ceftazidime + avibactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 2 g in vial + 0.5 g in vial powder for injection	Carbapenem-resistant <i>Pseudomonas aeruginosa</i> Carbapenem resistant Enterobacteriales

## Ceftolozane + tazobactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 1 g + 0.5 g powder for injection	Carbapenem-resistant <i>Pseudomonas aeruginosa</i>

## Colistin (injection)

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 1 million IU in vial (as colistemethate sodium) Parenteral > General injections > unspecified: 1 million IU in vial (as colistemethate sodium) (equivalent to 34 mg colistin base activity)	Other specified bacterial diseases Carbapenem resistant <i>Acinetobacter baumannii</i> Carbapenem-resistant <i>Pseudomonas aeruginosa</i> Carbapenem resistant Enterobacteriales

## Daptomycin

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 350 mg in vial powder for injection; 500 mg in vial powder for injection	Other specified bacterial diseases

## Eravacycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 50 mg lyophilized powder for injection	Carbapenem resistant Enterobacteriales

## Fifth generation cephalosporins

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases

## Fosfomycin (injection)

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases
Parenteral > General injections > IV: 2 g in vial (as sodium) powder for injection; 4 g in vial (as sodium) powder for injection	Carbapenem resistant Acinetobacter baumannii
	Carbapenem-resistant Pseudomonas aeruginosa
	Carbapenem resistant Enterobacteriales

## Imipenem + cilastatin + relebactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 500 mg + 500 mg + 250 mg powder for injection	Other specified other antibiotic resistant Enterobacteriales

## Linezolid

Section	Indications
<p>Reserve group antibiotics</p> <p>Parenteral &gt; General injections &gt; IV: 2 mg per mL in 300 mL bag</p> <p>Oral &gt; Liquid: 100 mg per 5 mL powder for oral liquid</p> <p>Oral &gt; Solid: 400 mg tablet; 600 mg tablet</p> <p>Oral &gt; Solid &gt; dispersible tablet: 150 mg</p> <p>Oral &gt; Solid &gt; tablet: 600 mg</p>	<p>Other specified bacterial diseases</p> <p>Methicillin resistant <i>Staphylococcus aureus</i></p> <p>Vancomycin resistant <i>Staphylococcus aureus</i></p> <p>Vancomycin resistant <i>Enterococcus</i></p>

## Meropenem + vaborbactam

Section	Indications
<p>Reserve group antibiotics</p> <p>Parenteral &gt; General injections &gt; IV: 1 g in vial + 1 g in vial powder for injection</p>	<p>Carbapenem-resistant <i>Pseudomonas aeruginosa</i></p> <p>Carbapenem resistant <i>Enterobacteriales</i></p> <p>Carbapenem resistant <i>Acinetobacter baumannii</i></p>

## Omadacycline

Section	Indications
<p>Reserve group antibiotics</p> <p>Parenteral &gt; General injections &gt; IV: 100 mg lyophilized powder for injection</p> <p>Oral &gt; Solid: 300 mg</p>	<p>Methicillin resistant <i>Staphylococcus aureus</i></p>

## Oxazolidinones

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases

## Plazomicin

Section	Indications
<p>Reserve group antibiotics</p> <p>Parenteral &gt; General injections &gt; IV: 500 mg per 10 mL (Injection)</p>	<p>Carbapenem resistant <i>Acinetobacter baumannii</i></p> <p>Carbapenem-resistant <i>Pseudomonas aeruginosa</i></p> <p>Carbapenem resistant <i>Enterobacteriales</i></p>

## Polymyxin B (injection)

Section	Indications
Reserve group antibiotics Parenteral > General injections > unspecified: 500000 IU in vial powder for injection (equivalent to 50 mg polymyxin B base)	Carbapenem resistant <i>Acinetobacter baumannii</i> Carbapenem-resistant <i>Pseudomonas aeruginosa</i> Carbapenem resistant <i>Enterobacteriales</i>

## Polymyxins

Section	Indications
Reserve group antibiotics	Other specified bacterial diseases

## Tedizolid

Section	Indications
Reserve group antibiotics	Therapeutic equivalent to <i>linezolid</i> pour <i>Methicillin resistant Staphylococcus aureus</i> Therapeutic equivalent to <i>linezolid</i> pour <i>Vancomycin resistant Staphylococcus aureus</i> Therapeutic equivalent to <i>linezolid</i> pour <i>Vancomycin resistant Enterococcus</i>

## Tigecycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 50 mg in vial powder for injection	Other specified bacterial diseases