

Acetylsalicylic acid

Section	Indications
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Acute rheumatic fever without mention of heart involvement Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis

Adalimumab

Section	Indications
Medicines for juvenile joint diseases Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

Certolizumab pegol

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Etanercept

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Golimumab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Infliximab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Methotrexate

Section

Medicines for juvenile joint diseases

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Indications

Juvenile idiopathic arthritis