

## Abemaciclib

## Section

Targeted therapies

## Indications

Therapeutic equivalent to [palbociclib](#) pour

Other specified malignant neoplasms of breast

## Abiraterone

## Section

Hormones and antihormones

Oral &gt; Solid: 250 mg; 500 mg

## Indications

Malignant neoplasms of prostate

## Afatinib

## Section

Targeted therapies

## Indications

Therapeutic equivalent to [erlotinib](#) pour

Other specified malignant neoplasms of bronchus or lung

## All-trans retinoic acid

## Section

Targeted therapies

Oral &gt; Solid: 10 mg capsule

## Indications

Acute myeloid leukaemia with recurrent genetic abnormalities

## Anastrozole

## Section

Hormones and antihormones

Oral &gt; Solid: 1 mg tablet

## Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

## Asparaginase

## Section

Cytotoxic medicines

Powder for injection: 10 000 IU in vial.

## Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

## Atezolizumab

### Section

Immunomodulators

### Indications

Therapeutic equivalent to pembrolizumab pour

Other specified malignant neoplasms of bronchus or lung

## Azacitidine

### Section

Cytotoxic medicines

Parenteral > General injections > IV: 100 mg in vial powder for injection

### Indications

Myeloid leukaemia

## Bendamustine

### Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL

### Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Follicular lymphoma

## Bicalutamide

### Section

Hormones and antihormones

Oral > Solid: 50 mg

### Indications

Malignant neoplasms of prostate

## Binimetinib

### Section

Targeted therapies

### Indications

Therapeutic equivalent to trametinib pour

Other specified melanoma of skin

## Bleomycin

### Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)

### Indications

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

## Bortezomib

### Section

#### Targeted therapies

Parenteral > General injections > unspecified: 3.5 mg in vial powder for injection

### Indications

Plasma cell myeloma

## Calcium folinate (leucovorin calcium)

### Section

#### Cytotoxic medicines

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule  
Oral > Solid > tablet: 15 mg; 5 mg; 25 mg

### Indications

Malignant neoplasm metastasis in large intestine

Malignant trophoblastic neoplasms of placenta

Burkitt lymphoma including Burkitt leukaemia

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of rectum

Malignant neoplasms of colon

## Capecitabine

### Section

#### Cytotoxic medicines

Oral > Solid: 150 mg tablet; 500 mg tablet

### Indications

Malignant neoplasms of colon

Other specified malignant neoplasms of breast

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of rectum

## Carboplatin

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL

### Indications

Other specified gliomas of brain

Other specified carcinomas of ovary

Retinoblastoma

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Malignant neoplasms of kidney, except renal pelvis

Malignant neoplasms of breast

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

## Cemiplimab

### Section

#### Immunomodulators

### Indications

Therapeutic equivalent to pembrolizumab pour

Other specified malignant neoplasms of bronchus or lung

## Chlorambucil

### Section

#### Cytotoxic medicines

Oral > Solid: 2 mg tablet

### Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

## Cisplatin

### Section

#### Cytotoxic medicines

Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection

Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL

### Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Germ cell tumour of testis

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

## Cobimetinib

### Section

#### Targeted therapies

### Indications

Therapeutic equivalent to trametinib pour

Other specified melanoma of skin

## Cyclophosphamide

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection

Oral > Solid: 25 mg; 50 mg

Oral > Solid: 25 mg; 50 mg tablet

### Indications

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of breast

Ewing sarcoma of bone and articular cartilage of unspecified sites

Lymphoid leukaemia, not elsewhere classified

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Diffuse large B-cell lymphomas

Plasma cell myeloma

Other specified gliomas of brain

Other specified malignant neoplasms of breast

Malignant trophoblastic neoplasms of placenta

Follicular lymphoma

Malignant neoplasms of kidney, except renal pelvis

Rhabdomyosarcoma primary site

Hodgkin lymphoma

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

## Cytarabine

### Section

#### Cytotoxic medicines

Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial

### Indications

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Acute myeloid leukaemia with recurrent genetic abnormalities

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

## Dabrafenib

### Section

#### Targeted therapies

Oral > Solid: 50 mg; 75 mg

### Indications

Other specified melanoma of skin

## Dacarbazine

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection

### Indications

Hodgkin lymphoma

## Dactinomycin

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 500 µg in vial powder for injection

### Indications

Malignant trophoblastic neoplasms of placenta

Malignant neoplasms of kidney, except renal pelvis

Rhabdomyosarcoma primary site

Ewing sarcoma of bone and articular cartilage of unspecified sites

## Daratumumab

### Section

#### Immunomodulators

Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL

### Indications

Plasma cell myeloma

## Dasatinib

### Section

#### Targeted therapies

Oral > Solid: 20 mg tablet; 50 mg tablet; 70 mg tablet; 80 mg tablet; 100 mg tablet; 140 mg tablet

### Indications

Chronic myeloid leukaemia, not elsewhere classified

## Daunorubicin

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)

### Indications

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Acute myeloid leukaemia with recurrent genetic abnormalities

## Dexamethasone

### Section

#### Hormones and antihormones

Oral > Solid: 4 mg

Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

Oral > Liquid: 2 mg per 5 mL

Oral > Solid: 2 mg; 4 mg

### Indications

Plasma cell myeloma

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

## Docetaxel

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL

### Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

Malignant neoplasms of prostate

## Doxorubicin

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride)

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride); 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Parenteral > General injections > IV: 50 mg in vial (hydrochloride) +; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride); 10 mg in vial (hydrochloride)

### Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Rhabdomyosarcoma primary site

Follicular lymphoma

Burkitt lymphoma including Burkitt leukaemia

Ewing sarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of breast

Lymphoid leukaemia, not elsewhere classified

Diffuse large B-cell lymphomas

Malignant neoplasms of kidney, except renal pelvis

Osteosarcoma of bone and articular cartilage of unspecified sites

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive



## Doxorubicin (as pegylated liposomal)

### Section

Cytotoxic medicines

Parenteral > General injections > IV: 2 mg per mL in 10 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

### Indications

Kaposi sarcoma of unspecified primary site

## Encorafenib

### Section

Targeted therapies

### Indications

Therapeutic equivalent to [dabrafenib](#) pour

Other specified melanoma of skin

## Enzalutamide

### Section

Hormones and antihormones

### Indications

Therapeutic equivalent to [abiraterone](#) pour

Malignant neoplasms of prostate

## Erlotinib

### Section

Targeted therapies

Oral > Solid: 100 mg; 150 mg

### Indications

Other specified malignant neoplasms of bronchus or lung

## Etoposide

| Section                                                                                                                                                                                                                                                                                                   | Indications                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Cytotoxic medicines</b><br>Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule<br>Oral > Solid: 100 mg capsule<br>Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate)<br>Oral > Solid: 100 mg capsule; 50 mg capsule | <div>Unspecified malignant neoplasms of ill-defined or unspecified sites</div> <div>Burkitt lymphoma including Burkitt leukaemia</div> <div>Ewing sarcoma of bone and articular cartilage of unspecified sites</div> <div>Lymphoid leukaemia, not elsewhere classified</div> <div>Myeloid leukaemia</div> <div>Other specified malignant neoplasms of the ovary</div> <div>Other specified malignant neoplasms of bronchus or lung</div> <div>Germ cell tumour of testis</div> <div>Malignant trophoblastic neoplasms of placenta</div> <div>Malignant neoplasms of kidney, except renal pelvis</div> <div>Osteosarcoma of bone and articular cartilage of unspecified sites</div> <div>Hodgkin lymphoma</div> <div>Retinoblastoma</div> <div>Anaplastic large cell lymphoma, ALK-negative</div> <div>Anaplastic large cell lymphoma, ALK-positive</div> |

## Everolimus

| Section                                                                                                                                                 | Indications                                 |
|---------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------|
| <b>Targeted therapies</b><br>Oral > Solid: 2.5 mg; 5 mg; 7.5 mg; 10 mg; 2 mg tablet (dispersible); 3 mg tablet (dispersible); 5 mg tablet (dispersible) | <div>Other specified gliomas of brain</div> |

## Fentanyl

| Section                                                                                                                      | Indications                    |
|------------------------------------------------------------------------------------------------------------------------------|--------------------------------|
| <b>Opioid analgesics</b><br>Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour | <div>Chronic cancer pain</div> |

## Filgrastim

### Section

#### Immunomodulators

Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial

### Indications

Acquired neutropaenia

## Fludarabine

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection

Oral > Solid: 10 mg tablet (fludarabine phosphate)

### Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

## Fluorouracil

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule

Parenteral > General injections > IV: 50 mg per mL in vial

### Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Malignant neoplasms of nasopharynx

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of breast

Malignant neoplasms of rectum

Malignant neoplasms of colon

## Flutamide

### Section

#### Hormones and antihormones

### Indications

Therapeutic equivalent to bicalutamide pour

Malignant neoplasms of prostate

## Fulvestrant

### Section

#### Hormones and antihormones

Parenteral > General injections > IV: 250 mg per 5 mL

### Indications

Malignant neoplasms of breast

## Gefitinib

### Section

Targeted therapies

### Indications

Therapeutic equivalent to erlotinib pour

Other specified malignant neoplasms of bronchus or lung

## Gemcitabine

### Section

Cytotoxic medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection

### Indications

Other specified carcinomas of ovary

Other specified malignant neoplasms of bronchus or lung

## Goserelin

### Section

Hormones and antihormones

### Indications

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of breast

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of prostate

## Hydrocortisone

### Section

Hormones and antihormones

Parenteral > General injections > unspecified: 100 mg in vial (as sodium succinate)

### Indications

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

## Hydroxycarbamide (hydroxyurea)

### Section

Cytotoxic medicines

Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g

Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg

### Indications

Squamous cell carcinoma of oropharynx

Chronic myeloid leukaemia, not elsewhere classified

## Ibrutinib

### Section

Targeted therapies

Oral > Solid: 140 mg

### Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

## Ifosfamide

| Section                                                                                                                                                                                                                                                                                 | Indications                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Cytotoxic medicines</b><br>Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection<br>Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection | <div>Other specified malignant neoplasms of the ovary</div> <div>Germ cell tumour of testis</div> <div>Ewing sarcoma of bone and articular cartilage of unspecified sites</div> <div>Osteosarcoma of bone and articular cartilage of unspecified sites</div> <div>Rhabdomyosarcoma primary site</div> <div>Unspecified malignant neoplasms of ill-defined or unspecified sites</div> <div>Burkitt lymphoma including Burkitt leukaemia</div> <div>Malignant neoplasms of kidney, except renal pelvis</div> <div>Anaplastic large cell lymphoma, ALK-negative</div> <div>Anaplastic large cell lymphoma, ALK-positive</div> |

## Imatinib

| Section                                                   | Indications                                                                                                                                                                                                                         |
|-----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Targeted therapies</b><br>Oral > Solid: 100 mg; 400 mg | <div>B lymphoblastic leukaemia or lymphoma with t(9:22) (q34;q11.2); BCR-ABL1</div> <div>Gastrointestinal stromal tumour of unspecified gastrointestinal sites</div> <div>Chronic myeloid leukaemia, not elsewhere classified</div> |

## Irinotecan

| Section                                                                                                                                                       | Indications                                                                                                                                                        |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Cytotoxic medicines</b><br>Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial | <div>Malignant neoplasm metastasis in large intestine</div> <div>Malignant neoplasms of kidney, except renal pelvis</div> <div>Rhabdomyosarcoma primary site</div> |

## Lenalidomide

| Section                                        | Indications                    |
|------------------------------------------------|--------------------------------|
| <b>Immunomodulators</b><br>Oral > Solid: 25 mg | <div>Plasma cell myeloma</div> |

## Leuporelin

### Section

#### Hormones and antihormones

Parenteral > General injections > IM: 7.5 mg in prefilled syringe; 22.5 mg in prefilled syringe

### Indications

Malignant neoplasms of breast

Malignant neoplasms of prostate

## Mercaptopurine

### Section

#### Cytotoxic medicines

Oral > Liquid: 20 mg per mL

Oral > Solid: 50 mg tablet

### Indications

Acute myeloid leukaemia with recurrent genetic abnormalities

Lymphoid leukaemia, not elsewhere classified

Langerhans cell histiocytosis

## Mesna

### Section

#### Supportive medicines

Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule

Oral > Solid: 400 mg; 600 mg

### Indications

Osteosarcoma of bone and articular cartilage of other specified sites

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Ewing sarcoma of bone and articular cartilage of unspecified sites

Rhabdomyosarcoma primary site

## Methotrexate

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection

Oral > Solid: 2.5 mg (as sodium salt)

### Indications

Lymphoid leukaemia, not elsewhere classified

Acute myeloid leukaemia with recurrent genetic abnormalities

Malignant trophoblastic neoplasms of placenta

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of breast

Osteosarcoma of bone and articular cartilage of unspecified sites

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

## Methylprednisolone

### Section

#### Hormones and antihormones

Parenteral > General injections > IV: 40 mg per mL in 1 mL single-dose vial (as sodium succinate); 40 mg per mL in 5 mL multi-dose vial (as sodium succinate); 80 mg per mL in 1 mL single-dose vial (as sodium succinate)

### Indications

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

## Nilotinib

### Section

#### Targeted therapies

Oral > Solid: 150 mg capsule; 200 mg capsule

### Indications

Chronic myeloid leukaemia, not elsewhere classified

## Nilutamide

### Section

#### Hormones and antihormones

### Indications

Therapeutic equivalent to bicalutamide pour

Malignant neoplasms of prostate

## Nivolumab

### Section

#### Immunomodulators

Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion

### Indications

Other specified malignant neoplasms of bronchus or lung

Melanoma of skin

## Normal immunoglobulin

### Section

#### Human immunoglobulins

Parenteral > General injections > IV: 5% protein solution; 10% protein solution

### Indications

Langerhans cell histiocytosis

## Osimertinib

### Section

#### Targeted therapies

Oral > Solid: 40 mg (as mesylate); 80 mg (as mesylate)

### Indications

Other specified malignant neoplasms of bronchus or lung

## Oxaliplatin

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial

### Indications

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of colon

## Paclitaxel

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 6 mg per mL

### Indications

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Kaposi sarcoma of unspecified primary site

Other specified malignant neoplasms of breast

Other specified carcinomas of ovary

Malignant neoplasms of cervix uteri

## Palbociclib

### Section

#### Targeted therapies

Oral > Solid: 75 mg; 100 mg; 125 mg

### Indications

Other specified malignant neoplasms of breast

## Pegaspargase

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection

### Indications

Lymphoid leukaemia, not elsewhere classified

## Pegfilgrastim

### Section

#### Immunomodulators

Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe

### Indications

Acquired neutropaenia



## Pembrolizumab

### Section

#### Immunomodulators

Parenteral > General injections > IV: 50 mg powder for injection

### Indications

Therapeutic equivalent to nivolumab pour

Melanoma of skin

Other specified malignant neoplasms of bronchus or lung

## Pertuzumab

### Section

#### Targeted therapies

Parenteral > General injections > IV: 240 mg per 14 mL in vial concentrated solution

### Indications

Carcinoma of breast, specialised type

## Prednisolone

### Section

#### Hormones and antihormones

Oral > Solid: 5 mg; 25 mg

Oral > Liquid: 5 mg per mL

### Indications

Plasma cell myeloma

Follicular lymphoma

Lymphoid leukaemia, not elsewhere classified

Malignant neoplasms of prostate

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Hodgkin lymphoma

Diffuse large B-cell lymphomas

## Prednisone

### Section

Hormones and antihormones

### Indications

Therapeutic equivalent to prednisolone pour

Plasma cell myeloma

Therapeutic equivalent to prednisolone pour

Follicular lymphoma

Therapeutic equivalent to prednisolone pour

Lymphoid leukaemia, not elsewhere classified

Therapeutic equivalent to prednisolone pour

Malignant neoplasms of prostate

Therapeutic equivalent to prednisolone pour

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Therapeutic equivalent to prednisolone pour

Hodgkin lymphoma

Therapeutic equivalent to prednisolone pour

Diffuse large B-cell lymphomas

## Procarbazine

### Section

Cytotoxic medicines

Oral > Solid: 50 mg (as hydrochloride)

### Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Hodgkin lymphoma

## Ribociclib

### Section

Targeted therapies

### Indications

Therapeutic equivalent to palbociclib pour

Other specified malignant neoplasms of breast

## Rituximab

### Section

Targeted therapies

Parenteral > General injections > IV: 100 mg per 10 mL in 10 mL vial; 500 mg per 50 mL in 50 mL vial

### Indications

Follicular lymphoma

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Diffuse large B-cell lymphomas

Burkitt lymphoma including Burkitt leukaemia

## Tamoxifen

### Section

Hormones and antihormones

Oral > Solid: 10 mg (as citrate); 20 mg (as citrate)

### Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

## Thalidomide

### Section

Immunomodulators

Oral > Solid: 50 mg

### Indications

Plasma cell myeloma

## Tioguanine

### Section

Cytotoxic medicines

Oral > Solid: 40 mg

### Indications

Lymphoid leukaemia, not elsewhere classified

## Tislelizumab

### Section

Immunomodulators

Parenteral > General injections > IV: 100 mg per 10 mL

### Indications

Malignant neoplasms of urinary tract

Hodgkin lymphoma

Other specified malignant neoplasms of bronchus or lung

## Toripalimab

### Section

Immunomodulators

Parenteral > General injections > IV: 240 mg per 6 mL in vial

### Indications

Malignant neoplasms of nasopharynx

Malignant neoplasms of oesophagus

## Trametinib

### Section

Targeted therapies

Oral > Solid: 0.5 mg; 2 mg

### Indications

Other specified melanoma of skin

## Trastuzumab

### Section

#### Targeted therapies

Parenteral > General injections > IV: 60 mg in vial powder for injection; 150 mg in vial powder for injection; 440 mg in vial powder for injection

### Indications

Carcinoma of breast, specialised type

## Triptorelin

### Section

#### Hormones and antihormones

### Indications

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of breast

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of prostate

## Vemurafenib

### Section

#### Targeted therapies

### Indications

Therapeutic equivalent to dabrafenib pour

Other specified melanoma of skin

## Vinblastine

### Section

#### Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)

### Indications

Other specified gliomas of brain

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

## Vincristine

| Section                                                                                                                                                                                                                            | Indications                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Cytotoxic medicines</b><br>Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate) | <ul style="list-style-type: none"><li>Other specified gliomas of brain</li><li>Malignant trophoblastic neoplasms of placenta</li><li>Follicular lymphoma</li><li>Kaposi sarcoma of unspecified primary site</li><li>Rhabdomyosarcoma primary site</li><li>Hodgkin lymphoma</li><li>Diffuse large B-cell lymphomas</li><li>Retinoblastoma</li><li>Burkitt lymphoma including Burkitt leukaemia</li><li>Malignant neoplasms of kidney, except renal pelvis</li><li>Ewing sarcoma of bone and articular cartilage of unspecified sites</li><li>Lymphoid leukaemia, not elsewhere classified</li><li>Langerhans cell histiocytosis</li></ul> |

## Vinorelbine

| Section                                                                                                                                                                                                                                                       | Indications                                                                                                                                                                                         |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Cytotoxic medicines</b><br>Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 50 mg per 5 mL in 5 mL vial<br>Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial<br>Oral > Solid: 20 mg; 30 mg; 80 mg | <ul style="list-style-type: none"><li>Other specified malignant neoplasms of bronchus or lung</li><li>Other specified malignant neoplasms of breast</li><li>Rhabdomyosarcoma primary site</li></ul> |

## Zanubrutinib

| Section                                          | Indications                                                                                                                              |
|--------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Targeted therapies</b><br>Oral > Solid: 80 mg | <ul style="list-style-type: none"><li>Mantle cell lymphoma</li><li>Chronic lymphocytic leukaemia or small lymphocytic lymphoma</li></ul> |

## Zoledronic acid

| Section                                                                                                                           | Indications                                                                                          |
|-----------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| <b>Supportive medicines</b><br>Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle | <ul style="list-style-type: none"><li>Malignant neoplasm metastasis in bone or bone marrow</li></ul> |

