

BCG vaccine

Section

Recommendations for all immunization programmes
All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against tuberculosis

Japanese encephalitis vaccine

Section

Recommendations for certain regions
All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against certain single viral diseases

Medicines for COVID-19

Section

Medicines for COVID-19
Refer to WHO living guidelines

Indications

COVID-19

Abacavir

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors
Oral > Solid: 300 mg tablet (as sulfate)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abacavir + lamivudine

Section

Fixed-dose combinations of antiretrovirals
Oral > Solid: 120 mg (as sulfate) + 60 mg tablet (dispersible, scored)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abacavir + lamivudine + lopinavir + ritonavir

Section

Fixed-dose combinations of antiretrovirals
Oral > Solid: 30 mg + 15 mg + 40 mg + 10 mg capsule containing oral granules

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Abemaciclib

Section

Targeted therapies

Indications

Therapeutic equivalent to **palbociclib** pour

Other specified malignant neoplasms of breast

Abiraterone

Section

Hormones and antihormones

Oral > Solid: 250 mg; 500 mg

Indications

Malignant neoplasms of prostate

Acamprosate calcium

Section

Medicines for alcohol use disorders

Oral > Solid > tablet: 333 mg

Indications

Disorders due to use of alcohol

Acetazolamide

Section

Ophthalmological preparations > Miotics and antiglaucoma medicines

Oral > Solid: 250 mg

Indications

Primary open-angle glaucoma

Acetic acid

Section

Ear, nose and throat medicines [c]

Local > Otological > drops: 2% in alcohol

Indications

Infectious diseases of external ear

Acetylcysteine

Section

Antidotes and other substances used in poisonings > Specific

Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule

Oral > Liquid: 10%; 20%

Indications

Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Acute or subacute hepatic failure

Acetylsalicylic acid

Section	Indications
Antimigraine medicines > For treatment of acute attack Oral > Solid: 300 to 500 mg	Migraine
Anti-platelet medicines Oral > Solid: 100 mg	Acute myocardial infarction Cerebral ischaemic stroke
Medicines for juvenile joint diseases Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Acute rheumatic fever without mention of heart involvement Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs) Oral > Solid: 100 to 500 mg Local > Rectal > Suppository: 50 to 150 mg	Pain

Acetylsalicylic acid + atorvastatin + ramipril

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 100 mg + 20 mg + 2.5 mg; 100 mg + 20 mg + 5 mg; 100 mg + 20 mg + 10 mg; 100 mg + 40 mg + 2.5 mg; 100 mg + 40 mg + 5 mg; 100 mg + 40 mg + 10 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Acetylsalicylic acid + simvastatin + ramipril + atenolol + hydrochlorothiazide

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease Oral > Solid > tablet: 100 mg + 20 mg + 5 mg + 50 mg + 12.5 mg	Atherosclerotic chronic arterial occlusive disease Chronic ischaemic heart disease Cerebral ischaemic stroke

Aciclovir

Section	Indications
Antitherpes medicines	Zoster
Parenteral > General injections > IV: 250 mg in vial powder for injection (as sodium salt)	Varicella
Oral > Liquid: 200 mg per 5 mL	Herpes simplex infections
Oral > Solid: 200 mg tablet	
Ophthalmological preparations > Anti-infective agents	Herpes simplex keratitis
Local > Ophthalmological > Ointment: 3% w/w	

Acclidinium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour
	Chronic obstructive pulmonary disease

Activated charcoal

Section	Indications
Antidotes and other substances used in poisonings > Non-specific	Unspecified injury, poisoning or certain other consequences of external causes
Oral > Other: powder	

Adalimumab

Section	Indications
Immunomodulators for non-malignant disease	Axial spondyloarthritis
Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Crohn disease site
	Rheumatoid arthritis
Medicines for juvenile joint diseases	Juvenile idiopathic arthritis
Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	

Afatinib

Section	Indications
Targeted therapies	Therapeutic equivalent to erlotinib pour
	Other specified malignant neoplasms of bronchus or lung

Albendazole

Section	Indications
Intestinal anthelmintics Oral > Solid > tablet: 400 mg (chewable, scored) Oral > Solid > tablet: 400 mg (chewable, scored))	Echinococcosis Ascariasis Enterobiasis Trichuriasis Hookworm diseases Cysticercosis
Cysticidal medicines Oral > Solid > tablet: 400 mg (chewable, scored); 200 mg (chewable)	Other specified echinococcosis Cysticercosis of central nervous system
Antifilarials Oral > Solid > tablet: 400 mg (chewable, scored)	Filariasis

Alcohol based hand rub

Section	Indications
Disinfectants Local > Topical > Solution: 80% v/v ethanol; 75% v/v isopropyl alcohol	Denatured alcohol

Alcuronium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 5 mg per mL in 2 mL ampoule (alcuronium chloride)	Muscle relaxants

Alfacalcidol

Section	Indications
Medicines for endocrine disorders Oral > Liquid: 2 µg per mL Oral > Solid > capsule: 0.25 µg; 1 µg	Chronic kidney disease, stage unspecified Disorders of vitamin D metabolism or transport

All-trans retinoic acid

Section	Indications
Targeted therapies Oral > Solid: 10 mg capsule	Acute myeloid leukaemia with recurrent genetic abnormalities

Allopurinol

Section

Medicines used to treat gout

Oral > Solid: 100 mg

Supportive medicines

Oral > Solid: 100 mg; 300 mg

Indications

Gout

Tumour lysis syndrome

Alteplase

Section

Thrombolytic medicines

Parenteral > General injections > IV: 10 mg in vial powder for injection; 20 mg in vial powder for injection; 50 mg in vial powder for injection

Indications

Cerebral ischaemic stroke

Amidotrizoate

Section

Diagnostic agents > Radiocontrast media

Parenteral > General injections > unspecified: 140 to 420 mg iodine per mL in 20 mL ampoule (as sodium or meglumine salt)

Indications

Amidotrizoate

Amikacin

Section

Ophthalmological preparations > Anti-infective agents

Antituberculosis medicines

Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)

Access group antibiotics

Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate); 50 mg per mL in 2 mL vial (as sulfate)

Parenteral > General injections > unspecified: 250 mg per mL in 2 mL vial (as sulfate)

Indications

Therapeutic equivalent to [gentamicin](#) pour

Other specified conjunctivitis

Therapeutic equivalent to [gentamicin](#) pour

Infectious blepharitis

Multi-drug resistant *Mycobacterium tuberculosis*

Premier choix

Neutropenia (high-risk)

co-prescrire avec [cefotaxime](#)

Acute pyelonephritis (severe)

co-prescrire avec [cefotaxime](#)

Inflammatory and other diseases of prostate (severe)

co-prescrire avec [ceftriaxone](#)

Acute pyelonephritis (severe)

co-prescrire avec [ceftriaxone](#)

Inflammatory and other diseases of prostate (severe)

Second choix

co-prescrire avec [cloxacillin](#)

Sepsis without septic shock

Amiloride

Section

Diuretics

Oral > Solid: 5 mg (hydrochloride)

Indications

Oedema

Ascites

Amiodarone

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 50 mg per mL in 3 ampoule (hydrochloride)

Oral > Solid: 100 mg (hydrochloride); 200 mg (hydrochloride); 400 mg (hydrochloride)

Indications

Cardiac arrhythmia

Amitriptyline

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 25 mg; 10 mg; 75 mg	Palliative care
Medicines for depressive disorders Oral > Solid: 25 mg; 75 mg	Depressive disorders

Amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 5 mg (as maleate, mesylate or besylate)	Essential hypertension

Amodiaquine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 153 mg (as hydrochloride); 200 mg (as hydrochloride)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Amodiaquine + sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Solid: 76.5 mg (as hydrochloride) [3] + 250 mg + 12.5 mg [1] dispersible tablets in co-package; 153 mg (as hydrochloride) [3] + 500 mg + 25 mg [1] dispersible tablets in co-package	Malaria

Amoxicillin

Section

Access group antibiotics

Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid

Oral > Solid: 250 mg (anhydrous) capsule or tablet; 500 mg (anhydrous) capsule or tablet

Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid; 250 mg per 5 mL (as trihydrate) powder for oral liquid

Oral > Solid: 250 mg (as trihydrate); 500 mg (as trihydrate)

Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection

Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate); 500 mg (scored) (as trihydrate)

Oral > Solid > dosage form: 250 mg (as trihydrate); 500 mg (as trihydrate)

Indications

Premier choix

Acute sinusitis

Acute otitis media

Acute pharyngitis

Acute malnutrition in infants, children or adolescents (complicated) [children]

Acute malnutrition in infants, children or adolescents (uncomplicated) [children]

Chronic obstructive pulmonary disease with acute exacerbation

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

co-prescrire avec [gentamicin](#)

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec [gentamicin](#)

Sepsis without septic shock

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Infectious cystitis

Amoxicillin + clavulanic acid

Section

Access group antibiotics

Parenteral > General injections > IV: 500 mg (as sodium salt) + 100 mg (as potassium salt) powder for injection; 1000 mg (as sodium salt) + 200 mg (as potassium salt) powder for injection

Oral > Solid: 500 mg (as trihydrate) + 125 mg (as potassium salt)

Oral > Liquid: 250 mg + 62.5 mg per 5 mL powder for oral liquid; 125 mg + 31.25 mg per 5 mL powder for oral liquid

Oral > Solid: 500 mg + 125 mg tablet

Oral > Liquid: 125 mg (as trihydrate) + 31.25 mg (as potassium salt) powder for oral liquid; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) powder for oral liquid

Oral > Solid: 500 mg (as trihydrate) + 125 mg (as potassium salt); 875 mg (as trihydrate) + 125 mg (as potassium salt)

Oral > Solid > dispersible tablet: 200 mg (as trihydrate) + 28.5 mg (as potassium salt); 250 mg (as trihydrate) + 62.5 mg (as potassium salt)

Oral > Liquid: 125 mg + 31.25 mg powder for oral liquid; 250 mg + 62.5 mg powder for oral liquid

Antituberculosis medicines

Oral > Liquid: 250 mg (as trihydrate) + 62.5 mg per 5 mL (as potassium salt) powder for oral liquid

Oral > Solid: 500 mg (as trihydrate) + 125 mg (as potassium salt)

Indications

Premier choix

Peritoneal abscess (mild-moderate)

Neutropenia (low-risk)

Bacterial cellulitis, erysipelas or lymphangitis

Acute sinusitis

Peritonitis (mild-moderate)

Infectious cystitis

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Chronic obstructive pulmonary disease with acute exacerbation

Second choix

Other specified prophylactic measures

Acute otitis media

Osteomyelitis or osteitis

Bacterial infection of joint

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

co-prescrire avec clarithromycin

Bacterial pneumonia (Community-acquired pneumonia - severe)

Autres indications

Bacterial infection of unspecified site

Multi-drug resistant Mycobacterium tuberculosis

Amphotericin B

Section

Antifungal medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)

Antileishmaniasis medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)

Indications

Blastomycosis

Cryptococcosis

Candidosis

Coccidioidomycosis

Histoplasmosis

Sporotrichosis

Aspergillosis

Visceral leishmaniasis

Ampicillin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 1 g in vial (as sodium salt) powder for injection

Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral suspension

Oral > Solid: 250 mg (anhydrous); 500 mg (anhydrous)

Indications

Premier choix

co-prescrire avec [gentamicin](#)

Peritoneal abscess (severe)

co-prescrire avec [gentamicin](#)

Sepsis without septic shock

co-prescrire avec [gentamicin](#)

Peritoneal abscess (mild-moderate)

co-prescrire avec [gentamicin](#)

Peritonitis (severe)

co-prescrire avec [gentamicin](#)

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec [gentamicin](#)

Peritonitis (mild-moderate)

co-prescrire avec [gentamicin](#)

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Anakinra

Section

Medicines for juvenile joint diseases

Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe

Indications

Juvenile systemic arthritis

Anastrozole

Section

Hormones and antihormones

Oral > Solid: 1 mg tablet

Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

Anidulafungin

Section

Antifungal medicines

Indications

Therapeutic equivalent to micafungin pour

Systemic or invasive candidosis

Ansuvimab

Section

Medicines for Ebola virus disease

Parenteral > General injections > IV: 400 mg powder for injection

Indications

Ebola virus disease

Anti-d immunoglobulin

Section

Human immunoglobulins

Parenteral > General injections > IV: 250 µg in single-dose vial

Parenteral > General injections > IM: 250 µg in single-dose vial

Indications

Maternal care for red cell antibodies

Rh incompatibility reaction

Anti-rabies immunoglobulin

Section

Human immunoglobulins

Parenteral > General injections > IM: 150 IU per mL in vial

Parenteral > Locoregional injections > Other: 150 IU per mL in vial

Indications

Rabies

Anti-rabies virus monoclonal antibodies

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies
Parenteral > Locoregional injections > Intradermal: 40 IU per mL in 1.25 mL vial (human); 40 IU per mL in 2.5 mL vial (human); 100 IU per mL in 2.5 mL vial (human); 300 IU per mL in 10 mL vial (murine); 600 IU per mL in 1 mL vial (murine); 600 IU per mL in 2.5 mL vial (murine); 600 IU per mL in 5 mL vial (murine)

Indications

Rabies

Anti-tetanus immunoglobulin

Section

Human immunoglobulins
Parenteral > General injections > IM: 500 IU in vial

Indications

Tetanus

Antirabies hyperimmune serum

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies
Parenteral > General injections > unspecified: 1000 IU in 5 mL ampoule

Indications

Rabies

Antivenom immunoglobulin

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies
Parenteral > General injections > IV:

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Antivenom sera

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies
Parenteral > General injections > IV:

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Apixaban

Section

Medicines affecting coagulation

Indications

Therapeutic equivalent to [dabigatran](#) pour

Venous thromboembolism

Therapeutic equivalent to [dabigatran](#) pour

Other specified atrial fibrillation

Aprepitant

Section	Indications
Antiemetic medicines Oral > Liquid: 125 mg powder for oral suspension (in sachet) Oral > Solid: 80 mg; 125 mg; 165 mg	Nausea or vomiting

Aripiprazole

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders

Arsenic trioxide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg per mL concentrate for solution for infusion; 2 mg per mL concentrate for solution for infusion	Acute myeloid leukaemia with recurrent genetic abnormalities

Artemether

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IM: 80 mg per mL in 1 mL ampoule (oily injection)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artemether + lumefantrine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 20 mg + 120 mg tablet; 20 mg + 120 mg dispersible tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate

Section	Indications
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 60 mg in ampoule containing anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution Oral > Solid: 50 mg tablet Local > Rectal > Other: 50 mg capsules; 200 mg capsules; 100 mg capsules	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax

Artesunate + amodiaquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 67.5 mg tablet; 50 mg + 135 mg tablet; 100 mg + 270 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + mefloquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 25 mg + 55 mg tablet; 100 mg + 220 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Artesunate + pyronaridine tetraphosphate

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 60 mg + 180 mg tablet; 20 mg + 60 mg granules

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Ascorbic acid

Section

Vitamins and minerals

Oral > Solid: 50 mg

Indications

Scurvy

Asparaginase

Section

Cytotoxic medicines

Powder for injection: 10 000 IU in vial.

Parenteral > General injections > unspecified: 10000 IU in vial powder for injection

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Lymphoid leukaemia, not elsewhere classified

Atazanavir

Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 100 mg (as sulfate); 300 mg (as sulfate)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Atazanavir + ritonavir

Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 300 mg (as sulfate) + 100 mg tablet (heat stable)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Atenolol

Section

Antianginal medicines

Oral > Solid: 50 mg; 100 mg

Antihypertensive medicines

Oral > Solid: 50 mg; 100 mg

Indications

Angina pectoris

Essential hypertension

Therapeutic equivalent to [bisoprolol](#) pour

Essential hypertension

Atezolizumab

Section

Immunomodulators

Parenteral > General injections > IV: 1.2 g per 20 mL concentrate solution for infusion

Indications

Other specified malignant neoplasms of bronchus or lung

Therapeutic equivalent to [pembrolizumab](#) pour

Other specified malignant neoplasms of bronchus or lung

Atoltivimab + maftivimab + odesivimab

Section

Medicines for Ebola virus disease

Parenteral > General injections > IV: 241.7 mg + 241.7 mg + 241.7 mg in 14.5 mL vial

Indications

Ebola virus disease

Atorvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Atorvastatin + perindopril + amlodipine

Section	Indications
Fixed-dose combinations for prevention of atherosclerotic cardiovascular disease	Atherosclerotic chronic arterial occlusive disease
Oral > Solid > tablet: 20 mg + 5 mg + 5 mg; 20 mg + 10 mg + 10 mg; 40 mg + 5 mg + 5 mg; 40 mg + 10 mg + 10 mg	Chronic ischaemic heart disease
	Cerebral ischaemic stroke

Atracurium

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors	Muscle relaxants
Parenteral > General injections > IV: 10 mg per mL (besylate)	Therapeutic equivalent to vecuronium pour
	Muscle relaxants

Atropine

Section	Indications
Antidotes and other substances used in poisonings > Specific	Exposure to or harmful effects of undetermined intent of pesticides
Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	
Preoperative medication and sedation for short-term procedures	Parasympatholytics [anticholinergics and antimuscarinics] and spasmolytics
Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (atropine sulfate)	
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide pour
	Cycloplegic drug
Ophthalmological preparations > Mydriatics	Anterior uveitis
Local > Ophthalmological > Solution (eye drops): 0.1% (sulfate); 0.5% (sulfate); 1% (sulfate)	

Axicabtagene ciloleucel

Section	Indications
Antineoplastics and supportive medicines	Diffuse large B-cell lymphomas
Cell suspension for infusion: 2×10^6 anti-CD19 CAR-positive viable T-cells per kg of body weight.	

Azacitidine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 100 mg in vial powder for injection

Indications

Myeloid leukaemia

Azathioprine

Section

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 50 mg

Immunomodulators for non-malignant disease

Oral > Solid: 50 mg

Parenteral > General injections > IV: 100 mg in vial powder for injection (as sodium salt); 50 mg in vial powder for injection (as sodium salt)

Oral > Liquid: 10 mg per mL

Oral > Solid > tablet: 50 mg (scored); 25 mg

Indications

Rheumatoid arthritis, serology unspecified

Psoriatic arthritis

Relapsing-remitting multiple sclerosis

Failure or rejection of transplanted organs or tissues

Azithromycin

Section

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Solution (eye drops): 1.5% eye drops

Watch group antibiotics

Oral > Liquid: 200 mg per 5 mL oral liquid

Oral > Solid: 250 mg (anhydrous) capsule; 500 mg (anhydrous) capsule

Oral > Liquid: 200 mg per 5 mL (anhydrous) powder for oral liquid

Oral > Solid > dosage form: 250 mg (anhydrous); 500 mg (anhydrous)

Oral > Solid > dosage form: 500 mg (anhydrous); 250 mg (anhydrous)

Indications

Trachoma

Premier choix

Typhoid fever

Trachoma

Chlamydia trachomatis

Yaws

Paratyphoid fever

co-prescrire avec ceftriaxone

Gonococcal infection

Second choix

Infectious gastroenteritis or colitis without specification of infectious agent

co-prescrire avec cefixime

Gonococcal infection

Aztreonam

Section

Reserve group antibiotics

Parenteral > General injections > unspecified: 1 g in vial powder for injection; 2 g in vial powder for injection

Indications

Other specified bacterial diseases

Barium sulfate

Section

Diagnostic agents > Radiocontrast media

Oral > Liquid: Aqueous suspension

Indications

Barium sulfate with suspending agents

Beclometasone

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Respiratory > Inhalation > aerosol: 50 µg per dose (dipropionate); 250 µg per dose (dipropionatae)

Indications

Therapeutic equivalent to budesonide pour

Asthma

Asthma

Beclometasone + formoterol

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to budesonide + formoterol pour

Asthma

Bedaquiline

Section

Antituberculosis medicines

Oral > Solid: 100 mg tablet; 20 mg tablet

Indications

Multi-drug resistant Mycobacterium tuberculosis

Bendamustine

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 45 mg per 0.5 mL; 180 mg per 2 mL

Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Follicular lymphoma

Benzathine benzylpenicillin

Section

Access group antibiotics

Parenteral > General injections > IM: 900 mg in vial benzylpenicillin powder for injection (= 1.2 million IU); 1.44 g in vial benzylpenicillin powder for injection (= 2.4 million IU)

Indications

Premier choix

Syphilis

Congenital syphilis [children]

Autres indications

Bacterial infection of unspecified site

Benznidazole

Section

American trypanosomiasis

Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg

Indications

Chagas disease

Benzoic acid + salicylic acid

Section

Dermatological medicines > Antifungal medicines

Local > Topical > Cream: 6% + 3%

Local > Topical > Ointment: 6% + 3%

Indications

Fungal infection of the skin

Benzoyl peroxide

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Local > Topical > Cream: 5%

Local > Topical > Lotion: 5%

Indications

Acne

Benzyl benzoate

Section

Dermatological medicines > Scabicides and pediculicides

Local > Topical > Lotion: 25%

Indications

Pthiriasis

Pediculosis corporis

Pediculosis capitis

Scabies

Benzylpenicillin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 600 mg in vial powder for injection (=1 million IU as sodium or potassium salt); 3 g in vial powder for injection (=5 million IU as sodium or potassium salt)

Parenteral > General injections > unspecified: 600 mg in vial powder for injection (= 1 million IU as sodium or potassium salt); 3 g in vial powder for injection (= 5 million IU as sodium or potassium salt)

Indications

Premier choix

Congenital syphilis [children]

Syphilis

co-prescrire avec [gentamicin](#)

Sepsis without septic shock

co-prescrire avec [gentamicin](#)

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec [gentamicin](#)

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Second choix

Bacterial meningitis

Autres indications

Bacterial infection of unspecified site

Betamethasone

Section

Dermatological medicines > Anti-inflammatory and antipruritic medicines

Local > Topical > Cream: 0.1% (as valerate)

Local > Topical > Ointment: 0.1% (as valerate)

Indications

Pruritus due to skin disorder

Psoriasis of unspecified type

Dermatitis or eczema

Lichen planus

Bevacizumab

Section

Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations

Parenteral > Locoregional injections > Intravitreal: 25 mg per mL

Indications

Age related macular degeneration

Bicalutamide

Section

Hormones and antihormones

Oral > Solid: 50 mg

Indications

Malignant neoplasms of prostate

Binimetinib

Section

Targeted therapies

Indications

Therapeutic equivalent to [trametinib](#) pour

Other specified melanoma of skin

Biperiden

Section

Medicines for parkinsonism

Parenteral > General injections > IV: 5 mg per mL in 1 mL ampoule (lactate)

Oral > Solid: 2 mg tablet (hydrochloride)

Indications

Parkinson disease

Bisacodyl

Section

Laxatives

Indications

Therapeutic equivalent to [senna](#) pour

Constipation

Bisoprolol

Section

Antianginal medicines

Oral > Solid: 1.25 mg; 5 mg

Medicines used in heart failure

Oral > Solid: 1.25 mg; 5 mg

Antihypertensive medicines

Oral > Solid: 1.25 mg; 5 mg

Antiarrhythmic medicines

Oral > Solid: 1.25 mg; 5 mg

Indications

Angina pectoris

Heart failure

Essential hypertension

Cardiac arrhythmia

Bleomycin

Section	Indications
Cytotoxic medicines Parenteral > General injections > unspecified: 15 mg in vial (as sulfate) Parenteral > General injections > unspecified: 15000 IU in vial (as sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Kaposi sarcoma of unspecified primary site Hodgkin lymphoma Other specified malignant neoplasms of the ovary Germ cell tumour of testis

Bortezomib

Section	Indications
Targeted therapies Parenteral > General injections > unspecified: 3.5 mg in vial powder for injection	Plasma cell myeloma

Bromocriptine

Section	Indications
Antiseizure medicines Oral > Solid: 2.5 mg; 5 mg	Parkinson disease
Medicines for disorders of the pituitary hormone system	Therapeutic equivalent to cabergoline pour Other specified benign neoplasm of endocrine glands

Budesonide

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > aerosol: 100 µg per dose; 200 µg per dose	Asthma
Ear, nose and throat medicines [c] Local > Nasal > Spray: 100 µg per dose	Allergic rhinitis

Budesonide + formoterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease Respiratory > Inhalation > dry powder: 100 µg + 6 µg per dose; 200 µg + 6 µg per dose	Asthma

Budesonide + salmeterol

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to budesonide + formoterol pour Asthma

Bumetanide

Section	Indications
Medicines used in heart failure	Therapeutic equivalent to furosemide pour Heart failure
Diuretics	Therapeutic equivalent to furosemide pour Anuria or oliguria Therapeutic equivalent to furosemide pour Oedema

Bupivacaine

Section	Indications
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 0.5% in 4 mL ampoule (hydrochloride) + 7.5% glucose solution Parenteral > General injections > unspecified: 0.25% in vial (hydrochloride); 0.5% in vial (hydrochloride)	Local anaesthetics

Buprenorphine

Section	Indications
Medicines for opioid use disorders	Therapeutic equivalent to methadone pour Opioid dependence

Bupropion

Section	Indications
Medicines for nicotine use disorders Oral > Solid: 150 mg sustained-release (hydrochloride)	Nicotine dependence

C1 esterase inhibitor

Section

Plasma-derived medicines

Parenteral > General injections > IV: 500 IU in vial powder for injection

Indications

Hereditary angioedema

Cabergoline

Section

Antiseizure medicines

Oral > Solid: 0.5 mg

Medicines for disorders of the pituitary hormone system

Oral > Solid > tablet: 0.5 mg; 1 mg

Indications

Parkinson disease

Other specified benign neoplasm of endocrine glands

Caffeine citrate

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 20 mg per mL (equivalent to 10 mg caffeine base per mL)

Oral > Liquid: 20 mg per mL (equivalent to 10 mg caffeine base per mL)

Indications

Apnoea of newborn

Calamine

Section

Dermatological medicines > Anti-inflammatory and antipruritic medicines

Local > Topical > Lotion:

Indications

Pruritus

Calcipotriol

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Local > Topical > Cream: 50 µg per mL (0.005%)

Local > Topical > Lotion: 50 µg per mL (0.005%)

Local > Topical > Ointment: 50 µg per mL (0.005%)

Indications

Psoriasis of unspecified type

Calcitriol

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Medicines for endocrine disorders

Oral > Solid > capsule: 0.25 µg; 0.5 µg

Indications

Therapeutic equivalent to calcipotriol pour

Psoriasis of unspecified type

Chronic kidney disease, stage unspecified

Disorders of vitamin D metabolism or transport

Calcium

Section

Vitamins and minerals

Oral > Solid: 500 mg (elemental calcium)

Indications

Calcium deficiency

Calcium folinate (leucovorin calcium)

Section

Cytotoxic medicines

Oral > Solid: 15 mg tablet

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule

Parenteral > General injections > unspecified: 3 mg per mL in 10 mL ampoule; 7.5 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Solid > tablet: 15 mg; 5 mg; 25 mg

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Malignant neoplasm metastasis in large intestine

Malignant trophoblastic neoplasms of placenta

Burkitt lymphoma including Burkitt leukaemia

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of rectum

Malignant neoplasms of colon

Calcium gluconate

Section

Antidotes and other substances used in poisonings > Specific

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule

Vitamins and minerals

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule

Indications

Hypermagnesaemia

Tetany due to acute calcium deficiency

Canagliflozin

Section

Oral hypoglycaemic agents

Indications

Therapeutic equivalent to empagliflozin pour

Type 2 diabetes mellitus

Capecitabine

Section

Cytotoxic medicines

Oral > Solid: 150 mg tablet; 500 mg tablet

Indications

Malignant neoplasms of colon

Other specified malignant neoplasms of breast

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of rectum

Capreomycin

Section

Antituberculosis medicines

Parenteral > General injections > IV: 1000 mg powder for injection (vial)

Indications

Multi-drug resistant Mycobacterium tuberculosis

Captopril

Section

Medicines used in heart failure

Oral > Solid: 25 mg

Indications

Heart failure

Carbachol

Section

Ophthalmological preparations > Miotics and antiglaucoma medicines

Indications

Therapeutic equivalent to pilocarpine pour

Acute angle closure with pupillary block

Therapeutic equivalent to pilocarpine pour

Ocular hypertension

Therapeutic equivalent to pilocarpine pour

Primary open-angle glaucoma

Carbamazepine

Section

Antiseizure medicines

Oral > Liquid: 100 mg per 5 mL

Oral > Solid > tablet: 200 mg (scored); 100 mg (scored); 100 mg (chewable); 200 mg (chewable); 400 mg (scored)

Medicines for bipolar disorders

Oral > Solid > tablet: 100 mg (scored); 200 mg (scored); 400 mg

Indications

Epilepsy or seizures

Bipolar or related disorders

Carbetocin

Section

Uterotonics

Parenteral > General injections > IV: 100 µg per mL (heat stable)

Indications

Postpartum haemorrhage

Carbimazole

Section

Thyroid hormones and antithyroid medicines

Indications

Therapeutic equivalent to methimazole pour

Thyrotoxicosis

Carboplatin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg per 5 mL; 150 mg per 15 mL; 450 mg per 45 mL; 600 mg per 60 mL

Indications

Other specified gliomas of brain

Other specified carcinomas of ovary

Retinoblastoma

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Malignant neoplasms of kidney, except renal pelvis

Malignant neoplasms of breast

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Carmellose

Section

Ophthalmological preparations

Indications

Therapeutic equivalent to [hypromellose](#) pour

Keratoconjunctivitis sicca

Carvedilol

Section

Antianginal medicines

Indications

Therapeutic equivalent to [bisoprolol](#) pour

Angina pectoris

Medicines used in heart failure

Therapeutic equivalent to [bisoprolol](#) pour

Heart failure

Antihypertensive medicines

Therapeutic equivalent to [bisoprolol](#) pour

Essential hypertension

Antiarrhythmic medicines

Therapeutic equivalent to [bisoprolol](#) pour

Cardiac arrhythmia

Caspofungin

Section

Antifungal medicines

Indications

Therapeutic equivalent to [micafungin](#) pour

Systemic or invasive candidosis

Cefalexin

Section

[Access group antibiotics](#)

Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid; 250 mg per 5 mL (anhydrous) powder for oral liquid

Oral > Solid: 250 mg (as monohydrate)

Oral > Solid: 250 mg (as monohydrate); 500 mg (as monohydrate)

Oral > Solid > dispersible tablet: 125 mg; 250 mg

Indications

Premier choix

Bacterial cellulitis, erysipelas or lymphangitis

Second choix

Acute pharyngitis

Chronic obstructive pulmonary disease with acute exacerbation

Autres indications

Bacterial infection of unspecified site

Cefazolin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 1 g in vial (as sodium salt) powder for injection

Indications

Premier choix

Other specified prophylactic measures

co-prescrire avec metronidazole

Other specified prophylactic measures

Second choix

Osteomyelitis or osteitis

Bacterial infection of joint

Autres indications

Surgical site infection

Cefepime

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection; 2 g in vial (as hydrochloride) powder for injection

Indications

Other specified bacterial diseases

Cefixime

Section

Watch group antibiotics

Oral > Liquid: 100 mg per 5 mL powder for oral liquid

Oral > Solid: 200 mg (as trihydrate); 400 mg (as trihydrate)

Indications

Second choix

Infectious gastroenteritis or colitis without specification of infectious agent

co-prescrire avec azithromycin

Gonococcal infection

Cefotaxime

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)

Indications

Premier choix

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

Other specified pneumonia (Hospital-acquired pneumonia)

Bacterial meningitis

co-prescrire avec amikacin

Inflammatory and other diseases of prostate (severe)

co-prescrire avec amikacin

Acute pyelonephritis (severe)

co-prescrire avec clarithromycin

Bacterial pneumonia (Community-acquired pneumonia - severe)

co-prescrire avec metronidazole

Peritoneal abscess (mild-moderate)

co-prescrire avec metronidazole

Peritonitis (mild-moderate)

co-prescrire avec metronidazole

Peritonitis (severe)

co-prescrire avec metronidazole

Peritoneal abscess (severe)

Second choix

Acute pyelonephritis (mild to moderate)

Osteomyelitis or osteitis

Bacterial infection of joint

Sepsis without septic shock

Inflammatory and other diseases of prostate (mild to moderate)

Autres indications

Bacterial infection of unspecified site

Ceftaroline

Section

Reserve group antibiotics

Parenteral > General injections > IV: 400 mg in vial (as fosamil) powder for injection; 600 mg in vial (as fosamil) powder for injection

Indications

Other specified bacterial diseases

Ceftazidime

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 250 mg in vial (as pentahydrate) powder for injection; 1 g in vial (as pentahydrate) powder for injection

Indications

Premier choix

co-prescrire avec [vancomycin](#)

[Endophthalmitis](#)

Autres indications

[Pseudomonas aeruginosa resistant to other antibiotic](#)

Ceftazidime + avibactam

Section

Reserve group antibiotics

Parenteral > General injections > IV: 2 g in vial + 0.5 g in vial powder for injection

Indications

[Carbapenem-resistant Pseudomonas aeruginosa](#)

[Carbapenem resistant Enterobacterales](#)

Ceftolozane + tazobactam

Section

Reserve group antibiotics

Parenteral > General injections > IV: 1 g + 0.5 g powder for injection

Indications

[Carbapenem-resistant Pseudomonas aeruginosa](#)

Ceftriaxone

Section

Watch group antibiotics

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 2 g in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt)

Parenteral > General injections > unspecified: 2 g in vial powder for injection (as sodium salt); 500 mg in vial powder for injection (as sodium salt); 250 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injection (as sodium salt)

Indications

Premier choix

[Other specified pneumonia \(Hospital-acquired pneumonia\)](#)

[Paratyphoid fever](#)

[Typhoid fever](#)

[Bacterial pneumonia \(Community-acquired pneumonia - severe\) \[children\]](#)

[Bacterial meningitis](#)

co-prescrire avec [amikacin](#)

[Inflammatory and other diseases of prostate \(severe\)](#)

co-prescrire avec [amikacin](#)

[Acute pyelonephritis \(severe\)](#)

co-prescrire avec [azithromycin](#)

[Gonococcal infection](#)

co-prescrire avec [clarithromycin](#)

[Bacterial pneumonia \(Community-acquired pneumonia - severe\)](#)

co-prescrire avec [metronidazole](#)

[Peritonitis \(severe\)](#)

co-prescrire avec [metronidazole](#)

[Peritonitis \(mild-moderate\)](#)

co-prescrire avec [metronidazole](#)

[Peritoneal abscess \(mild-moderate\)](#)

co-prescrire avec [metronidazole](#)

[Necrotising fasciitis](#)

co-prescrire avec [metronidazole](#)

[Peritoneal abscess \(severe\)](#)

co-prescrire avec [vancomycin](#)

[Endophthalmitis](#)

Second choix

[Acute pyelonephritis \(mild to moderate\)](#)

[Osteomyelitis or osteitis](#)

[Inflammatory and other diseases of prostate \(mild to moderate\)](#)

[Bacterial infection of joint](#)

[Sepsis without septic shock](#)

[Infectious gastroenteritis or colitis without specification of infectious agent](#)

Autres indications

[Bacterial infection of unspecified site](#)

Cefuroxime

Section

Watch group antibiotics

Parenteral > General injections > IV: 250 mg in vial (as sodium salt) powder for injection; 750 mg in vial (as sodium salt) powder for injection; 1.5 g in vial (as sodium salt) powder for injection

Indications

Second choix

[Other specified prophylactic measures](#)

Cemiplimab

Section

Immunomodulators

Indications

Therapeutic equivalent to [pembrolizumab](#) pour

[Other specified malignant neoplasms of bronchus or lung](#)

Certolizumab pegol

Section	Indications
Immunomodulators for non-malignant disease	Therapeutic equivalent to adalimumab pour Axial spondyloarthritis Therapeutic equivalent to adalimumab pour Crohn disease site Therapeutic equivalent to adalimumab pour Rheumatoid arthritis
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab pour Juvenile idiopathic arthritis

Cetirizine

Section	Indications
Antiallergics and medicines used in anaphylaxis	Therapeutic equivalent to loratadine pour Allergic or hypersensitivity conditions of unspecified type

Chlorambucil

Section	Indications
Cytotoxic medicines Oral > Solid: 2 mg tablet	Unspecified malignant neoplasms of ill-defined or unspecified sites Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Chloramphenicol

Section	Indications
Access group antibiotics Parenteral > General injections > IV: 1 g in vial powder for injection (sodium succinate) Parenteral > General injections > IM: 500 mg in 2 mL ampoule oily suspension for injection Oral > Liquid: 150 mg per 5 mL (as palmitate) Oral > Solid: 250 mg capsule	Second choix Bacterial meningitis Autres indications Bacterial infection of unspecified site

Chlorhexidine

Section	Indications
Medicines administered to the neonate [c] Local > Topical > Solution: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine) Local > Topical > Gel: 7.1% chlorhexidine digluconate (delivering 4% chlorhexidine) Antiseptics Local > Topical > Solution: 5% (digluconate) for dilution	Omphalitis of newborn Chlorhexidine

Chlorine base compound

Section	Indications
Disinfectants Local > Topical > Powder: 0.1% available chlorine for solution Local > Topical > Liquid: 0.1% available chlorine for solution Local > Topical > Solid: 0.1% available chlorine for solution	Disinfectant, not elsewhere classified

Chloroquine

Section	Indications
Antimalarial medicines > For chemoprevention Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) Oral > Solid: 150 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax Malaria due to Plasmodium ovale Malaria due to Plasmodium malariae
Antimalarial medicines > For curative treatment Parenteral > General injections > IV: 40 mg per mL in 5 mL ampoule (as hydrochloride, phosphate or sulfate) Oral > Liquid: 50 mg per 5 mL syrup (as phosphate or sulfate) Oral > Solid: 150 mg tablet (as phosphate or sulfate); 100 mg tablet (as phosphate or sulfate)	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)	Rheumatoid arthritis

Chlorothiazide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chloroxylonol

Section	Indications
Disinfectants Local > Topical > Solution: 4.8%	Phenol disinfectant, not elsewhere classified

Chlorpromazine

Section	Indications
Medicines for psychotic disorders Oral > Liquid: 25 mg per 5 mL (hydrochloride) Oral > Solid: 100 mg (hydrochloride); 10 mg (hydrochloride); 25 mg (hydrochloride); 50 mg (hydrochloride) Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (hydrochloride) Oral > Solid: 100 mg (hydrochloride)	Schizophrenia or other primary psychotic disorders Therapeutic equivalent to haloperidol pour Schizophrenia or other primary psychotic disorders

Chlortalidone

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Chlortetracycline

Section

Ophthalmological preparations > Anti-infective agents

Indications

Therapeutic equivalent to tetracycline pour

Other specified conjunctivitis

Therapeutic equivalent to tetracycline pour

Infectious keratitis

Therapeutic equivalent to tetracycline pour

Infectious blepharitis

Therapeutic equivalent to tetracycline pour

Trachoma

Cholera vaccine

Section

Recommendations for some high-risk populations

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against cholera alone

Ciclesonide

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to budesonide pour

Asthma

Ciclosporin

Section

Immunomodulators for non-malignant disease

Parenteral > General injections > IV: 50 mg per mL in 1 mL ampoule concentrate for injection

Oral > Liquid: 100 mg per mL

Oral > Solid > capsule: 25 mg

Indications

Failure or rejection of transplanted organs or tissues

Cimetidine

Section

Antiulcer medicines

Oral > Solid: 200 mg

Parenteral > General injections > unspecified: 200 mg in 2 mL ampoule

Indications

Peptic ulcer, site unspecified

Gastro-oesophageal reflux disease

Ciprofloxacin

Section

Watch group antibiotics

Parenteral > General injections > IV: 2 mg per mL (as hyclate)

Oral > Liquid: 250 mg per 5 mL (anhydrous)

Oral > Solid: 250 mg (as hydrochloride)

Oral > Solid: 250 mg (as hydrochloride); 500 mg (as hydrochloride)

Oral > Solid > dosage form: 250 mg (as hydrochloride); 500 mg (as hydrochloride); 100 mg (as hydrochloride)

Ear, nose and throat medicines [c]

Local > Otological > drops: 0.3% (as hydrochloride)

Indications

Premier choix

Paratyphoid fever

Typhoid fever

Acute pyelonephritis (mild to moderate)

Neutropenia (low-risk)

Infectious gastroenteritis or colitis without specification of infectious agent

Inflammatory and other diseases of prostate (mild to moderate)

Second choix

Cholera [children]

Cholera

co-prescrire avec metronidazole

Peritoneal abscess (mild-moderate)

co-prescrire avec metronidazole

Peritonitis (mild-moderate)

Autres indications

Bacterial infection of unspecified site

Infectious diseases of external ear

Cisplatin

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 10 mg in vial powder for injection; 50 mg in vial powder for injection

Parenteral > General injections > IV: 50 mg per 50 mL; 100 mg per 100 mL; 10 mg per 10 mL; 20 mg per 20 mL

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Germ cell tumour of testis

Osteosarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of lip, oral cavity or pharynx

Malignant neoplasms of cervix uteri

Cladribine

Section

Medicines for multiple sclerosis

Oral > Solid > tablet: 10 mg

Cytotoxic medicines

Parenteral > General injections > IV: 1 mg per mL in 10 mL vial; 2 mg per mL in 5 mL vial

Indications

Multiple sclerosis

Langerhans cell histiocytosis

Clarithromycin

Section

Watch group antibiotics

Oral > Liquid: 125 mg per 5 mL powder for oral liquid; 250 mg per 5 mL powder for oral liquid

Parenteral > General injections > unspecified: 500 mg in vial powder for injection

Oral > Solid > dosage form: 500 mg; 250 mg

Oral > Solid: 500 mg

Indications

Premier choix

co-prescrire avec cefotaxime

Bacterial pneumonia (Community-acquired pneumonia - severe)

co-prescrire avec ceftriaxone

Bacterial pneumonia (Community-acquired pneumonia - severe)

Second choix

Acute pharyngitis

co-prescrire avec amoxicillin + clavulanic acid

Bacterial pneumonia (Community-acquired pneumonia - severe)

Clindamycin

Section

Access group antibiotics

Oral > Liquid: 75 mg per 5 mL (as palmitate hydrochloride)

Oral > Solid: 150 mg (as hydrochloride) capsule

Parenteral > General injections > unspecified: 150 mg per mL (as phosphate) injection

Parenteral > General injections > IV: 150 mg per mL (as phosphate); 600 mg per 4 mL (as phosphate); 900 mg per 6 mL (as phosphate)

Oral > Liquid: 75 mg per 5 mL powder for oral liquid (as palmitate hydrochloride)

Oral > Solid: 150 mg (as hydrochloride)

Indications

Premier choix

co-prescrire avec piperacillin + tazobactam

Necrotising fasciitis

Second choix

Osteomyelitis or osteitis

Bacterial infection of joint

Autres indications

Bacterial infection of unspecified site

Clofazimine

Section	Indications
Antituberculosis medicines Oral > Solid: 100 mg; 50 mg	Multi-drug resistant Mycobacterium tuberculosis
Antileprosy medicines Oral > Solid: 100 mg; 50 mg	Leprosy

Clomifene

Section	Indications
Ovulation inducers Oral > Solid: 50 mg (citrate)	Female infertility without specification whether primary or secondary

Clomipramine

Section	Indications
Medicines for obsessive compulsive disorders Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride)	Obsessive-compulsive disorder

Clonazepam

Section	Indications
Antiseizure medicines Oral > Solid: 500 µg	Generalised myoclonic seizure

Clopidogrel

Section	Indications
Anti-platelet medicines Oral > Solid: 75 mg; 300 mg	Acute ischaemic heart disease Presence of coronary angioplasty implant or graft

Clotrimazole

Section	Indications
Antifungal medicines Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet	Vulvovaginal candidosis

Cloxacillin

Section

Access group antibiotics

Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid

Oral > Solid: 500 mg (as sodium salt) capsule; 1 g (as sodium salt) capsule

Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection

Oral > Liquid: 125 mg per 5 mL (as sodium salt) powder for oral liquid; 250 mg per 5 mL (as sodium salt) powder for oral liquid

Parenteral > General injections > unspecified: 500 mg in vial (as sodium salt) powder for injection; 250 mg in vial (as sodium salt) powder for injection

Oral > Solid > capsule: 500 mg (as sodium salt); 1 g (as sodium salt); 250 mg (as sodium salt)

Indications

Premier choix

Bacterial cellulitis, erysipelas or lymphangitis

Bacterial infection of joint

Osteomyelitis or osteitis

Second choix

co-prescrire avec amikacin

Sepsis without septic shock

Autres indications

Bacterial infection of unspecified site

Clozapine

Section

Medicines for psychotic disorders

Oral > Solid: 25 mg; 50 mg; 100 mg; 200 mg

Indications

Schizophrenia or other primary psychotic disorders

Coagulation factor IX

Section

Blood coagulation factors

Parenteral > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection

Indications

Coagulation defects, purpura or other haemorrhagic or related conditions

Haemophilia B

Coagulation factor IX complex

Section

Blood coagulation factors

Indications

Therapeutic equivalent to coagulation factor IX pour

Coagulation defects, purpura or other haemorrhagic or related conditions

Therapeutic equivalent to coagulation factor IX pour

Haemophilia B

Coagulation factor VIII

Section

Blood coagulation factors

Parenteral > General injections > IV: 500 IU in vial powder for injection;
250 IU in vial powder for injection; 1000 IU in vial powder for injection

Indications

Haemophilia A

Coal tar

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Local > Topical > Solution: 5%

Indications

Psoriasis of unspecified type

Cobicistat + elvitegravir + emtricitabine + tenofovir disoproxil fumarate

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 150 mg + 150 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Cobicistat + elvitegravir + emtricitabine+ tenofovir alafenamide

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 150 mg + 150 mg + 200 mg + 10 mg

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Cobimetinib

Section

Targeted therapies

Indications

Therapeutic equivalent to trametinib pour

Other specified melanoma of skin

Codeine

Section

Opioid analgesics

Oral > Solid: 30 mg tablet (codeine phosphate)

Antitussives

Oral > Solid: 10 mg tablet (codeine phosphate)

Medicines for diarrhoea

Oral > Solid: 30 mg tablet (codeine phosphate)

Indications

Pain

Cough

Diarrhoea

Colchicine

Section

Medicines used to treat gout

Oral > Solid: 0.5 mg

Indications

Gout

Colecalciferol

Section

Vitamins and minerals

Oral > Liquid: 400 IU per mL

Oral > Solid: 400 IU; 1000 IU

Indications

Calcium deficiency

Vitamin D deficiency

Therapeutic equivalent to [ergocalciferol](#) pour

Vitamin D deficiency

Therapeutic equivalent to [ergocalciferol](#) pour

Calcium deficiency

Colistin (injection)

Section

Reserve group antibiotics

Parenteral > General injections > unspecified: 1 million IU in vial (as colisthemethate sodium)

Parenteral > General injections > unspecified: 1 million IU in vial (as colisthemethate sodium) (equivalent to 34 mg colistin base activity)

Indications

Other specified bacterial diseases

Carbapenem resistant *Acinetobacter baumannii*

Carbapenem-resistant *Pseudomonas aeruginosa*

Carbapenem resistant Enterobacterales

Compound sodium lactate solution

Section

Solutions correcting water, electrolyte and acid-base disturbances >

Parenteral

Parenteral > General injections > IV: injectable solution

Indications

Other specified disorders of fluid, electrolyte or acid-base balance

Hypovolaemic shock

Condoms

Section

Barrier methods

-

Indications

Contact with health services for contraceptive management

Copper-containing intrauterine device

Section

Intrauterine devices

-

Indications

Contact with health services for insertion of contraceptive device

Crizotinib

Section

Targeted therapies

Oral > Solid: 200 mg; 250 mg

Oral > Solid > capsule: 200 mg; 250 mg

Indications

Other specified malignant neoplasms of bronchus or lung

Anaplastic large cell lymphoma, ALK-positive

Cryoprecipitate (pathogen-reduced)

Section

Blood and blood components

Injection: frozen liquid in bag or lyophilized powder in vial containing:

> 50 IU Factor VIII

> 100 IU vWF

> 140 mg clottable fibrinogen per unit

Indications

Coagulation defects, purpura or other haemorrhagic or related conditions

Von Willebrand disease

Obstetric haemorrhage

Cyanocobalamin

Section

Antianaemia medicines

.

Indications

Anaemias or other erythrocyte disorders

Cyclizine

Section

Medicines for other common symptoms in palliative care

Oral > Solid: 50 mg tablet

Parenteral > General injections > unspecified: 50 mg per mL

Indications

Palliative care

Cyclopentolate

Section	Indications
Diagnostic agents > Ophthalmic medicines	Therapeutic equivalent to tropicamide pour Cycloplegic drug
Ophthalmological preparations > Mydriatics	Therapeutic equivalent to atropine pour Anterior uveitis

Cyclophosphamide

Section	Indications
Cytotoxic medicines Oral > Solid: 25 mg tablet Parenteral > General injections > unspecified: 500 mg in vial powder for injection Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection Oral > Solid: 25 mg; 50 mg Oral > Solid: 25 mg; 50 mg tablet	Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of breast Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Chronic lymphocytic leukaemia or small lymphocytic lymphoma Diffuse large B-cell lymphomas Plasma cell myeloma Other specified gliomas of brain Other specified malignant neoplasms of breast Malignant trophoblastic neoplasms of placenta Follicular lymphoma Malignant neoplasms of kidney, except renal pelvis Rhabdomyosarcoma primary site Hodgkin lymphoma Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive

Cycloserine

Section	Indications
Antituberculosis medicines Oral > Solid: 250 mg; 125 mg	Multi-drug resistant Mycobacterium tuberculosis

Cytarabine

Section

Cytotoxic medicines

Parenteral > General injections > unspecified: 100 mg in vial powder for injection

Parenteral > General injections > unspecified: 100 mg in vial powder for injection; 100 mg per mL in vial

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Acute myeloid leukaemia with recurrent genetic abnormalities

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Dabigatran

Section

Medicines affecting coagulation

Oral > Solid: 110 mg; 150 mg

Indications

Venous thromboembolism

Other specified atrial fibrillation

Dabrafenib

Section

Targeted therapies

Oral > Solid: 50 mg; 75 mg

Indications

Other specified melanoma of skin

Dacarbazine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 100 mg in vial powder for injection; 200 mg in vial powder for injection

Indications

Hodgkin lymphoma

Daclatasvir

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations

Oral > Solid: 30 mg tablet (as hydrochloride); 60 mg tablet (as hydrochloride)

Indications

Chronic hepatitis C

Daclatasvir + sofosbuvir

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations
Oral > Solid: 60 mg + 400 mg

Indications

Chronic hepatitis C

Dactinomycin

Section

Cytotoxic medicines
Parenteral > General injections > IV: 500 µg in vial powder for injection

Indications

Malignant trophoblastic neoplasms of placenta

Unspecified malignant neoplasms of ill-defined or unspecified sites

Malignant neoplasms of kidney, except renal pelvis

Rhabdomyosarcoma primary site

Ewing sarcoma of bone and articular cartilage of unspecified sites

Dalteparin

Section

Medicines affecting coagulation

Indications

Therapeutic equivalent to [enoxaparin](#) pour

Acute ischaemic heart disease

Therapeutic equivalent to [enoxaparin](#) pour

Venous thromboembolism

Dapagliflozin

Section

Oral hypoglycaemic agents

Indications

Therapeutic equivalent to [empagliflozin](#) pour

Type 2 diabetes mellitus

Dapsone

Section

Antileprosy medicines
Oral > Solid: 100 mg tablet; 50 mg tablet; 25 mg tablet

Indications

Leprosy

Daptomycin

Section

Reserve group antibiotics

Parenteral > General injections > IV: 350 mg in vial powder for injection; 500 mg in vial powder for injection

Indications

Other specified bacterial diseases

Daratumumab

Section

Immunomodulators

Parenteral > General injections > IV: 100 mg per 5 mL; 400 mg per 20 mL

Indications

Plasma cell myeloma

Darbepoetin alfa

Section

Antianaemia medicines

Indications

Therapeutic equivalent to erythropoiesis-stimulating agents pour

Anaemia due to chronic disease

Darunavir

Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 75 mg; 400 mg; 600 mg; 800 mg

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Dasabuvir

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

Oral > Solid: 250 mg tablet

Indications

Chronic hepatitis C

Dasatinib

Section

Targeted therapies

Oral > Solid: 20 mg tablet; 50 mg tablet; 70 mg tablet; 80 mg tablet; 100 mg tablet; 140 mg tablet

Indications

Chronic myeloid leukaemia, not elsewhere classified

Daunorubicin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as hydrochloride) powder for injection; 20 mg in vial (as hydrochloride) powder for injection; 2 mg per mL in vial (as hydrochloride); 5 mg per mL in vial (as hydrochloride)

Indications

Lymphoid leukaemia, not elsewhere classified

Myeloid leukaemia

Acute myeloid leukaemia with recurrent genetic abnormalities

Deferasirox

Section

Other medicines for haemoglobinopathies

Oral > Solid > dispersible tablet: 100 mg; 125 mg; 250 mg; 400 mg; 500 mg

Oral > Solid > tablet: 90 mg; 180 mg; 360 mg

Indications

Therapeutic equivalent to [deferoxamine](#) pour

Thalassaemias

Other specified sickle cell disorders or other haemoglobinopathies

Thalassaemias

Deferoxamine

Section

Antidotes and other substances used in poisonings > Specific

Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)

Other medicines for haemoglobinopathies

Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)

Indications

Harmful effects of drugs, medicaments or biological substances, not elsewhere classified

Thalassaemias

Other specified sickle cell disorders or other haemoglobinopathies

Delafloxacin

Section

Watch group antibiotics

Parenteral > General injections > IV: 300 mg lyophilized powder for injection

Oral > Solid: 450 mg

Indications

Methicillin resistant *Staphylococcus aureus*

Delamanid

Section

Antituberculosis medicines

Oral > Solid: 50 mg tablet; 25 mg tablet (dispersible)

Indications

Multi-drug resistant *Mycobacterium tuberculosis*

Dengue vaccine

Section

Recommendations for some high-risk populations

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against certain single viral diseases

Desmopressin

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 4 µg per mL in 1 mL ampoule (acetate)

Local > Nasal > Spray: 10 µg per dose (acetate)

Indications

Haemophilia A

Von Willebrand disease

Dexamethasone

Section

Other medicines administered to the mother

Parenteral > General injections > unspecified: 4 mg per mL (dexamethasone phosphate (as sodium))

Hormones and antihormones

Oral > Solid: 4 mg

Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

Oral > Liquid: 2 mg per 5 mL

Oral > Solid: 2 mg; 4 mg

Antiallergics and medicines used in anaphylaxis

Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per 5 mL

Oral > Solid: 2 mg; 4 mg

Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

Antiemetic medicines

Oral > Liquid: 0.5 mg per 5 mL; 2 mg per 5 mL

Oral > Solid: 0.5 mg; 0.75 mg; 1.5 mg; 4 mg

Parenteral > General injections > unspecified: 4 mg per mL in 1 mL ampoule (as disodium phosphate salt)

Indications

Respiratory distress syndrome of the newborn, altered by maternal corticosteroid therapy

Plasma cell myeloma

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Allergic or hypersensitivity conditions of unspecified type

Palliative care

Nausea or vomiting

Dextran 40

Section

Plasma substitutes

.

Indications

Hypovolaemia

Dextran 70

Section

Plasma substitutes

Parenteral > General injections > IV: 6% injectable solution

Indications

Hypovolaemia

Diaphragms

Section

Barrier methods

-

Indications

Contact with health services for contraceptive management

Diazepam

Section

Antiseizure medicines

Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system

Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube

Medicines for other common symptoms in palliative care

Oral > Liquid: 2 mg per 5 mL

Oral > Solid: 5 mg; 10 mg

Local > Rectal > Gel: 5 mg per mL in 0.5 mL rectal delivery system; 5 mg per mL in 2 mL rectal delivery system; 5 mg per mL in 4 mL rectal delivery system

Local > Rectal > Solution: 2 mg per mL in 1.25 mL rectal tube; 2 mg per mL in 2.5 mL rectal tube; 4 mg per mL in 2.5 mL rectal tube

Parenteral > General injections > unspecified: 5 mg per mL

Medicines for anxiety disorders

Oral > Solid: 5 mg (scored); 2 mg (scored)

Indications

Therapeutic equivalent to lorazepam pour

Status epilepticus

Status epilepticus

Palliative care

Anxiety

Diazoxide

Section	Indications
Medicines for hypoglycaemia Oral > Liquid: 50 mg per mL Oral > Solid: 50 mg	Persistent hyperinsulinaemic hypoglycaemia of infancy

Didanosine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 100 mg buffered powder for oral solution; 167 mg buffered powder for oral solution; 250 mg buffered powder for oral solution Oral > Solid: 25 mg buffered chewable dispersible tablet; 50 mg buffered chewable dispersible tablet; 100 mg buffered chewable dispersible tablet; 150 mg buffered chewable dispersible tablet; 200 mg buffered chewable dispersible tablet; 125 mg unbuffered enteric coated capsule; 200 mg unbuffered enteric coated capsule; 250 mg unbuffered enteric coated capsule; 400 mg unbuffered enteric coated capsule	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Diethylcarbamazine

Section	Indications
Antifilarials Oral > Solid: 50 mg tablet (dihydrogen citrate); 100 mg tablet (dihydrogen citrate)	Lymphatic filariasis

Digitoxin

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 0.2 mg in 1 mL ampoule Oral > Solid: 50 µg; 100 µg	Heart failure

Digoxin

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid: 62.5 µg; 250 µg

Medicines used in heart failure

Parenteral > General injections > IV: 250 µg per mL in 2 mL ampoule; 100 µg per mL in 1 mL ampoule

Oral > Liquid: 50 µg per mL

Oral > Solid > tablet: 62.5 µg; 250 µg; 125 µg

Indications

Cardiac arrhythmia

Heart failure

Dihydroartemisinin + piperazine phosphate

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 40 mg + 320 mg tablet; 20 mg + 160 mg tablet

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Dihydroergocryptine mesylate

Section

Antiseizure medicines

Oral > Solid:

Indications

Parkinson disease

Diloxanide

Section

Antiamoebic and anti giardiasis medicines

Oral > Solid: 500 mg (furoate)

Indications

Amoebiasis

Dimercaprol

Section

Antidotes and other substances used in poisonings > Specific

Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Diphtheria antitoxin

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies

Parenteral > General injections > IM: 10000 IU in vial; 20000 IU in vial

Indications

Need for immunization against diphtheria alone

Diphtheria vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against diphtheria alone

Diphtheria-pertussis-tetanus vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against diphtheria-tetanus-pertussis, combined

Diphtheria-tetanus vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against combinations of infectious diseases

Docetaxel

Section

Cytotoxic medicines

Parenteral > General injections > IV: 20 mg per mL; 40 mg per mL

Indications

Other specified malignant neoplasms of breast

Malignant neoplasms of breast

Malignant neoplasms of prostate

Unspecified malignant neoplasms of ill-defined or unspecified sites

Docusate sodium

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 50 mg per 5 mL oral liquid

Oral > Solid: 100 mg capsule

Indications

Palliative care

Dolasetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Dolutegravir

Section	Indications
Antiretrovirals > Integrase inhibitors Oral > Solid: 50 mg tablet; 10 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Dolutegravir + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Donepezil

Section	Indications
Medicines for diseases of the nervous system Oral > Liquid: 1 mg per mL (hydrochloride) Oral > Solid > tablet: 5 mg (hydrochloride); 10 mg (hydrochloride); 5 mg (hydrochloride) (orodispersible); 10 mg (hydrochloride) (orodispersible)	Dementia due to Alzheimer disease

Dopamine

Section	Indications
Medicines used in heart failure Parenteral > General injections > IV: 40 mg per mL in 5 mL vial (hydrochloride)	Heart failure Cardiogenic shock

Doxorubicin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride)

Parenteral > General injections > IV: 10 mg in vial (hydrochloride); 50 mg in vial (hydrochloride); 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Parenteral > General injections > IV: 50 mg in vial (hydrochloride) +; 2 mg per mL in 5 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride); 10 mg in vial (hydrochloride)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Rhabdomyosarcoma primary site

Follicular lymphoma

Burkitt lymphoma including Burkitt leukaemia

Ewing sarcoma of bone and articular cartilage of unspecified sites

Malignant neoplasms of breast

Lymphoid leukaemia, not elsewhere classified

Other specified malignant neoplasms of breast

Diffuse large B-cell lymphomas

Plasma cell myeloma

Malignant neoplasms of kidney, except renal pelvis

Osteosarcoma of bone and articular cartilage of unspecified sites

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Doxorubicin (as pegylated liposomal)

Section

Cytotoxic medicines

Parenteral > General injections > IV: 2 mg per mL in 10 mL vial (hydrochloride); 2 mg per mL in 25 mL vial (hydrochloride)

Indications

Kaposi sarcoma of unspecified primary site

Doxycycline

Section

Access group antibiotics

Oral > Solid: 100 mg (as hyclate)

Parenteral > General injections > unspecified: 100 mg in vial powder for injection

Oral > Liquid: 25 mg per 5 mL (anhydrous); 50 mg per 5 mL (anhydrous)

Oral > Solid: 100 mg (as hyclate); 50 mg (as hyclate)

Oral > Liquid: 25 mg per 5 mL powder for oral liquid (monohydrate); 50 mg per 5 mL (calcium)

Oral > Solid: 50 mg (as hyclate); 100 mg (as hyclate)

Oral > Solid > dispersible tablet: 100 mg (as monohydrate)

Antimalarial medicines > For curative treatment

Oral > Solid: 100 mg (hydrochloride or hyclate); 100 mg dispersible tablet (as monohydrate)

Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg (hydrochloride or hyclate)

Indications

Premier choix

Cholera

Chlamydia trachomatis

Second choix

Bacterial pneumonia (Community-acquired pneumonia - mild to moderate)

Cholera [children]

Chronic obstructive pulmonary disease with acute exacerbation

Autres indications

Bacterial infection of unspecified site

Malaria due to Plasmodium falciparum

Malaria due to Plasmodium falciparum

Durvalumab

Section

Immunomodulators

Parenteral > General injections > IV: 120 mg per 2.4 mL in vial; 500 mg per 10 mL in vial

Indications

Other specified malignant neoplasms of bronchus or lung

Edoxaban

Section

Medicines affecting coagulation

Indications

Therapeutic equivalent to dabigatran pour

Venous thromboembolism

Therapeutic equivalent to dabigatran pour

Other specified atrial fibrillation

Efavirenz

Section

Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors

Oral > Solid: 600 mg tablet

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Efavirenz + emtricitabine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 600 mg + 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Efavirenz + lamivudine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate, equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Therapeutic equivalent to efavirenz + emtricitabine + tenofovir pour

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Eflornithine

Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)

Indications

African trypanosomiasis

Elbasvir + grazoprevir

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

Oral > Solid: 50 mg + 100 mg

Indications

Chronic hepatitis C

Empagliflozin

Section

Oral hypoglycaemic agents

Oral > Solid: 10 mg; 25 mg

Indications

Type 2 diabetes mellitus

Emtricitabine

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

Oral > Liquid: 10 mg per mL

Oral > Solid: 200 mg

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir alafenamide

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 200 mg + 25 mg + 25 mg

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + rilpivirine + tenofovir disoproxil fumarate

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 200 mg + 25 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Emtricitabine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 200 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Contact with or exposure to human immunodeficiency virus

Emtricitabine + tenofovir alafenamide

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 200 mg + 10 mg; 200 mg + 25 mg

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Enalapril

Section

Antihypertensive medicines

Oral > Liquid: 1 mg per mL (as hydrogen maleate)

Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)

Medicines used in heart failure

Oral > Solid > tablet: 2.5 mg (as hydrogen maleate); 5 mg (as hydrogen maleate); 10 mg (as hydrogen maleate)

Indications

Essential hypertension

Heart failure

Encorafenib

Section

Targeted therapies

Indications

Therapeutic equivalent to [dabrafenib](#) pour

Other specified melanoma of skin

Enoxaparin

Section

Medicines affecting coagulation

Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 120 mg per 0.8 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule

Parenteral > General injections > SC: 20 mg per 0.2 mL in prefilled syringe; 40 mg per 0.4 mL in prefilled syringe; 60 mg per 0.6 mL in prefilled syringe; 80 mg per 0.8 mL in prefilled syringe; 100 mg per 1 mL in prefilled syringe; 120 mg per 0.8 mL in prefilled syringe; 150 mg per 1 mL in prefilled syringe; 150 mg per 1 mL in ampoule; 20 mg per 0.2 mL in ampoule; 40 mg per 0.4 mL in ampoule; 60 mg per 0.6 mL in ampoule; 80 mg per 0.8 mL in ampoule; 120 mg per 0.8 mL in ampoule; 100 mg per 1 mL in ampoule

Indications

Acute ischaemic heart disease

Venous thromboembolism

Entecavir

Section

Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors

Oral > Liquid: 0.05 mg per mL

Oral > Solid: 0.5 mg; 1 mg

Indications

Chronic hepatitis B

Enzalutamide

Section

Hormones and antihormones

Oral > Solid: 40 mg

Indications

Therapeutic equivalent to [abiraterone](#) pour

Malignant neoplasms of prostate

Malignant neoplasms of prostate

Ephedrine

Section

Local anaesthetics

Parenteral > Locoregional injections > Spinal anaesthesia: 30 mg per mL in 1 mL ampoule (ephedrine hydrochloride)

Indications

Cardiac complications of anaesthesia during labour or delivery

Epinephrine

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 100 µg per mL in 10 mL ampoule (as acid tartrate or hydrochloride)

Ophthalmological preparations > Mydriatics

Local > Ophthalmological > Solution (eye drops): 2% (as hydrochloride)

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Parenteral > General injections > unspecified: 1 mg per mL in ampoule (as hydrochloride or hydrogen tartrate)

Antiallergics and medicines used in anaphylaxis

Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule (as hydrochloride or hydrogen tartrate)

Indications

Cardiac arrest

Primary open-angle glaucoma

Asthma

Chronic obstructive pulmonary disease

Anaphylaxis

Epoetin alfa

Section

Antianaemia medicines

Indications

Therapeutic equivalent to erythropoiesis-stimulating agents pour

Anaemia due to chronic disease

Epoetin beta

Section

Antianaemia medicines

Indications

Therapeutic equivalent to erythropoiesis-stimulating agents pour

Anaemia due to chronic disease

Epoetin theta

Section

Antianaemia medicines

Indications

Therapeutic equivalent to erythropoiesis-stimulating agents pour

Anaemia due to chronic disease

Equine rabies immunoglobulin

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies
Parenteral > Locoregional injections > Intradermal: 150 IU per mL in vial;
200 IU per mL in vial; 300 IU per mL in vial; 400 IU per mL in vial

Indications

Rabies

Eravacycline

Section

Reserve group antibiotics
Parenteral > General injections > IV: 50 mg lyophilized powder for
injection

Indications

Carbapenem resistant Enterobacterales

Ergocalciferol

Section

Vitamins and minerals
Oral > Liquid: 250 µg per mL (10 000 IU per mL)
Oral > Solid: 1.25 mg (50 000 IU)

Indications

Therapeutic equivalent to [colecalfiferol](#) pour

Calcium deficiency

Therapeutic equivalent to [colecalfiferol](#) pour

Vitamin D deficiency

Vitamin D deficiency

Calcium deficiency

Ergometrine

Section

Uterotonics
Parenteral > General injections > unspecified: 200 µg in 1 mL ampoule
(hydrogen maleate)

Indications

Postpartum haemorrhage

Erlotinib

Section

Targeted therapies
Oral > Solid: 100 mg; 150 mg

Indications

Other specified malignant neoplasms of
bronchus or lung

Erythromycin

Section

Watch group antibiotics

Parenteral > General injections > IV: 500 mg in vial powder for injection (as lactobionate)

Oral > Liquid: 125 mg per 5 mL (as stearate or ethylsuccinate)

Oral > Solid: 250 mg (as stearate or ethylsuccinate)

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Ointment: 0.5%

Indications

Second choix

Therapeutic equivalent to [clarithromycin](#) pour

Acute pharyngitis

Autres indications

Bacterial infection of unspecified site

Chlamydial conjunctivitis

Gonococcal infection of eye

Erythropoiesis-stimulating agents

Section

Antianaemia medicines

Parenteral > General injections > unspecified: 1000 IU per 0.5 mL pre-filled syringe; 2000 IU per 0.5 mL pre-filled syringe; 3000 IU per 0.3 mL pre-filled syringe; 4000 IU per 0.4 mL pre-filled syringe; 5000 IU per 0.5 mL pre-filled syringe; 6000 IU per 0.6 mL pre-filled syringe; 8000 IU per 0.8 mL pre-filled syringe; 10000 IU per 1 mL pre-filled syringe; 20000 IU per 0.5 mL pre-filled syringe; 40000 IU per 1 mL pre-filled syringe

Indications

Anaemia due to chronic disease

Estradiol

Section

Estrogens

Oral > Solid > tablet: 0.5 mg; 1 mg; 2 mg

Indications

Hypopituitarism

Gonadotropin deficiency

Turner syndrome

Estradiol cypionate + medroxyprogesterone acetate

Section

Injectable hormonal contraceptives

Parenteral > General injections > IM: 5 mg + 25 mg

Indications

Contact with health services for reasons associated with reproduction

Etanercept

Section

Immunomodulators for non-malignant disease

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) pour

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) pour

Crohn disease site

Therapeutic equivalent to [adalimumab](#) pour

Rheumatoid arthritis

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

Ethambutol

Section

Antituberculosis medicines

Parenteral > General injections > IV: 1000 mg per 10 mL; 2000 mg per 20 mL

Oral > Solid: 100 mg (hydrochloride); 400 mg (hydrochloride); 100 mg tablet (dispersible)

Indications

Other specified tuberculosis

Tuberculosis

Ethambutol + isoniazid

Section

Antituberculosis medicines

Oral > Solid: 400 mg + 150 mg tablet

Indications

Tuberculosis

Ethambutol + isoniazid + pyrazinamide + rifampicin

Section

Antituberculosis medicines

Oral > Solid: 275 mg + 75 mg + 400 mg + 150 mg tablet

Indications

Tuberculosis

Ethambutol + isoniazid + rifampicin

Section

Antituberculosis medicines

Oral > Solid: 275 mg + 75 mg + 150 mg tablet

Indications

Tuberculosis

Ethanol

Section

Antiseptics

Local > Topical > Solution: 70% (denatured)

Indications

Denatured alcohol

Ether

Section

General anaesthetics and oxygen > Inhalational medicines

Respiratory > Inhalation > solution:

Indications

Anaesthetics and therapeutic gases

Ethinylestradiol + etonogestrel

Section

Intravaginal contraceptives

Local > Vaginal > vaginal ring: 2.7 mg + 11.7 mg

Indications

Contact with health services for contraceptive management

Ethinylestradiol + levonorgestrel

Section

Oral hormonal contraceptives

Oral > Solid: 50 µg + 250 µg [4]

Oral > Solid: 30 µg + 150 µg

Indications

Contact with health services for postcoital contraception

Contact with health services for contraceptive management

Ethinylestradiol + norethisterone

Section

Oral hormonal contraceptives

Oral > Solid: 35 µg + 1 mg

Indications

Contact with health services for contraceptive management

Ethionamide

Section

Antituberculosis medicines

Oral > Solid: 250 mg tablet; 125 mg tablet (dispersible)

Indications

Multi-drug resistant Mycobacterium tuberculosis

Tuberculous meningitis

Ethosuximide

Section	Indications
Antiseizure medicines Oral > Liquid: 250 mg per 5 mL Oral > Solid: 250 mg	Absence seizures, typical

Etonogestrel-releasing implant

Section	Indications
Implantable contraceptives Implant > Subdermal: 68 mg single rod	Contact with health services for insertion of contraceptive device

Etoposide

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule Oral > Solid: 100 mg capsule Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule; 100 mg in vial powder for injection (as phosphate) Oral > Solid: 100 mg capsule; 50 mg capsule	Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Myeloid leukaemia Other specified malignant neoplasms of the ovary Other specified malignant neoplasms of bronchus or lung Germ cell tumour of testis Malignant trophoblastic neoplasms of placenta Malignant neoplasms of kidney, except renal pelvis Osteosarcoma of bone and articular cartilage of unspecified sites Hodgkin lymphoma Retinoblastoma Anaplastic large cell lymphoma, ALK-negative Anaplastic large cell lymphoma, ALK-positive

Everolimus

Section	Indications
Targeted therapies Oral > Solid: 2.5 mg; 5 mg; 7.5 mg; 10 mg; 2 mg tablet (dispersible); 3 mg tablet (dispersible); 5 mg tablet (dispersible)	Other specified gliomas of brain

Fentanyl

Section

Opioid analgesics

Local > Topical > Transdermal patch: 12 µg/hour; 25 µg/hour; 50 µg/hour; 75 µg/hour; 100 µg/hour

Oral > Solid: 100 to 800 µg sub-lingual tablet (as citrate); 200 to 1600 µg lozenge (as citrate)

Oral > Other: 200 to 1200 µg buccal film (as citrate)

Indications

Chronic cancer pain

Other specified chronic cancer related pain

Ferrous salt

Section

Antianaemia medicines

Oral > Liquid: 25 mg per mL iron (as sulfate)

Oral > Solid: 60 mg iron

Indications

Iron deficiency anaemia

Ferrous salt + folic acid

Section

Antianaemia medicines

Oral > Solid: 60 mg iron + 2.8 mg

Oral > Solid > tablet: 60 mg elemental iron + 400 µg tablet; 60 mg elemental iron + 2.8 mg

Indications

Iron deficiency anaemia

Clefts of lip, alveolus or palate

Iron deficiency anaemia

Anencephaly or similar anomalies

Spina bifida

Fexinidazole

Section

Medicines for the treatment of 1st stage African trypanosomiasis

Oral > Solid: 600 mg

Medicines for the treatment of 2nd stage African trypanosomiasis

Oral > Solid: 600 mg

Indications

African trypanosomiasis

African trypanosomiasis

Fexofenadine

Section

Antiallergics and medicines used in anaphylaxis

Indications

Therapeutic equivalent to loratadine pour

Allergic or hypersensitivity conditions of unspecified type

Fifth generation cephalosporins

Section

Reserve group antibiotics

Indications

Other specified bacterial diseases

Filgrastim

Section

Immunomodulators

Parenteral > General injections > SC: 120 µg per 0.2 mL in prefilled syringe; 300 µg per 0.5 mL in prefilled syringe; 480 µg per 0.8 mL in prefilled syringe; 300 µg per mL in 1 mL vial; 480 µg per 1.6 mL in 1.6 mL vial

Indications

Acquired neutropaenia

Fingolimod

Section

Immunomodulators for non-malignant disease

Oral > Solid: 0.25 mg; 0.5 mg

Indications

Relapsing-remitting multiple sclerosis

Flomoxef

Section

Watch group antibiotics

Parenteral > General injections > IV: 0.5 g in vial powder for injection; 1 g in vial powder for injection

Indications

Other specified other antibiotic resistant Enterobacterales

Fluconazole

Section

Antifungal medicines

Parenteral > General injections > IV: 2 mg per mL in vial

Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid

Oral > Solid > capsule: 50 mg

Indications

Cryptococcosis

Candidosis

Flucytosine

Section

Antifungal medicines

Parenteral > General injections > IV: 2.5 g per 250 mL infusion

Oral > Solid: 250 mg capsule

Indications

Cryptococcosis

Fludarabine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (fludarabine phosphate) powder for injection

Oral > Solid: 10 mg tablet (fludarabine phosphate)

Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Fludrocortisone

Section

Adrenal hormones and synthetic substitutes

Oral > Solid: 100 µg (acetate)

Indications

Adrenocortical insufficiency

Flunisolide

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to [budesonide](#) pour

Asthma

Fluorescein

Section

Diagnostic agents > Ophthalmic medicines

Local > Ophthalmological > Solution (eye drops): 1% (sodium salt)

Indications

Fluorescein

Fluoride

Section

Dental medicines and preparations

Local > Dental > Gel: 2500 to 12500 ppm fluoride (any type)

Local > Dental > Toothpaste, cream or gel: 1000 to 1500 ppm fluoride (any type)

Local > Dental > Mouthrinse: 230 to 900 ppm fluoride (any type)

Local > Dental > Varnish: 22500 ppm fluoride (any type)

Indications

Dental caries

Fluorouracil

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Local > Topical > Ointment: 5%

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg per mL in 5 mL ampoule

Parenteral > General injections > IV: 50 mg per mL in vial

Indications

Plantar warts

Unspecified malignant neoplasms of ill-defined or unspecified sites

Malignant neoplasms of nasopharynx

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of breast

Malignant neoplasms of rectum

Malignant neoplasms of colon

Fluoxetine

Section

Medicines for depressive disorders

Oral > Solid: 20 mg (as hydrochloride)

Medicines for anxiety disorders

Oral > Solid: 20 mg (as hydrochloride)

Medicines for other common symptoms in palliative care

Oral > Solid: 20 mg (as hydrochloride)

Medicines for obsessive compulsive disorders

Oral > Solid: 20 mg (as hydrochloride)

Indications

Depressive disorders

Social anxiety disorder

Generalised anxiety disorder

Panic disorder

Palliative care

Obsessive-compulsive disorder

Fluphenazine

Section

Medicines for psychotic disorders

Parenteral > General injections > IM: 25 mg per mL in ampoule (decanoate); 25 mg per mL in ampoule (enanthate)

Indications

Schizophrenia or other primary psychotic disorders

Flutamide

Section

Hormones and antihormones

Indications

Therapeutic equivalent to bicalutamide pour

Malignant neoplasms of prostate

Fluticasone

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to [budesonide](#) pour

Asthma

Fluticasone + formoterol

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to [budesonide + formoterol](#) pour

Asthma

Fluticasone furoate + vilanterol

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to [budesonide + formoterol](#) pour

Asthma

Fluvastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Folic acid

Section

Antianaemia medicines

Oral > Solid: 1 mg tablet; 5 mg tablet

Oral > Solid: 400 µg

Indications

Folate deficiency anaemia

Anencephaly or similar anomalies

Spina bifida

Cephalocele

Fomepizole

Section

Antidotes and other substances used in poisonings > Specific
Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule (sulfate); 1 g per mL in 1.5 mL ampoule (base)

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Fosfomycin (injection)

Section

Reserve group antibiotics
Parenteral > General injections > IV: 2 g in vial (as sodium) powder for injection; 4 g in vial (as sodium) powder for injection

Indications

Other specified bacterial diseases

Carbapenem resistant *Acinetobacter baumannii*

Carbapenem-resistant *Pseudomonas aeruginosa*

Carbapenem resistant Enterobacterales

Fourth generation cephalosporins

Section

Watch group antibiotics
.

Indications

Other specified bacterial diseases

Fresh-frozen plasma

Section

Blood and blood components
Parenteral > General injections > IV:

Indications

Blood transfusion without reported diagnosis

Fulvestrant

Section

Hormones and antihormones
Parenteral > General injections > IV: 250 mg per 5 mL

Indications

Malignant neoplasms of breast

Furosemide

Section

Medicines used in heart failure

Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL

Oral > Solid > tablet: 40 mg; 20 mg

Diuretics

Parenteral > General injections > IV: 10 mg per mL in 2 mL ampoule; 10 mg per mL in 5 mL ampoule

Oral > Liquid: 20 mg per 5 mL; 50 mg per 5 mL

Oral > Solid > tablet: 40 mg; 20 mg

Indications

Heart failure

Anuria or oliguria

Oedema

Gabapentin

Section

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral > Solid: 100 mg; 200 mg; 300 mg; 400 mg; 600 mg; 800 mg

Indications

Neuropathic pain

Gallamine

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

Parenteral > General injections > IV: 40 mg per mL in 2 mL ampoule (gallamine triethiodide)

Indications

Muscle relaxants

Gatifloxacin

Section

Antituberculosis medicines

Oral > Solid: 200 mg; 400 mg

Indications

Multi-drug resistant *Mycobacterium tuberculosis*

Gefitinib

Section

Targeted therapies

Indications

Therapeutic equivalent to erlotinib pour

Other specified malignant neoplasms of bronchus or lung

Gemcitabine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 200 mg in vial powder for injection; 1 g in vial powder for injection

Indications

Other specified carcinomas of ovary

Other specified malignant neoplasms of bronchus or lung

Gentamicin

Section

Access group antibiotics

Parenteral > General injections > unspecified: 10 mg per mL in 2 mL vial (as sulfate); 40 mg per mL in 2 mL vial (as sulfate)

Indications

Premier choix

Neonatal meningitis

co-prescrire avec amoxicillin

Sepsis without septic shock

co-prescrire avec amoxicillin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec ampicillin

Peritoneal abscess (mild-moderate)

co-prescrire avec ampicillin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec ampicillin

Peritoneal abscess (severe)

co-prescrire avec ampicillin

Peritonitis (severe)

co-prescrire avec ampicillin

Sepsis without septic shock

co-prescrire avec ampicillin

Peritonitis (mild-moderate)

co-prescrire avec ampicillin

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec benzylpenicillin

Bacterial pneumonia (Community-acquired pneumonia - severe) [children]

co-prescrire avec benzylpenicillin

Acute malnutrition in infants, children or adolescents (complicated) [children]

co-prescrire avec benzylpenicillin

Sepsis without septic shock

Second choix

Other specified prophylactic measures

Gonococcal infection

Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 0.3% (gentamicin sulfate)	Autres indications
	Bacterial infection of unspecified site
	Other specified conjunctivitis
	Infectious blepharitis

Glass ionomer cement

Section	Indications
Dental medicines and preparations Local > Dental > Powder + liquid: Single-use capsules: 0.4 g powder + 0.09 mL liquid; Multi-use bottle: powder + liquid	Dental caries

Glatiramer acetate

Section	Indications
Medicines for multiple sclerosis Parenteral > General injections > SC: 20 mg per mL; 40 mg per mL Parenteral > General injections > SC: 20 mg per mL in pre-filled syringe; 40 mg per mL in pre-filled syringe	Relapsing-remitting multiple sclerosis Multiple sclerosis

Glecaprevir + pibrentasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid: 100 mg + 40 mg tablet; 50 mg + 20 mg granules	Chronic hepatitis C

Glibenclamide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 5 mg; 2.5 mg	Type 2 diabetes mellitus

Gliclazide

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release	Type 2 diabetes mellitus

Glucagon

Section

Medicines for hypoglycaemia

Parenteral > General injections > unspecified: 1 mg per mL

Indications

Hypoglycaemia in the context of diabetes

Glucose

Section

Solutions correcting water, electrolyte and acid-base disturbances >

Parenteral

Parenteral > General injections > IV: 5% isotonic; 10% hypertonic

Parenteral > General injections > IV: 50% hypertonic

Indications

Other specified disorders of fluid, electrolyte or acid-base balance

Hypoglycaemia without associated diabetes

Glucose + sodium chloride

Section

Solutions correcting water, electrolyte and acid-base disturbances >

Parenteral

Parenteral > General injections > IV: 4% + 0.18% (equivalent to Na⁺ 30 mmol/L and Cl⁻ 30 mmol/L; 5% + 0.9% (equivalent to Na⁺ 150 mmol/L and Cl⁻ 150 mmol/L); 5% + 0.45% (equivalent to Na⁺ 75 mmol/L and Cl⁻ 75 mmol/L)

Indications

Other specified disorders of fluid, electrolyte or acid-base balance

Glutaral

Section

Disinfectants

Local > Topical > Solution: 2%

Indications

Glutaral nonmedicinal

Glyceryl trinitrate

Section

Antianginal medicines

Oral > Solid: 500 µg sublingual tablet

Indications

Angina pectoris

Glycopyrronium

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to tiotropium bromide pour

Chronic obstructive pulmonary disease

Golimumab

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) pour

[Axial spondyloarthritis](#)

Therapeutic equivalent to [adalimumab](#) pour

[Crohn disease site](#)

Therapeutic equivalent to [adalimumab](#) pour

[Rheumatoid arthritis](#)

Medicines for juvenile joint diseases

Therapeutic equivalent to [adalimumab](#) pour

[Juvenile idiopathic arthritis](#)

Goserelin

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [leuporelin](#) pour

[Malignant neoplasms of breast](#)

Therapeutic equivalent to [leuporelin](#) pour

[Malignant neoplasms of prostate](#)

Granisetron

Section

Medicines for other common symptoms in palliative care

Indications

Therapeutic equivalent to [ondansetron](#) pour

[Palliative care](#)

Antiemetic medicines

Therapeutic equivalent to [ondansetron](#) pour

[Nausea or vomiting](#)

Griseofulvin

Section

Antifungal medicines

Oral > Liquid: 125 mg per 5 mL

Oral > Solid: 125 mg; 250 mg

Indications

[Dermatophytosis](#)

Haemophilus influenzae type b vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against single bacterial diseases

Haloperidol

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 2 mg per mL Oral > Solid: 0.5 mg; 2 mg; 5 mg Parenteral > General injections > unspecified: 5 mg per mL in 1-mL ampoule	Palliative care
Medicines for psychotic disorders Parenteral > General injections > IM: 5 mg per mL in ampoule Oral > Solid: 2 mg; 5 mg	Schizophrenia or other primary psychotic disorders

Haloperidol decanoate

Section	Indications
Medicines for psychotic disorders	Therapeutic equivalent to fluphenazine pour Schizophrenia or other primary psychotic disorders

Halothane

Section	Indications
General anaesthetics and oxygen > Inhalational medicines Respiratory > Inhalation > liquid:	Anaesthetics and therapeutic gases

Heparin sodium

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 1000 IU per mL in 5 mL ampoule; 20000 IU per mL in 5 mL ampoule; 5000 IU per mL in 5 mL ampoule	Acute ischaemic heart disease Unstable angina Venous thromboembolism

Hepatitis a vaccine

Section

Recommendations for some high-risk populations

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against viral hepatitis

Hepatitis b vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against viral hepatitis

Homatropine

Section

Ophthalmological preparations > Mydriatics

Local > Ophthalmological > Solution (eye drops): 2% (hydrobromide)

Indications

Anterior uveitis

Therapeutic equivalent to *atropine* pour

Anterior uveitis

Human papilloma virus (HPV) vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against certain single viral diseases

Hydralazine

Section

Antihypertensive medicines

Parenteral > General injections > IV: 20 mg in ampoule powder for injection (hydrochloride)

Oral > Solid: 50 mg tablet; 25 mg tablet (hydrochloride)

Oral > Solid: 50 mg (hydrochloride); 25 mg (hydrochloride)

Indications

Essential hypertension

Gestational hypertension

Hydrochlorothiazide

Section

Antihypertensive medicines

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg; 12.5 mg

Medicines used in heart failure

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 25 mg

Diuretics

Oral > Solid: 25 mg

Indications

Essential hypertension

Heart failure

Oedema

Hydrocortisone

Section

Hormones and antihormones

Parenteral > General injections > unspecified: 100 mg in vial (as sodium succinate)

Adrenal hormones and synthetic substitutes

Oral > Solid: 5 mg; 10 mg; 20 mg

Ophthalmological preparations > Anti-inflammatory agents

Local > Ophthalmological > Ointment: 1% (acetate)

Antiallergics and medicines used in anaphylaxis

Parenteral > General injections > unspecified: 100 mg in vial powder for injection (as sodium succinate)

Anti-inflammatory medicines

Local > Rectal > Suppository: 25 mg (acetate)

Local > Rectal > Retention enema: 100 mg per 60 mL

Dermatological medicines > Anti-inflammatory and antipruritic medicines

Local > Topical > Cream: 1% (acetate)

Local > Topical > Ointment: 1% (acetate)

Indications

Lymphoid leukaemia, not elsewhere classified

Unspecified malignant neoplasms of ill-defined or unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Adrenocortical insufficiency

Scleritis

Anterior uveitis

Anaphylaxis

Crohn disease site

Ulcerative colitis

Dermatitis or eczema

Lichen planus

Pruritus due to skin disorder

Psoriasis of unspecified type

Hydromorphone

Section

Opioid analgesics

Indications

Therapeutic equivalent to [morphine](#) pour

Pain

Hydroxocobalamin

Section

Antianaemia medicines

Parenteral > General injections > IM: 1 mg per mL in 1 mL ampoule (as acetate); 1 mg per mL in 1 mL ampoule (as hydrochloride); 1 mg per mL in 1 mL ampoule (as sulfate)

Indications

Megaloblastic anaemia due to vitamin B12 deficiency

Hydroxycarbamide (hydroxyurea)

Section

Cytotoxic medicines

Oral > Solid: 200 mg; 250 mg; 300 mg; 400 mg; 500 mg; 1 g

Oral > Solid > dosage form: 200 mg; 300 mg; 400 mg; 500 mg; 1 g; 100 mg

Other medicines for haemoglobinopathies

Oral > Solid > dosage form: 200 mg; 500 mg; 1 g; 100 mg

Indications

Squamous cell carcinoma of oropharynx

Chronic myeloid leukaemia, not elsewhere classified

Other specified sickle cell disorders or other haemoglobinopathies

Hydroxychloroquine

Section

Antimalarial medicines > For chemoprevention

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 200 mg (as sulfate)

Indications

Therapeutic equivalent to [chloroquine](#) pour

Malaria due to *Plasmodium falciparum*

Systemic lupus erythematosus

Cutaneous lupus erythematosus of unspecified type

Hyoscine butylbromide

Section

Medicines for other common symptoms in palliative care

Parenteral > General injections > unspecified: 20 mg per mL injection

Indications

Palliative care

Hyoscine hydrobromide

Section

Medicines for other common symptoms in palliative care

Local > Topical > Transdermal patch: 1 mg per 72 hours transdermal patch

Parenteral > General injections > unspecified: 400 µg per mL injection; 600 µg per mL injection

Indications

Palliative care

Hypochlorous acid

Section

Antiseptics

Local > Topical > Solution:

Indications

Disinfectant, not elsewhere classified

Hypromellose

Section

Ophthalmological preparations

Local > Ophthalmological > Solution (eye drops): 0.3%

Indications

Keratoconjunctivitis sicca

Ibrutinib

Section

Targeted therapies

Oral > Solid: 140 mg

Indications

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Ibuprofen

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 5 mg per mL

Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs)

Oral > Liquid: 200 mg per 5 mL; 100 mg per 5 mL

Oral > Solid > tablet: 200 mg; 400 mg; 600 mg

Antimigraine medicines > For treatment of acute attack

Oral > Liquid: 100 mg per 5 mL; 200 mg per 5 mL

Oral > Solid > tablet: 200 mg; 400 mg

Indications

Patent arterial duct

Pain

Migraine

Idoxuridine

Section

Ophthalmological preparations > Anti-infective agents
Local > Ophthalmological > Solution (eye drops): 0.1%
Local > Ophthalmological > Ointment: 0.2%

Indications

Herpes simplex keratitis

Ifosfamide

Section

Cytotoxic medicines
Parenteral > General injections > IV: 500 mg in vial powder for injection; 1 g in vial powder for injection; 2 g in vial powder for injection
Parenteral > General injections > IV: 1 g in vial powder for injection; 2 g in vial powder for injection

Indications

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Ewing sarcoma of bone and articular cartilage of unspecified sites

Osteosarcoma of bone and articular cartilage of unspecified sites

Rhabdomyosarcoma primary site

Unspecified malignant neoplasms of ill-defined or unspecified sites

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of kidney, except renal pelvis

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Imatinib

Section

Targeted therapies
Oral > Solid: 100 mg; 400 mg

Indications

B lymphoblastic leukaemia or lymphoma with t(9:22) (q34;q11.2); BCR-ABL1

Gastrointestinal stromal tumour of unspecified gastrointestinal sites

Chronic myeloid leukaemia, not elsewhere classified

Imipenem + cilastatin

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 250 mg (as monohydrate) + 250 mg (as sodium salt) powder for injection; 500 mg (as monohydrate) + 500 mg (as sodium salt) powder for injection	Second choix Therapeutic equivalent to meropenem pour Neutropenia (high-risk) Therapeutic equivalent to meropenem pour Peritoneal abscess (severe) Therapeutic equivalent to meropenem pour Peritonitis (severe)
Antituberculosis medicines	Autres indications Pseudomonas aeruginosa resistant to other antibiotic Antibiotic resistant Acinetobacter baumannii Therapeutic equivalent to meropenem pour Multi-drug resistant Mycobacterium tuberculosis

Imipenem + cilastatin + relebactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 500 mg + 500 mg + 250 mg powder for injection	Other specified other antibiotic resistant Enterobacterales Carbapenem resistant Enterobacterales Carbapenem-resistant Pseudomonas aeruginosa

Indapamide

Section	Indications
Antihypertensive medicines	Therapeutic equivalent to hydrochlorothiazide pour Essential hypertension
Medicines used in heart failure	Therapeutic equivalent to hydrochlorothiazide pour Heart failure

Indinavir

Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 400 mg (as sulfate)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Indometacin

Section

Medicines administered to the neonate [c]

Indications

Therapeutic equivalent to [ibuprofen](#) pour

Patent arterial duct

Infliximab

Section

Immunomodulators for non-malignant disease

Indications

Therapeutic equivalent to [adalimumab](#) pour

Axial spondyloarthritis

Therapeutic equivalent to [adalimumab](#) pour

Crohn disease site

Therapeutic equivalent to [adalimumab](#) pour

Rheumatoid arthritis

Medicines for juvenile joint diseases

Therapeutic equivalent to [adalimumab](#) pour

Juvenile idiopathic arthritis

Influenza vaccine (seasonal)

Section

Recommendations for immunization programmes with certain characteristics

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against influenza

Insulin

Section

Insulins

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial (soluble); 100 IU per mL in 10 mL vial (soluble); 100 IU per mL in 3 mL cartridge (soluble); 100 IU per mL in 3 mL pre-filled pen (soluble)

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial (soluble); 100 IU per mL in 10 mL vial (soluble); 100 IU per mL in 3 mL cartridge (soluble); 100 IU per mL in 3 mL pre-filled pen

Indications

Type 2 diabetes mellitus

Type 1 diabetes mellitus

Insulin analogues

Section

Insulins

Not specified

Indications

Type 1 diabetes mellitus

Type 2 diabetes mellitus

Insulin degludec

Section

Insulins

Parenteral > General injections > SC: 100 units per mL

Indications

Therapeutic equivalent to [insulin glargine](#) pour

Type 1 diabetes mellitus

Therapeutic equivalent to [long-acting insulin analogues](#) pour

Type 1 diabetes mellitus

Therapeutic equivalent to [long-acting insulin analogues](#) pour

Type 2 diabetes mellitus

Insulin detemir

Section

Insulins

Indications

Therapeutic equivalent to [insulin glargine](#) pour

Type 1 diabetes mellitus

Therapeutic equivalent to [long-acting insulin analogues](#) pour

Type 1 diabetes mellitus

Therapeutic equivalent to [long-acting insulin analogues](#) pour

Type 2 diabetes mellitus

Insulin glargine

Section

Insulins

Parenteral > General injections > SC: 100 units per mL

Indications

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 1 diabetes mellitus

Therapeutic equivalent to long-acting insulin analogues pour

Type 2 diabetes mellitus

Intermediate-acting insulin

Section

Insulins

Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin

Indications

Type 2 diabetes mellitus

Intraperitoneal dialysis solution

Section

Peritoneal dialysis solution
parenteral solution

Indications

Care involving peritoneal dialysis

Iodine

Section

Vitamins and minerals

Parenteral > General injections > IM: 0.5 mL (240 mg iodine) iodized oil in ampoule; 1 mL (480 mg iodine) iodized oil in ampoule

Oral > Liquid: 1 mL (480 mg iodine) iodized oil in ampoule; 0.5 mL (240 mg iodine) iodized oil in ampoule; 0.57 mL (308 mg iodine) iodized oil in dispenser bottle

Oral > Solid: 190 mg

Antiseptics

Indications

Iodine deficiency

Therapeutic equivalent to povidone iodine pour

Povidone iodine

Iohexol

Section

Diagnostic agents > Radiocontrast media

Parenteral > General injections > unspecified: 140 to 350 mg iodine per mL in 5 mL ampoule; 140 to 350 mg iodine per mL in 10 mL ampoule; 140 to 350 mg iodine per mL in 20 mL ampoule

Indications

X-ray contrast media, iodinated

Ipecacuanha

Section

Antidotes and other substances used in poisonings > Non-specific

Oral > Liquid: 0.14% ipecacuanha alkaloids calculated as emetine

Indications

Unspecified injury, poisoning or certain other consequences of external causes

Ipratropium bromide

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Respiratory > Inhalation > aerosol: 20 µg per dose

Indications

Asthma

Chronic obstructive pulmonary disease

Irinotecan

Section

Cytotoxic medicines

Parenteral > General injections > IV: 40 mg per 2 mL in 2 mL vial; 100 mg per 5 mL in 5 mL vial; 500 mg per 25 mL in 25 mL vial

Indications

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of kidney, except renal pelvis

Rhabdomyosarcoma primary site

Isoflurane

Section

General anaesthetics and oxygen > Inhalational medicines

Respiratory > Inhalation > liquid:

Indications

Anaesthetics and therapeutic gases

Isoniazid

Section

Antituberculosis medicines

Parenteral > General injections > IV: 100 mg per mL

Oral > Solid: 100 mg tablet; 100 mg tablet (dispersible); 300 mg tablet

Indications

Other specified tuberculosis

Tuberculosis

Isoniazid + pyrazinamide + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid: 50 mg + 150 mg + 75 mg tablet (dispersible)	Tuberculosis

Isoniazid + pyridoxine + sulfamethoxazole + trimethoprim

Section	Indications
Medicines for prevention of HIV-related opportunistic infections Oral > Solid: 300 mg + 25 mg + 800 mg + 160 mg tablet (scored)	Other specified prophylactic measures

Isoniazid + rifampicin

Section	Indications
Antituberculosis medicines Oral > Solid: 50 mg + 75 mg tablet (dispersible); 75 mg + 150 mg tablet; 150 mg + 300 mg tablet	Tuberculosis

Isoniazid + rifapentine

Section	Indications
Antituberculosis medicines Oral > Solid: 300 mg + 300 mg tablet (scored)	Latent tuberculosis

Isoprenaline

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 µg per mL in ampoule (hydrochloride) Oral > Solid: 10 mg (hydrochloride or sulfate); 15 mg (hydrochloride or sulfate)	Bradycardia Cardiogenic shock

Isosorbide dinitrate

Section	Indications
Antianginal medicines Oral > Solid: 5 mg sublingual tablet	Angina pectoris

Itraconazole

Section

Antifungal medicines

Oral > Liquid: 10 mg per mL

Oral > Solid: 100 mg

Indications

Chronic pulmonary aspergillosis

Chromoblastomycosis

Histoplasmosis

Talaromycosis

Sporotrichosis

Paracoccidioidomycosis

Ivermectin

Section

Intestinal anthelmintics

Oral > Solid > tablet: 3 mg

Indications

Strongyloidiasis

Ascariasis

Trichuriasis

Hookworm diseases

Ancylostomiasis

Antifilarials

Oral > Solid > tablet: 3 mg

Onchocerciasis

Lymphatic filariasis

Medicines for ectoparasitic infections

Oral > Solid > tablet: 3 mg

Scabies

Kanamycin

Section

Ophthalmological preparations > Anti-infective agents

Indications

Therapeutic equivalent to [gentamicin](#) pour

Other specified conjunctivitis

Therapeutic equivalent to [gentamicin](#) pour

Infectious blepharitis

Kanamycin (injection)

Section

Antituberculosis medicines

Parenteral > General injections > IV: 1000 mg powder for injection (vial)

Indications

Multi-drug resistant *Mycobacterium tuberculosis*

Ketamine

Section

General anaesthetics and oxygen > Injectable medicines

Parenteral > General injections > unspecified: 50 mg per mL in 10 mL vial (as hydrochloride)

Indications

Anaesthetics and therapeutic gases

Ketoconazole

Section

Medicines for endocrine disorders

Oral > Solid > tablet: 200 mg

Indications

Cushing syndrome

Lactulose

Section

Medicines for other common symptoms in palliative care

Oral > Liquid: 3.1 to 3.7 g per 5 mL oral liquid

Indications

Palliative care

Lamivudine

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

Oral > Liquid: 50 mg per 5 mL

Oral > Solid: 150 mg tablet

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamivudine + nevirapine + zidovudine

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 30 mg + 50 mg + 60 mg Tablet; 150 mg + 200 mg + 300 mg Tablet

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamivudine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

Indications

Therapeutic equivalent to emtricitabine + tenofovir pour

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Therapeutic equivalent to emtricitabine + tenofovir pour

Contact with or exposure to human immunodeficiency virus

Lamivudine + zidovudine

Section

Fixed-dose combinations of antiretrovirals

Oral > Solid: 30 mg + 60 mg Tablet; 150 mg + 300 mg Tablet

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Lamotrigine

Section

Antiseizure medicines

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet; 200 mg tablet; 2 mg tablet (chewable, dispersible); 5 mg tablet (chewable, dispersible); 25 mg tablet (chewable, dispersible); 50 mg tablet (chewable, dispersible); 100 mg tablet (chewable, dispersible); 200 mg tablet (chewable, dispersible)

Indications

Epilepsy or seizures

Latanoprost

Section

Ophthalmological preparations > Miotics and antiglaucoma medicines

Local > Ophthalmological > Solution (eye drops): 50 µg per mL eye drops

Indications

Primary open-angle glaucoma

Ledipasvir + sofosbuvir

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations

Oral > Solid: 90 mg + 400 mg tablet

Indications

Chronic hepatitis C

Lenalidomide

Section

Immunomodulators

Oral > Solid: 25 mg

Indications

Plasma cell myeloma

Letrozole

Section

Ovulation inducers

Oral > Solid > dosage form: 2.5 mg

Indications

Female infertility without specification whether primary or secondary

Leuprorelin

Section

Hormones and antihormones

Parenteral > General injections > IM: 7.5 mg in prefilled syringe; 22.5 mg in prefilled syringe

Indications

Malignant neoplasms of breast

Malignant neoplasms of prostate

Levamisole

Section

Intestinal anthelmintics

Oral > Solid > tablet: 50 mg (as hydrochloride); 150 mg (as hydrochloride)

Indications

Ascariasis

Levetiracetam

Section

Antiseizure medicines

Oral > Liquid: 100 mg per mL

Oral > Solid > tablet: 250 mg; 500 mg; 750 mg; 1000 mg

Parenteral > General injections > IV: 500 mg per 5 mL in 5 mL vial; 5 mg per mL in 100 mL bag; 10 mg per mL in 100 mL bag; 15 mg per mL in 100 mL bag

Indications

Epilepsy or seizures

Status epilepticus

Levodopa

Section

Antiseizure medicines

Oral > Solid: 250 mg

Indications

Parkinson disease

Levodopa + benserazide

Section

Medicines for parkinsonism

Indications

Therapeutic equivalent to levodopa + carbidopa pour

Parkinson disease

Levodopa + carbidopa

Section

Medicines for parkinsonism

Oral > Solid: 100 mg + 10 mg tablet; 250 mg + 25 mg tablet; 100 mg + 25 mg tablet

Indications

Parkinson disease

Levofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 250 mg tablet; 500 mg tablet; 750 mg tablet; 100 mg tablet (dispersible)	Multi-drug resistant Mycobacterium tuberculosis

Levonorgestrel

Section	Indications
Oral hormonal contraceptives Oral > Solid: 750 µg [2]; 1.5 mg [1] Oral > Solid: 30 µg	Contact with health services for postcoital contraception Contact with health services for contraceptive management
Intrauterine devices Intrauterine system with reservoir containing 52 mg of levonorgestrel.	Contact with health services for insertion of contraceptive device

Levonorgestrel-releasing implant

Section	Indications
Implantable contraceptives Implant > Subdermal: 75 mg per rod (two-rods)	Contact with health services for insertion of contraceptive device

Levothyroxine

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 50 µg (sodium salt); 100 µg (sodium salt); 25 µg (sodium salt)	Hypothyroidism

Lidocaine

Section	Indications
Antiarrhythmic medicines Parenteral > General injections > IV: 20 mg per mL in 5 mL ampoule (hydrochloride)	Ventricular tachyarrhythmia
Local anaesthetics Parenteral > Locoregional injections > Spinal anaesthesia: 5% in 2 mL ampoule (hydrochloride) + 7.5% glucose solution Parenteral > Locoregional injections > Other: 1% in vial (hydrochloride); 2% in vial (hydrochloride) Local > Topical > unspecified: 2 to 4% (hydrochloride)	Local anaesthetics

Lidocaine + epinephrine

Section	Indications
Local anaesthetics Parenteral > Locoregional injections > Other: 1% (lidocaine hydrochloride OR lidocaine sulfate) + 1:200 000 epinephrine in vial; 2% (lidocaine hydrochloride OR lidocaine sulfate) + 1:200 000 epinephrine in vial Local > Dental > Cartridge: 2% (lidocaine hydrochloride) + 1:80 000 epinephrine	Local anaesthetics

Lindane

Section	Indications
Dermatological medicines > Scabicides and pediculicides Local > Topical > Cream: 1% Local > Topical > Lotion: 1%	Scabies

Linezolid

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 2 mg per mL in 300 mL bag Oral > Liquid: 100 mg per 5 mL powder for oral liquid Oral > Solid: 400 mg tablet; 600 mg tablet Oral > Solid > dispersible tablet: 150 mg Oral > Solid > tablet: 600 mg	Other specified bacterial diseases Methicillin resistant Staphylococcus aureus Vancomycin resistant Staphylococcus aureus Vancomycin resistant Enterococcus
Antituberculosis medicines Oral > Solid: 600 mg tablet; 150 mg tablet (dispersible)	Multi-drug resistant Mycobacterium tuberculosis

Liraglutide

Section	Indications
Medicines for endocrine disorders Parenteral > General injections > SC: 6 mg per mL in 3 mL pre-filled pen	Obesity

Lisinopril + amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 5 mg; 20 mg + 5 mg; 20 mg + 10 mg	Essential hypertension

Lisinopril + hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Solid: 10 mg + 12.5 mg; 20 mg + 12.5 mg; 20 mg + 25 mg	Essential hypertension

Lisocabtagene maraleucel

Section	Indications
Antineoplastics and supportive medicines	Therapeutic equivalent to axicabtagene ciloleucel pour Diffuse large B-cell lymphomas

Lithium carbonate

Section	Indications
Medicines for bipolar disorders Oral > Solid: 300 mg	Bipolar or related disorders

Long-acting insulin analogues

Section	Indications
Insulins Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen	Type 1 diabetes mellitus Type 2 diabetes mellitus

Loperamide

Section	Indications
Medicines for other common symptoms in palliative care Oral > Solid: 2 mg solid oral dosage form	Palliative care

Lopinavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors Oral > Solid: 100 mg + 25 mg tablet (heat-stable); 200 mg + 50 mg tablet (heat-stable); 40 mg + 10 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Loratadine

Section	Indications
Antiallergics and medicines used in anaphylaxis Oral > Liquid: 1 mg per mL Oral > Solid: 10 mg	Allergic or hypersensitivity conditions of unspecified type

Lorazepam

Section

Antiseizure medicines

Parenteral > General injections > IV: 2 mg per mL in 1 mL ampoule; 4 mg per mL in 1 mL ampoule

Medicines for anxiety disorders

Indications

Status epilepticus

Therapeutic equivalent to [diazepam](#) pour

Anxiety

Losartan

Section

Antihypertensive medicines

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Medicines used in heart failure

Oral > Solid: 25 mg tablet; 50 mg tablet; 100 mg tablet

Indications

Essential hypertension

Heart failure

Lovastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Lugol's solution

Section

Thyroid hormones and antithyroid medicines

Oral > Liquid: 13% w/v (about 130 mg/mL total iodine)

Indications

Thyrotoxicosis

Magnesium sulfate

Section

Antiseizure medicines

Parenteral > General injections > IV: 500 mg per mL in 2 mL ampoule (equivalent to 1 g in 2 mL; 50% weight/volume)

Parenteral > General injections > IM: 500 mg per mL in 10 mL ampoule (equivalent to 5 g in 10 mL; 50% weight/volume)

Indications

Severe pre-eclampsia

Eclampsia

Mannitol

Section

Diuretics

Parenteral > General injections > IV: 10% solution; 20% solution

Indications

Ocular hypertension

Cerebral oedema

Measles vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against measles alone

Mebendazole

Section

Intestinal anthelmintics

Oral > Solid > tablet: 100 mg (chewable); 500 mg (chewable)

Indications

Ascariasis

Enterobiasis

Trichuriasis

Hookworm diseases

Cysticidal medicines

Oral > Solid > tablet: 500 mg (chewable); 100 mg (chewable)

Other specified echinococcosis

Medroxyprogesterone acetate

Section

Injectable hormonal contraceptives

Parenteral > General injections > IM: 150 mg per mL in 1 mL vial

Parenteral > General injections > SC: 104 mg per 0.65 mL in pre-filled syringe; 104 mg per 0.65 mL in single dose injection delivery system

Indications

Contact with health services for reasons associated with reproduction

Progestogens

Oral > Solid: 5 mg

Excessive menstruation with irregular cycle

Mefloquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 250 mg tablet (hydrochloride)

Indications

Malaria due to *Plasmodium falciparum*

Malaria due to *Plasmodium vivax*

Antimalarial medicines > For chemoprevention

Oral > Solid: 250 mg tablet (hydrochloride)

Malaria due to *Plasmodium falciparum*

Meglumine antimoniate

Section

Antileishmaniasis medicines

Parenteral > General injections > IM: 1.5 g per 5 mL in 5 mL ampoule

Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

Meglumine iotroxate

Section

Diagnostic agents > Radiocontrast media

Parenteral > General injections > IV: 5 to 8 g iodine (as meglumine) in 100 to 250 mL

Indications

X-ray contrast media, iodinated

Melarsoprol

Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)

Indications

African trypanosomiasis

Melphalan

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection

Oral > Solid: 2 mg

Indications

Plasma cell myeloma

Meningococcal meningitis vaccine

Section

Recommendations for some high-risk populations

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against single bacterial diseases

Mercaptopurine

Section

Cytotoxic medicines

Oral > Solid: 50 mg tablet

Oral > Liquid: 20 mg per mL

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Acute myeloid leukaemia with recurrent genetic abnormalities

Lymphoid leukaemia, not elsewhere classified

Langerhans cell histiocytosis

Meropenem

Section	Indications
Watch group antibiotics Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	Second choix Neutropenia (high-risk) Peritoneal abscess (severe) Peritonitis (severe) Neonatal meningitis
Antituberculosis medicines Parenteral > General injections > IV: 500 mg in vial (as trihydrate) powder for injection; 1 g in vial (as trihydrate) powder for injection	Multi-drug resistant Mycobacterium tuberculosis

Meropenem + vaborbactam

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 1 g in vial + 1 g in vial powder for injection	Carbapenem-resistant Pseudomonas aeruginosa Carbapenem resistant Enterobacterales Carbapenem resistant Acinetobacter baumannii

Mesalazine

Section	Indications
Anti-inflammatory medicines	Therapeutic equivalent to sulfasalazine pour Crohn disease site Therapeutic equivalent to sulfasalazine pour Ulcerative colitis

Mesna

Section	Indications
Supportive medicines Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule Oral > Solid: 400 mg; 600 mg	Osteosarcoma of bone and articular cartilage of other specified sites Other specified malignant neoplasms of the ovary Germ cell tumour of testis Ewing sarcoma of bone and articular cartilage of unspecified sites Rhabdomyosarcoma primary site Unspecified malignant neoplasms of ill-defined or unspecified sites Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis

Metformin

Section	Indications
Oral hypoglycaemic agents Oral > Solid: 500 mg (hydrochloride)	Type 2 diabetes mellitus

Methadone

Section	Indications
Medicines for opioid use disorders Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)	Opioid dependence
Opioid analgesics Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride) Oral > Solid: 5 mg (methadone hydrochloride); 10 mg (methadone hydrochloride)	Chronic cancer pain

Methimazole

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 5 mg; 10 mg; 20 mg	Thyrotoxicosis

Methionine

Section

Antidotes and other substances used in poisonings > Specific
Oral > Solid: 250 mg (DL)

Indications

Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Methotrexate

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium salt)

Oral > Solid: 2.5 mg (as sodium salt)

Parenteral > General injections > IV: 50 mg in vial (as sodium salt); 50 mg per 2 mL; 1000 mg per 10 mL concentrated injection

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Oral > Solid > tablet: 2.5 mg (as sodium salt); 10 mg (as sodium salt)

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Immunomodulators for non-malignant disease

Parenteral > General injections > SC: pre-filled syringe in multiple strengths

Medicines for juvenile joint diseases

Oral > Solid > tablet: 2.5 mg (as sodium salt)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Lymphoid leukaemia, not elsewhere classified

Acute myeloid leukaemia with recurrent genetic abnormalities

Malignant trophoblastic neoplasms of placenta

Burkitt lymphoma including Burkitt leukaemia

Malignant neoplasms of breast

Osteosarcoma of bone and articular cartilage of unspecified sites

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Psoriasis of unspecified type

Rheumatoid arthritis, serology unspecified

Crohn disease site

Psoriatic arthritis

Juvenile idiopathic arthritis

Methoxy polyethylene glycol-epoetin beta

Section

Antianaemia medicines

Indications

Therapeutic equivalent to erythropoiesis-stimulating agents pour

Anaemia due to chronic disease

Methyldopa

Section

Antihypertensive medicines

Oral > Solid: 250 mg

Indications

Essential hypertension

Gestational hypertension

Methylergometrine

Section

Uterotonics

Indications

Therapeutic equivalent to [ergometrine](#) pour

Postpartum haemorrhage

Methylphenidate

Section

Medicines for mental and behavioural disorders

Multiple formulations and strengths.

Indications

Attention deficit hyperactivity disorder

Methylprednisolone

Section

Hormones and antihormones

Parenteral > General injections > IV: 40 mg per mL in 1 mL single-dose vial (as sodium succinate); 40 mg per mL in 5 mL multi-dose vial (as sodium succinate); 80 mg per mL in 1 mL single-dose vial (as sodium succinate)

Indications

Lymphoid leukaemia, not elsewhere classified

Burkitt lymphoma including Burkitt leukaemia

Methylthioninium chloride

Section

Antidotes and other substances used in poisonings > Specific

Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule

Indications

Acquired methaemoglobinaemia

Metoclopramide

Section	Indications
Medicines for other common symptoms in palliative care Oral > Liquid: 5 mg per 5 mL Oral > Solid: 10 mg (hydrochloride) Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	Palliative care
Antiemetic medicines Oral > Liquid: 5 mg per 5 mL Oral > Solid: 10 mg (hydrochloride) Parenteral > General injections > unspecified: 5 mg per mL in 2 mL ampoule (hydrochloride)	Nausea or vomiting

Metoprolol

Section	Indications
Antianginal medicines	Therapeutic equivalent to bisoprolol pour Angina pectoris
Medicines used in heart failure	Therapeutic equivalent to bisoprolol pour Heart failure
Antihypertensive medicines	Therapeutic equivalent to bisoprolol pour Essential hypertension
Antiarrhythmic medicines	Therapeutic equivalent to bisoprolol pour Cardiac arrhythmia

Metronidazole

Section

Access group antibiotics

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Oral > Solid: 200 to 500 mg tablet

Parenteral > General injections > unspecified: 500 mg per 100 mL in vial

Injection

Local > Rectal > Suppository: 500 mg suppository; 1 g suppository

Antiamoebic and anti giardiasis medicines

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Parenteral > General injections > unspecified: 500 mg in 100 mL vial

Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg

Indications

Premier choix

co-prescrire avec cefazolin

Other specified prophylactic measures

co-prescrire avec cefotaxime

Peritoneal abscess (severe)

co-prescrire avec cefotaxime

Peritonitis (severe)

co-prescrire avec cefotaxime

Peritonitis (mild-moderate)

co-prescrire avec cefotaxime

Peritoneal abscess (mild-moderate)

co-prescrire avec ceftriaxone

Peritoneal abscess (severe)

co-prescrire avec ceftriaxone

Peritoneal abscess (mild-moderate)

co-prescrire avec ceftriaxone

Peritonitis (severe)

co-prescrire avec ceftriaxone

Peritonitis (mild-moderate)

Second choix

co-prescrire avec ciprofloxacin

Peritoneal abscess (mild-moderate)

co-prescrire avec ciprofloxacin

Peritonitis (mild-moderate)

Autres indications

Bacterial infection of unspecified site

Amoebiasis

Micafungin

Section

Antifungal medicines

Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection

Indications

Systemic or invasive candidosis

Miconazole

Section

[Dermatological medicines](#) > [Antifungal medicines](#)

Local > Topical > Cream: 2% (nitrate)

Local > Topical > Ointment: 2% (nitrate)

Indications

Fungal infection of the skin

Midazolam

Section

[Medicines for other common symptoms in palliative care](#)

Oral > Liquid: 2 mg per mL oral liquid

Oral > Solid: 7.5 mg solid oral dosage form; 15 mg solid oral dosage form

Parenteral > General injections > unspecified: 1 mg per mL injection; 5 mg per mL injection

[Antiseizure medicines](#)

Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial (for buccal administration); 5 mg per mL in 1 mL vial (for buccal administration); 5 mg per mL in 3 mL vial (for buccal administration)

Local > Buccal > Solution: 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration); 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration); 10 mg per mL in 0.75 mL pre-filled syringe; 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)

[Preoperative medication and sedation for short-term procedures](#)

Parenteral > General injections > IV: 1 mg per mL

Oral > Liquid: 2 mg per mL

Oral > Solid: 15 mg tablet; 7.5 mg tablet

Indications

Palliative care

Therapeutic equivalent to [lorazepam](#) pour

Status epilepticus

Status epilepticus

Sedative-hypnotic and anxiolytic drugs

Mifepristone - misoprostol

Section

[Uterotonics](#)

Oral > Solid: 200 mg + 200 µg; 200 mg [1] + 200 µg [4] in co-package

Indications

Induced abortion

Unspecified time of fetal death, cause not specified

Miltefosine

Section

Antileishmaniasis medicines

Oral > Solid: 10 mg; 50 mg

Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

Misoprostol

Section

Uterotonics

Oral > Solid: 200 µg

Local > Vaginal > tablet: 25 µg

Indications

Spontaneous abortion, incomplete, without complication

Unspecified obstetric condition

Postpartum haemorrhage

Mometasone

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to [budesonide](#) pour

Asthma

Mometasone + formoterol

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to [budesonide + formoterol](#) pour

Asthma

Morphine

Section	Indications
Preoperative medication and sedation for short-term procedures Parenteral > General injections > IV: 10 mg in 1 mL ampoule (sulfate OR hydrochloride) Opioid analgesics Oral > Liquid: 10 mg per 5 mL (morphine hydrochloride or morphine sulfate) Oral > Solid: 10 mg immediate release (morphine sulfate); 10 to 200 mg slow release (morphine hydrochloride or morphine sulfate) Oral > Other: 20 to 200 mg slow release granules to mix with water (morphine sulfate) Parenteral > General injections > unspecified: 10 mg in 1 mL ampoule (morphine hydrochloride or morphine sulfate)	Opioids or related analgesics and agents affecting opioid receptors Pain

Moxifloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 400 mg tablet; 100 mg tablet (dispersible) Oral > Solid: 400 mg tablet	Multi-drug resistant Mycobacterium tuberculosis Tuberculosis

Multiple micronutrient powder

Section	Indications
Vitamins and minerals Oral > Other:	Iron deficiency anaemia

Multiple micronutrient supplement

Section	Indications
Other medicines administered to the mother Tablet containing: Vit A: 800 mcg retinol activity equivalent Vit C: 70 mg Vit D: 5 mcg (200 IU) Vit E: 10 mg alpha tocopherol equivalent Vit B1: 1.4 mg Vit B2: 1.4 mg Vit B3: 18 mg niacin equivalent Vit B6: 1.9 mg Folic acid: 400 mcg Vit B12: 2.6 mcg Iron: 30 mg Iodine: 150 mcg Zinc: 15 mg Selenium: 65 mcg Copper: 2 mg	Maternal care for other specified conditions predominantly related to pregnancy

Mumps vaccine

Section	Indications
Recommendations for immunization programmes with certain characteristics All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against mumps alone

Mupirocin

Section	Indications
Dermatological medicines > Anti-infective medicines Local > Topical > Cream: 2% (as mupirocin calcium) Local > Topical > Ointment: 2%	Superficial bacterial folliculitis Impetigo

Nadroparin

Section

Medicines affecting coagulation

Indications

Therapeutic equivalent to enoxaparin pour

Acute ischaemic heart disease

Therapeutic equivalent to enoxaparin pour

Venous thromboembolism

Nalidixic acid

Section

Anti-infective medicines

Oral > Solid: 250 mg; 500 mg

Indications

Shigella resistant to unspecified antibiotic

Naloxone

Section

Antidotes and other substances used in poisonings > Specific

Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule
(naloxone hydrochloride)

Indications

Unintentional exposure to or harmful effects of
opioids or related analgesics

Naltrexone

Section

Medicines for alcohol use disorders

Parenteral > General injections > IM: 380 mg in vial (extended-release)

Oral > Solid > tablet: 50 mg

Indications

Disorders due to use of alcohol

Natamycin

Section

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Suspension: 5% eye drops

Indications

Infectious keratitis

Nelfinavir

Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 250 mg (as mesilate)

Oral > Other: 50 mg per g oral powder

Indications

Human immunodeficiency virus disease
without mention of associated disease or
condition, clinical stage unspecified

Neostigmine

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

Oral > Solid: 15 mg (neostigmine bromide)

Parenteral > General injections > unspecified: 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate); 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate)

Indications

Myasthenia gravis

Anticholinesterase reversible

Netilmicin

Section

Ophthalmological preparations > Anti-infective agents

Indications

Therapeutic equivalent to gentamicin pour

Other specified conjunctivitis

Therapeutic equivalent to gentamicin pour

Infectious blepharitis

Nevirapine

Section

Antiretrovirals > Non-nucleoside reverse transcriptase inhibitors

Oral > Liquid: 50 mg per 5 mL oral solution

Oral > Solid: 200 mg tablet; 50 mg tablet (dispersible)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Niclosamide

Section

Intestinal anthelmintics

Oral > Solid: 500 mg tablet (chewable)

Indications

Hymenolepiasis

Diphyllobothriasis

Taeniasis due to Taenia saginata

Taeniasis due to Taenia solium

Nicotinamide

Section

Vitamins and minerals

Oral > Solid: 50 mg

Indications

Pellagra

Nicotine replacement therapy

Section

Medicines for nicotine use disorders

Oral > Other: 2 mg chewing gum; 4 mg chewing gum; 2 mg lozenge; 4 mg lozenge; 1 mg per actuation oral spray

Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour

Indications

Nicotine dependence

Nifedipine

Section

Antioxytotics (tocolytics)

Oral > Solid: 10 mg (immediate-release)

Antianginal medicines

Oral > Solid: 10 mg

Antihypertensive medicines

Oral > Solid: 10 mg tablet; sustained-release formulations

Indications

Preterm labour without delivery

Angina pectoris

Essential hypertension

Nifurtimox

Section

Medicines for the treatment of 2nd stage African trypanosomiasis

Oral > Solid > tablet: 120 mg (scored); 30 mg (scored)

American trypanosomiasis

Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

Indications

African trypanosomiasis

Chagas disease

Nilotinib

Section

Targeted therapies

Oral > Solid: 150 mg capsule; 200 mg capsule

Indications

Chronic myeloid leukaemia, not elsewhere classified

Nilutamide

Section

Hormones and antihormones

Indications

Therapeutic equivalent to [bicalutamide](#) pour

Malignant neoplasms of prostate

Nitrofurantoin

Section

Access group antibiotics

Oral > Liquid: 25 mg per 5 mL oral liquid

Oral > Solid > dosage form: 100 mg; 50 mg

Indications

Premier choix

Infectious cystitis

Nitrous oxide

Section

General anaesthetics and oxygen > Inhalational medicines

Respiratory > Inhalation:

Indications

Anaesthetics and therapeutic gases

Nivolumab

Section

Immunomodulators

Parenteral > General injections > IV: 10 mg per mL concentrate solution for infusion

Indications

Other specified malignant neoplasms of bronchus or lung

Melanoma of skin

Norethisterone

Section

Oral hormonal contraceptives

Oral > Solid: 350 µg

Progestogens

Oral > Solid: 5 mg

Indications

Contact with health services for contraceptive management

Excessive menstruation with irregular cycle

Therapeutic equivalent to medroxyprogesterone acetate pour

Excessive menstruation with irregular cycle

Norethisterone enantate

Section

Injectable hormonal contraceptives

Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule oily solution

Indications

Contact with health services for reasons associated with reproduction

Normal immunoglobulin

Section

Human immunoglobulins

Parenteral > General injections > IV: 5% protein solution; 10% protein solution

Parenteral > General injections > IM: 16% protein solution

Parenteral > General injections > SC: 15% protein solution; 16% protein solution

Indications

Primary immunodeficiencies

Mucocutaneous lymph node syndrome

Langerhans cell histiocytosis

Nystatin

Section

Antifungal medicines

Oral > Liquid: 100000 IU per mL

Local > Topical > Other: 100000 IU pessary

Oral > Solid > dosage form: 500000 IU

Oral > Solid > lozenge: 100000 IU

Indications

Candidosis

Ocrelizumab

Section

Medicines for multiple sclerosis

Parenteral > General injections > IV: 30 mg per mL in 10 vial concentrate for solution

Parenteral > General injections > IV: 30 mg per mL in 10 mL vial

Indications

Multiple sclerosis

Relapsing-remitting multiple sclerosis

Octreotide

Section

Medicines for disorders of the pituitary hormone system

Parenteral > General injections > IM: 20 mg in vial (modified-release, as acetate) plus diluent

Parenteral > General injections > SC: 0.05 mg per mL in 1 mL vial (immediate-release, as acetate); 0.1 mg per mL in 1 mL vial (immediate-release, as acetate); 0.5 mg per mL in 1 mL vial (immediate-release, as acetate)

Indications

Acromegaly or pituitary gigantism

Ofloxacin

Section	Indications
Antituberculosis medicines Oral > Solid: 200 mg; 400 mg	Multi-drug resistant Mycobacterium tuberculosis
Ear, nose and throat medicines [c]	Therapeutic equivalent to ciprofloxacin pour Infectious diseases of external ear
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Solution (eye drops): 0.3%	Other specified conjunctivitis Infectious keratitis

Olanzapine

Section	Indications
Medicines for psychotic disorders Parenteral > General injections > IM: 10 mg in vial powder for injection	Therapeutic equivalent to risperidone pour Schizophrenia or other primary psychotic disorders Schizophrenia or other primary psychotic disorders

Omadacycline

Section	Indications
Reserve group antibiotics Parenteral > General injections > IV: 100 mg lyophilized powder for injection Oral > Solid: 300 mg	Methicillin resistant Staphylococcus aureus

Ombitasvir + paritaprevir + ritonavir

Section	Indications
Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations Oral > Solid: 12.5 mg + 75 mg + 50 mg tablet	Chronic hepatitis C

Omeprazole

Section	Indications
Antiulcer medicines Oral > Liquid: 20 mg powder for oral liquid; 40 mg powder for oral liquid Oral > Solid: 10 mg; 20 mg; 40 mg Parenteral > General injections > IV: 40 mg in vial	Peptic ulcer, site unspecified Gastro-oesophageal reflux disease Gastrointestinal bleeding

Ondansetron

Section

Medicines for other common symptoms in palliative care

Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride)

Oral > Liquid: 4 mg per 5 mL (ondansetron base)

Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base)

Antiemetic medicines

Parenteral > General injections > IV: 2 mg per mL in 2 mL ampoule (ondansetron base, as hydrochloride)

Oral > Liquid: 4 mg per 5 mL (ondansetron base)

Oral > Solid: 4 mg (ondansetron base); 8 mg (ondansetron base); 24 mg (ondansetron base)

Indications

Palliative care

Nausea or vomiting

Oral rehydration salts	
Section	Indications
Oral rehydration Powder for dilution in 200 mL, 500 mL, 1 L: - glucose 75 mEq - sodium 75 mEq or mmol/L - chloride 65 mEq or mmol/L - potassium 20 mEq or mmol/L - citrate 10 mmol/L - osmolarity 245 mOsm/L - glucose 13.5 g/L - sodium chloride 2.6 g/L - potassium chloride 1.5 g/L - trisodium citrate dihydrate 2.9 g/L Solutions correcting water, electrolyte and acid-base disturbances > Oral Powder for dilution in 200 mL, 500 mL, 1 L: - glucose 75 mEq - sodium 75 mEq or mmol/L - chloride 65 mEq or mmol/L - potassium 20 mEq or mmol/L - citrate 10 mmol/L - osmolarity 245 mOsm/L - glucose 13.5 g/L - sodium chloride 2.6 g/L - potassium chloride 1.5 g/L - trisodium citrate dihydrate 2.9 g/L	Diarrhoea Other specified disorders of fluid, electrolyte or acid-base balance

Oral rehydration salts - zinc sulfate	
Section	Indications
Medicines used in diarrhoea Oral > Other: ORS powder for dilution and zinc sulfate solid oral dosage form 20 mg (co-packaged)	Diarrhoea

Oseltamivir	
Section	Indications
Other antivirals Oral > Solid: 30 mg capsule (as phosphate); 45 mg capsule (as phosphate); 75 mg capsule (as phosphate)	Influenza due to identified seasonal influenza virus

Osimertinib

Section

Targeted therapies

Oral > Solid: 40 mg (as mesylate); 80 mg (as mesylate)

Indications

Other specified malignant neoplasms of bronchus or lung

Oxaliplatin

Section

Cytotoxic medicines

Parenteral > General injections > IV: 50 mg in vial powder for injection; 100 mg in vial powder for injection; 50 mg per 10 mL in 10 mL vial; 100 mg per 20 mL in 20 mL vial; 200 mg per 40 mL in 40 mL vial

Indications

Malignant neoplasm metastasis in large intestine

Malignant neoplasms of colon

Oxamniquine

Section

Antischistosomes and other antitremitode medicines

Oral > Liquid: 250 mg per 5 mL

Oral > Solid: 250 mg capsule

Indications

Schistosomiasis due to *Schistosoma mansoni*

Oxazolindinones

Section

Reserve group antibiotics

.

Indications

Other specified bacterial diseases

Oxycodone

Section

Opioid analgesics

Indications

Therapeutic equivalent to [morphine](#) pour

Pain

Oxygen

Section

General anaesthetics and oxygen > Inhalational medicines

Respiratory > Inhalation: (medical gas)

Medical gases

Respiratory > Inhalation:

Indications

Anaesthetics and therapeutic gases

Respiratory failure as acute or chronic

Oxytetracycline

Section

Ophthalmological preparations > Anti-infective agents

Indications

Therapeutic equivalent to tetracycline pour

Other specified conjunctivitis

Therapeutic equivalent to tetracycline pour

Infectious keratitis

Therapeutic equivalent to tetracycline pour

Infectious blepharitis

Therapeutic equivalent to tetracycline pour

Trachoma

Oxytocin

Section

Uterotonics

Parenteral > General injections > unspecified: 10 IU per mL

Indications

Postpartum haemorrhage

P-aminosalicylate sodium

Section

Antituberculosis medicines

Oral > Liquid: 5.52 g in sachet (equivalent to 4 g p-aminosalicylic acid)

Indications

Multi-drug resistant Mycobacterium tuberculosis

P-aminosalicylic acid

Section

Antituberculosis medicines

Oral > Solid: 4 g granules in sachet

Indications

Multi-drug resistant Mycobacterium tuberculosis

Paclitaxel

Section

Cytotoxic medicines

Parenteral > General injections > IV: 6 mg per mL

Indications

Other specified malignant neoplasms of the ovary

Other specified malignant neoplasms of bronchus or lung

Malignant neoplasms of nasopharynx

Kaposi sarcoma of unspecified primary site

Other specified malignant neoplasms of breast

Other specified carcinomas of ovary

Malignant neoplasms of breast

Malignant neoplasms of cervix uteri

Palbociclib

Section

Targeted therapies

Oral > Solid: 75 mg; 100 mg; 125 mg

Indications

Other specified malignant neoplasms of breast

Paliperidone

Section

Medicines for psychotic disorders

Parenteral > General injections > IM: 25 mg in pre-filled syringe (as palmitate); 50 mg in pre-filled syringe (as palmitate); 75 mg in pre-filled syringe (as palmitate); 100 mg in pre-filled syringe (as palmitate); 150 mg in pre-filled syringe (as palmitate)

Indications

Therapeutic equivalent to risperidone pour

Schizophrenia or other primary psychotic disorders

Schizophrenia or other primary psychotic disorders

Palonosetron

Section

Medicines for other common symptoms in palliative care

Antiemetic medicines

Indications

Therapeutic equivalent to ondansetron pour

Palliative care

Therapeutic equivalent to ondansetron pour

Nausea or vomiting

Pancreatic enzymes

Section	Indications
Pancreatic enzymes Oral > Solid: Age-appropriate formulations and doses including lipase, protease and amylase.	Enzyme intestinal

Paracetamol (acetaminophen)

Section	Indications
Non-opioids and non-steroidal anti-inflammatory medicines (NSAIDs) Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 100 mg; 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg Antimigraine medicines > For treatment of acute attack Oral > Liquid: 125 mg per 5 mL; 120 mg per 5 mL; 250 mg per 5 mL Local > Rectal > Suppository: 250 mg Oral > Solid > dispersible tablet: 100 mg; 250 mg Oral > Solid > tablet: 250 mg; 325 mg; 500 mg	Pain Migraine

Paromomycin

Section	Indications
Antileishmaniasis medicines Parenteral > General injections > IM: 750 mg paromomycin base (as sulfate)	Visceral leishmaniasis
Antiamoebic and anti-giardiasis medicines Oral > Liquid: 125 mg per 5 mL as sulfate Oral > Solid: 250 mg as sulfate	Amoebiasis

Pegaspargase

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 3750 units per 5 mL in vial; 3750 units in vial powder for injection	Lymphoid leukaemia, not elsewhere classified

Pegfilgrastim

Section	Indications
Immunomodulators Parenteral > General injections > SC: 6 mg per 0.6 mL in pre-filled syringe	Acquired neutropaenia

Pegylated interferon alfa (2a)

Section

Medicines for hepatitis C > Other antivirals for hepatitis C

Parenteral > General injections > SC: 180 µg in vial; 180 µg in prefilled syringe

Indications

Chronic hepatitis C

Pegylated interferon alfa (2b)

Section

Medicines for hepatitis C > Other antivirals for hepatitis C

Parenteral > General injections > SC: 80 µg in prefilled syringe; 100 µg in prefilled syringe

Indications

Chronic hepatitis C

Pembrolizumab

Section

Immunomodulators

Parenteral > General injections > IV: 50 mg powder for injection

Indications

Therapeutic equivalent to nivolumab pour

Melanoma of skin

Other specified malignant neoplasms of bronchus or lung

Penicillamine

Section

Antidotes and other substances used in poisonings > Specific

Oral > Solid: 250 mg

Disease-modifying anti-rheumatic drugs (DMARDs)

Oral > Solid: 250 mg

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Rheumatoid arthritis, serology unspecified

Pentamidine

Section

Antileishmaniasis medicines

Parenteral > General injections > IM: 200 mg (as isetionate); 300 mg (as isetionate)

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 200 mg tablet; 300 mg tablet

Medicines for the treatment of 1st stage African trypanosomiasis

Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection

Indications

Visceral leishmaniasis

Pneumocystosis

African trypanosomiasis

Permethrin

Section

[Dermatological medicines > Scabicides and pediculicides](#)

Local > Topical > Lotion: 1%

Local > Topical > Cream: 5%

Indications

[Pediculosis corporis](#)

[Pediculosis capitis](#)

[Scabies](#)

Pertussis vaccine

Section

[Recommendations for all immunization programmes](#)

All vaccines should comply with the WHO requirements for biological substances.

Indications

[Need for immunization against pertussis alone](#)

Pertuzumab

Section

[Targeted therapies](#)

Parenteral > General injections > IV: 240 mg per 14 mL in vial concentrated solution

Indications

[Carcinoma of breast, specialised type](#)

Phenelzine

Section

[Medicines for depressive disorders](#)

Oral > Solid > dosage form: 15 mg (as sulfate)

Indications

[Other specified depressive disorders](#)

Phenobarbital

Section

[Antiseizure medicines](#)

Oral > Liquid: 15 mg per 5 mL

Oral > Solid: 15 to 100 mg

Parenteral > General injections > IV: 200 mg per mL (phenobarbital sodium); 30 mg per mL (phenobarbital sodium); 60 mg per mL (phenobarbital sodium)

Indications

[Epilepsy or seizures](#)

[Status epilepticus](#)

Phenoxymethylpenicillin

Section	Indications
Access group antibiotics Oral > Liquid: 250 mg per 5 mL (as potassium salt) powder for oral liquid Oral > Solid: 250 mg (as potassium salt) tablet Oral > Solid: 250 mg (as potassium salt); 500 mg (as potassium salt)	Premier choix Acute pharyngitis Periapical abscess without sinus Bacterial pneumonia (Community-acquired pneumonia - mild to moderate) Autres indications Bacterial infection of unspecified site

Phenytoin

Section	Indications
Antiseizure medicines Oral > Liquid: 30 mg per 5 mL (phenytoin) Parenteral > General injections > unspecified: 50 mg per mL in vial (phenytoin sodium) Oral > Solid > tablet: 50 mg (phenytoin) (chewable) Oral > Solid > dosage form: 50 mg (phenytoin sodium); 25 mg (phenytoin sodium); 100 mg (phenytoin sodium)	Epilepsy or seizures

Phosphorus

Section	Indications
Medicines for endocrine disorders Oral > Other: 500 mg in sachet (elemental phosphorus) granules Oral > Solid > tablet: 250 mg (elemental phosphorus); 500 mg (elemental phosphorus); 500 mg (elemental phosphorus) (effervescent)	Hypophosphataemic rickets

Phytomenadione

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 10 mg per mL in ampoule; 1 mg per mL in ampoule Oral > Solid: 10 mg tablet	Haemorrhagic disorder due to other specified circulating anticoagulants Haemorrhagic or haematological disorders of fetus or newborn

Pilocarpine

Section

Ophthalmological preparations > Miotics and antiglaucoma medicines
Local > Ophthalmological > Solution (eye drops): 2% (hydrochloride or nitrate); 4% (hydrochloride or nitrate)

Indications

Acute angle closure with pupillary block

Ocular hypertension

Primary open-angle glaucoma

Piperacillin + tazobactam

Section

Watch group antibiotics

Parenteral > General injections > IV: 2 g (as sodium salt) + 250 mg (as sodium salt) powder for injection; 4 g (as sodium salt) + 500 mg (as sodium salt) powder for injection

Indications

Premier choix

Peritoneal abscess (severe)

Other specified pneumonia (Hospital-acquired pneumonia)

Neutropenia (high-risk)

Peritonitis (severe)

co-prescrire avec clindamycin

Necrotising fasciitis

Platelets

Section

Blood and blood components
Parenteral > General injections > IV:

Indications

Blood transfusion without reported diagnosis

Plazomicin

Section

Reserve group antibiotics

Parenteral > General injections > IV: 500 mg per 10 mL (Injection)

Indications

Carbapenem resistant *Acinetobacter baumannii*

Carbapenem-resistant *Pseudomonas aeruginosa*

Carbapenem resistant Enterobacterales

Pneumococcal vaccine

Section

Recommendations for all immunization programmes
All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against single bacterial diseases

Podophyllotoxin

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Indications

Therapeutic equivalent to [podophyllum resin](#) pour

Anogenital warts

Therapeutic equivalent to [podophyllum resin](#) pour

Plantar warts

Podophyllum resin

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Local > Topical > Solution: 10 to 25%

Indications

Anogenital warts

Plantar warts

Poliomyelitis vaccine

Section

Recommendations for all immunization programmes

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against poliomyelitis

Polygeline

Section

Plasma substitutes

Parenteral > General injections > IV: 3.5% injectable solution

Parenteral > General injections > IV: 3.5% in solution

Indications

Therapeutic equivalent to [dextran 70](#) pour

Hypovolaemia

Hypovolaemia

Polymyxin B (injection)

Section

Reserve group antibiotics

Parenteral > General injections > unspecified: 500000 IU in vial powder for injection (equivalent to 50 mg polymyxin B base)

Indications

Carbapenem resistant *Acinetobacter baumannii*

Carbapenem-resistant *Pseudomonas aeruginosa*

Carbapenem resistant Enterobacterales

Polymyxins

Section

Reserve group antibiotics

Indications

Other specified bacterial diseases

Potassium chloride

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Oral Oral > Liquid: powder for solution	Hypokalaemia
Solutions correcting water, electrolyte and acid-base disturbances > Parenteral Parenteral > General injections > IV: 11.2% in 20 mL ampoule (equivalent to K ⁺ 1.5 mmol/mL and Cl ⁻ 1.5 mmol/mL); 7.5% solution for dilution (equivalent to K ⁺ 1 mmol/mL and Cl ⁻ 2 mmol/mL); 15% solution for dilution (equivalent to K ⁺ 2 mmol/mL and Cl ⁻ 2 mmol/mL)	Hypokalaemia

Potassium ferric hexacyanoferrate

Section	Indications
Antidotes and other substances used in poisonings > Specific Oral > Solid: Powder for oral administration	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Potassium iodide

Section	Indications
Antifungal medicines Local > Topical > Solution: Saturated solution	Sporotrichosis
Thyroid hormones and antithyroid medicines Oral > Solid: 60 mg	Thyrotoxicosis

Potassium permanganate

Section	Indications
Dermatological medicines > Anti-infective medicines Local > Topical > Solution: aqueous solution: 1:10 000	Tropical phagedaenic ulcer
	Pemphigus
	Other specified pyogenic bacterial infection of skin and subcutaneous tissue
	Impetigo

Povidone iodine

Section	Indications
Antiseptics Local > Topical > Solution: 10% (equivalent to 1% available iodine)	Povidone iodine

Pralidoxime

Section

Antidotes and other substances used in poisonings > Specific

Indications

Exposure to or harmful effects of undetermined intent of pesticides

Pramipexole

Section

Antiseizure medicines

Oral > Solid: 125 µg immediate-release; 250 mg immediate-release; 500 µg immediate-release; 750 µg immediate-release; 1 mg immediate-release; 1.5 mg immediate-release; 375 µg extended-release; 750 µg extended-release; 1.5 mg extended-release; 2.25 mg extended-release; 3 mg extended-release; 3.75 mg extended-release; 4.5 extended-release

Indications

Parkinson disease

Pravastatin

Section

Lipid-lowering agents

Indications

Therapeutic equivalent to [simvastatin](#) pour

Mixed hyperlipidaemia

Therapeutic equivalent to [simvastatin](#) pour

Coronary atherosclerosis

Praziquantel

Section

Antischistosomes and other antitremitode medicines

Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg

Indications

Paragonimiasis

Clonorchiasis

Opisthorchiasis

Schistosomiasis

Intestinal anthelmintics

Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg

Taeniasis due to *Taenia saginata*

Taeniasis due to *Taenia solium*

Diphyllobothriasis

Hymenolepiasis

Cysticidal medicines

Oral > Solid > tablet: 600 mg (scored); 500 mg; 150 mg

Cysticercosis of central nervous system

Precipitated sulfur

Section

[Dermatological medicines](#) > [Scabicides and pediculicides](#)

Local > Topical > Ointment:

Indications

Therapeutic equivalent to [benzyl benzoate](#) pour

Pthiriasis

Therapeutic equivalent to [benzyl benzoate](#) pour

Pediculosis corporis

Therapeutic equivalent to [benzyl benzoate](#) pour

Pediculosis capitis

Therapeutic equivalent to [benzyl benzoate](#) pour

Scabies

Prednisolone

Section

Hormones and antihormones

Oral > Liquid: 5 mg per mL

Oral > Solid: 5 mg; 25 mg

Anti-inflammatory medicines

Local > Rectal > Retention enema:

Local > Rectal > Retention enema: 20 mg per 100 mL (as sodium phosphate)

Antiallergics and medicines used in anaphylaxis

Oral > Liquid: 5 mg per mL

Oral > Solid: 5 mg; 25 mg

Ophthalmological preparations > Anti-inflammatory agents

Local > Ophthalmological > Solution (eye drops): 0.5% (sodium phosphate)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Plasma cell myeloma

Follicular lymphoma

Lymphoid leukaemia, not elsewhere classified

Malignant neoplasms of prostate

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Hodgkin lymphoma

Diffuse large B-cell lymphomas

Burkitt lymphoma including Burkitt leukaemia

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Therapeutic equivalent to hydrocortisone pour

Crohn disease site

Crohn disease site

Ulcerative colitis

Allergic or hypersensitivity conditions of unspecified type

Scleritis

Anterior uveitis

Prednisone

Section

Hormones and antihormones

Indications

Therapeutic equivalent to prednisolone pour

Plasma cell myeloma

Therapeutic equivalent to prednisolone pour

Follicular lymphoma

Therapeutic equivalent to prednisolone pour

Lymphoid leukaemia, not elsewhere classified

Therapeutic equivalent to prednisolone pour

Malignant neoplasms of prostate

Therapeutic equivalent to prednisolone pour

Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Therapeutic equivalent to prednisolone pour

Hodgkin lymphoma

Therapeutic equivalent to prednisolone pour

Diffuse large B-cell lymphomas

Therapeutic equivalent to prednisolone pour

Anaplastic large cell lymphoma, ALK-negative

Therapeutic equivalent to prednisolone pour

Anaplastic large cell lymphoma, ALK-positive

Therapeutic equivalent to prednisolone pour

Langerhans cell histiocytosis

Antiallergics and medicines used in anaphylaxis

Therapeutic equivalent to prednisolone pour

Allergic or hypersensitivity conditions of unspecified type

Pretomanid

Section

Antituberculosis medicines

Oral > Solid > tablet: 200 mg

Indications

Multi-drug resistant Mycobacterium tuberculosis

Primaquine

Section

Antimalarial medicines > For curative treatment

Oral > Solid: 7.5 mg tablet (as phosphate); 15 mg tablet (as phosphate)

Indications

Malaria due to Plasmodium ovale

Malaria due to Plasmodium vivax

Probenecid

Section

Medicines used to treat gout

Oral > Solid: 500 mg

Indications

Gout

Procainamide

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule (hydrochloride)

Oral > Solid: 500 mg (hydrochloride); 250 mg (hydrochloride)

Indications

Cardiac arrhythmia

Procaine benzylpenicillin

Section

Access group antibiotics

Parenteral > General injections > IM: 1 g in vial (=1 million IU) powder for injection; 3 g in vial (=3 million IU) powder for injection

Indications

Premier choix

Congenital syphilis [children]

Second choix

Syphilis

Procarbazine

Section

Cytotoxic medicines

Oral > Solid: 50 mg (as hydrochloride)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Hodgkin lymphoma

Progesterone vaginal ring

Section

Intravaginal contraceptives

Local > Vaginal > vaginal ring: 2.074 g micronized progesterone

Indications

Contact with health services for contraceptive management

Proguanil

Section

Antimalarial medicines > For chemoprevention

Oral > Solid: 100 mg tablet (hydrochloride)

Indications

Malaria due to Plasmodium falciparum

Promethazine

Section	Indications
Antiemetic medicines Oral > Liquid: 5 mg per 5 mL (hydrochloride) Oral > Solid: 10 mg (hydrochloride); 25 mg (hydrochloride) Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule	Nausea or vomiting

Propanol

Section	Indications
Antiseptics	Therapeutic equivalent to ethanol pour Denatured alcohol

Propofol

Section	Indications
General anaesthetics and oxygen > Injectable medicines Parenteral > General injections > IV: 10 mg per mL; 20 mg per mL	Anaesthetics and therapeutic gases

Propranolol

Section	Indications
Antihypertensive medicines Oral > Solid: 80 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Essential hypertension
Antianginal medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride) Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Angina pectoris
Antiarrhythmic medicines Parenteral > General injections > IV: 1 mg in 1 mL ampoule (hydrochloride) Oral > Solid: 10 mg tablet (hydrochloride); 40 mg tablet (hydrochloride)	Cardiac arrhythmia
Antimigraine medicines > For prophylaxis Oral > Solid: 40 mg tablet (hydrochloride); 20 mg tablet (hydrochloride)	Migraine

Propylthiouracil

Section	Indications
Thyroid hormones and antithyroid medicines Oral > Solid: 50 mg	Thyrotoxicosis

Prostaglandin E1

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 0.5 mg per mL in alcohol

Indications

Other specified congenital anomaly of great arteries including arterial duct

Prostaglandin E2

Section

Medicines administered to the neonate [c]

Parenteral > General injections > IV: 1 mg per mL

Indications

Therapeutic equivalent to [prostaglandin E1](#) pour

Other specified congenital anomaly of great arteries including arterial duct

Other specified congenital anomaly of great arteries including arterial duct

Protamine sulfate

Section

Medicines affecting coagulation

Parenteral > General injections > IV: 10 mg per mL in 5 mL ampoule

Indications

Haemorrhagic disorder due to other specified circulating anticoagulants

Protionamide

Section

Antituberculosis medicines

Indications

Therapeutic equivalent to [ethionamide](#) pour

Multi-drug resistant *Mycobacterium tuberculosis*

Pyrantel

Section

Intestinal anthelmintics

Oral > Solid > tablet: 250 mg (chewable) (as embonate or pamoate)

Indications

Ascariasis

Enterobiasis

Trichuriasis

Hookworm diseases

Pyrazinamide

Section

Antituberculosis medicines

Oral > Solid: 400 mg; 150 mg tablet (dispersible); 500 mg

Indications

Tuberculosis

Pyridostigmine

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

Parenteral > General injections > IV: 1 mg in 1 mL ampoule
(pyridostigmine bromide)

Oral > Solid: 60 mg (pyridostigmine bromide)

Indications

Myasthenia gravis

Pyridoxine

Section

Vitamins and minerals

Oral > Solid: 25 mg (hydrochloride)

Indications

Vitamin B6 deficiency

Sideroblastic anaemia

Other specified polyneuropathy

Pyrimethamine

Section

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 25 mg

Indications

Toxoplasmosis

Quetiapine

Section

Medicines for psychotic disorders

Medicines for bipolar disorders

Oral > Solid > tablet: 25 mg (immediate-release); 100 mg (immediate-release); 150 mg (immediate-release); 200 mg (immediate-release); 300 mg (immediate-release); 50 mg (modified-release); 150 mg (modified-release); 200 mg (modified-release); 300 mg (modified-release); 400 mg (modified-release)

Indications

Therapeutic equivalent to risperidone pour

Schizophrenia or other primary psychotic disorders

Bipolar or related disorders

Quinidine

Section

Antiarrhythmic medicines

Oral > Solid: 200 mg (sulfate)

Indications

Cardiac arrhythmia

Quinine

Section

Antimalarial medicines > For curative treatment

Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride)

Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)

Indications

Malaria due to *Plasmodium falciparum*

Rabies vaccine

Section

Recommendations for some high-risk populations

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against rabies

Raltegravir

Section

Antiretrovirals > Integrase inhibitors

Oral > Liquid: 100 mg granules for oral suspension

Oral > Solid: 400 mg tablet; 25 mg tablet (chewable)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Ranibizumab

Section

Ophthalmological preparations > Anti-vascular endothelial growth factor (VEGF) preparations

.

Indications

Retinal venous occlusions

Diabetic macular oedema

Degenerative high myopia

Age related macular degeneration

Ranitidine

Section

Antiulcer medicines

Oral > Liquid: 75 mg per 5 mL (as hydrochloride)

Oral > Solid: 150 mg (as hydrochloride)

Parenteral > General injections > unspecified: 25 mg per mL in 2 mL ampoule (as hydrochloride)

Indications

Gastro-oesophageal reflux disease

Peptic ulcer, site unspecified

Rasburicase

Section	Indications
Supportive medicines Parenteral > General injections > IV: 1.5 mg in vial powder and solvent for solution; 7.5 mg in vial powder and solvent for solution	Tumour lysis syndrome

Ravidasvir

Section	Indications
Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations Oral > Solid > tablet: 200 mg	Chronic hepatitis C

Ready to use therapeutic food

Section	Indications
Therapeutic foods Biscuit or paste of nutritional composition as determined by the UN joint statement on the community-based management of severe acute malnutrition and Codex alimentarius guidelines.	Acute malnutrition in infants, children or adolescents

Realgar-indigo naturalis formulation

Section	Indications
Cytotoxic medicines Oral > Solid: 270 mg (containing tetra-arsenic tetra-sulfide 30 mg)	Acute myeloid leukaemia with recurrent genetic abnormalities

Red blood cells

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Resin-based composite (high-viscosity)

Section	Indications
Dental medicines and preparations Local > Topical > Other: Single-use capsule or multi-use syringe	Dental caries

Resin-based composite (low-viscosity)

Section	Indications
Dental medicines and preparations Local > Topical > Other: Single-use applicator or multi-use bottle	Dental caries

Retinol

Section

Vitamins and minerals

Parenteral > General injections > IM: 100000 IU in 2 mL ampoule (as palmitate) water-miscible injection

Oral > Liquid: 100000 IU per mL (as palmitate) oral oily solution in multidose dispenser

Oral > Solid: 10000 IU (as palmitate) sugar-coated tablet; 200000 IU (as palmitate) capsule; 50000 IU (as palmitate) capsule; 100000 IU (as palmitate) capsule

Indications

Vitamin A deficiency

Ribavirin

Section

Other antivirals

Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution

Oral > Solid: 200 mg; 400 mg; 600 mg

Medicines for hepatitis C > Other antivirals for hepatitis C

Parenteral > General injections > IV: 1000 mg per 10 mL phosphate buffer solution; 800 mg per 10 mL phosphate buffer solution

Oral > Solid: 200 mg; 400 mg; 600 mg

Indications

Viral haemorrhagic fever, not elsewhere classified

Chronic hepatitis C

Ribociclib

Section

Targeted therapies

Indications

Therapeutic equivalent to [palbociclib](#) pour

Other specified malignant neoplasms of breast

Riboflavin

Section

Vitamins and minerals

Oral > Solid: 5 mg

Indications

Vitamin B2 deficiency

Rifabutin

Section

Antituberculosis medicines

Oral > Solid: 150 mg

Indications

Tuberculosis

Rifampicin

Section

Antituberculosis medicines

Oral > Liquid: 20 mg per mL

Oral > Solid: 150 mg; 300 mg

Parenteral > General injections > IV: 600 mg in vial powder for injection

Antileprosy medicines

Oral > Liquid: 20 mg per mL

Oral > Solid: 150 mg; 300 mg

Indications

Tuberculosis

Other specified tuberculosis

Leprosy

Rifapentine

Section

Antituberculosis medicines

Oral > Solid: 150 mg tablet; 300 mg tablet

Indications

Tuberculosis

Latent tuberculosis

Risdiplam

Section

Medicines for diseases of the nervous system

Oral > Liquid: 0.75 mg per mL powder for oral liquid

Indications

Spinal muscular atrophy

Risperidone

Section

Medicines for psychotic disorders

Oral > Solid: 0.25 mg; 0.5 mg; 1 mg; 2 mg; 3 mg; 4 mg; 6 mg

Parenteral > General injections > IM:

Indications

Schizophrenia or other primary psychotic disorders

Therapeutic equivalent to paliperidone pour

Schizophrenia or other primary psychotic disorders

Ritonavir

Section

Antiretrovirals > Protease inhibitors

Oral > Solid: 25 mg tablet (heat stable); 100 mg tablet (heat stable)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Rituximab

Section	Indications
Targeted therapies Parenteral > General injections > IV: 100 mg per 10 mL in 10 mL vial; 500 mg per 50 mL in 50 mL vial	Follicular lymphoma Chronic lymphocytic leukaemia or small lymphocytic lymphoma Diffuse large B-cell lymphomas Burkitt lymphoma including Burkitt leukaemia
Medicines for multiple sclerosis Parenteral > General injections > IV: 500 mg per 50 mL in 50 mL vial	Multiple sclerosis

Rivaroxaban

Section	Indications
Medicines affecting coagulation	Therapeutic equivalent to dabigatran pour Venous thromboembolism Therapeutic equivalent to dabigatran pour Other specified atrial fibrillation

Ropinirole

Section	Indications
Antiseizure medicines Oral > Solid: 0.25 mg immediate-release; 0.5 mg immediate-release; 1 mg immediate-release; 2 immediate-release; 3 mg immediate-release; 4 mg immediate-release; 5 mg immediate-release; 2 mg extended-release; 4 mg extended-release; 6 mg extended-release; 8 mg extended-release; 12 mg extended-release	Parkinson disease

Rotavirus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Rubella vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against rubella alone

Salbutamol

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease
Parenteral > General injections > IV: 50 µg per mL in 5 mL ampoule (as sulfate)
Respiratory > Inhalation > aerosol: 100 µg per dose (as sulfate); 100 µg per dose (as sulfate) metered-dose inhaler
Respiratory > Inhalation > solution: 5 mg per mL (as sulfate) for use in nebulizers

Indications

Chronic obstructive pulmonary disease

Asthma

Salicylic acid

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation
Local > Topical > Solution: 5%

Indications

Seborrhoeic dermatitis

Psoriasis of unspecified type

Ichthyoses

Saquinavir

Section

Antiretrovirals > Protease inhibitors
Oral > Solid: 200 mg; 500 mg

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Selenium sulfide

Section

Dermatological medicines > Antifungal medicines
Local > Topical > Suspension: 2% detergent-based

Indications

Seborrhoeic dermatitis

Pityriasis versicolor

Senna

Section

Medicines for other common symptoms in palliative care
Oral > Liquid: 7.5 mg per 5 mL oral liquid

Laxatives
Oral > Solid: 7.5 mg (sennosides) (or traditional dosage forms)

Indications

Palliative care

Constipation

Sevoflurane

Section

General anaesthetics and oxygen > Inhalational medicines
Respiratory > Inhalation > liquid:

Indications

Anaesthetics and therapeutic gases

Silver diamine fluoride

Section

Dental medicines and preparations
Local > Dental > Solution: 38% w/v

Indications

Dental caries

Silver nitrate

Section

Ophthalmological preparations > Anti-infective agents
Local > Ophthalmological > Solution (eye drops): 1%

Indications

Neonatal conjunctivitis or dacryocystitis

Silver sulfadiazine

Section

Dermatological medicines > Anti-infective medicines
Local > Topical > Cream: 1%

Indications

Burns

Simeprevir

Section

Medicines for hepatitis C > Non-pangenotypic direct-acting antiviral combinations
Oral > Solid: 150 mg

Indications

Chronic hepatitis C

Simvastatin

Section

Lipid-lowering agents
Oral > Solid: 5 mg; 10 mg; 20 mg; 40 mg

Indications

Mixed hyperlipidaemia

Coronary atherosclerosis

Medicines for endocrine disorders
Oral > Solid: 20 mg

Polycystic ovary syndrome

Smallpox vaccine

Section

Recommendations for all immunization programmes
All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against certain single viral diseases

Snake antivenom

Section

Immunologicals > Sera, immunoglobulins and monoclonal antibodies
.

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium calcium edetate

Section

Antidotes and other substances used in poisonings > Specific
Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium chloride

Section

Solutions correcting water, electrolyte and acid-base disturbances >
Parenteral
Parenteral > General injections > IV: 0.9% isotonic (equivalent to Na⁺ 154 mmol/L and Cl⁻ 154 mmol/L)

Indications

Other specified disorders of fluid, electrolyte or acid-base balance

Sodium hyaluronate

Section

Ophthalmological preparations

Indications

Therapeutic equivalent to hypromellose pour

Keratoconjunctivitis sicca

Sodium hydrogen carbonate

Section

Solutions correcting water, electrolyte and acid-base disturbances >
Parenteral
Parenteral > General injections > IV: 1.4% isotonic (equivalent to Na⁺ 167 mmol/L and HCO₃⁻ 167 mmol/L); 8.4% in 10 mL ampoule (equivalent to Na⁺ 1000 mmol/L and HCO₃⁻ 1000 mmol/L)

Indications

Acidosis

Sodium nitrite

Section

Antidotes and other substances used in poisonings > Specific
Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium nitroprusside

Section

Antihypertensive medicines
Parenteral > General injections > IV: 50 mg in ampoule powder for infusion

Indications

Hypertensive crisis

Sodium stibogluconate

Section

Antileishmaniasis medicines
Parenteral > General injections > unspecified: 100 mg per mL in 30 mL vial

Indications

Visceral leishmaniasis

Mucocutaneous leishmaniasis

Cutaneous leishmaniasis

Sodium thiosulfate

Section

Dermatological medicines > Antifungal medicines
Local > Topical > Solution: 15%

Antidotes and other substances used in poisonings > Specific
Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule

Indications

Pityriasis versicolor

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sofosbuvir

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations
Oral > Solid: 400 mg; 200 mg

Indications

Chronic hepatitis C

Sofosbuvir + velpatasvir

Section

Medicines for hepatitis C > Pangenotypic direct-acting antiviral combinations
Oral > Solid: 400 mg + 100 mg tablet; 200 mg + 50 mg tablet

Indications

Chronic hepatitis C

Somatropin

Section	Indications
Medicines for endocrine disorders Parenteral > General injections > SC: 5 mg in cartridge powder for injection; 5 mg per mL in 2 mL cartridge	Hypoglycaemia without associated diabetes

Spectinomycin

Section	Indications
Access group antibiotics Parenteral > General injections > unspecified: 2 g in vial (as hydrochloride) powder for injection	Second choix Gonococcal infection Autres indications Bacterial infection of unspecified site

Spironolactone

Section	Indications
Medicines used in heart failure Oral > Solid: 25 mg Diuretics Oral > Liquid: 10 mg per 5 mL; 25 mg per 5 mL; 5 mg per 5 mL Oral > Solid: 25 mg	Heart failure Oedema Ascites

Stavudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Liquid: 5 mg per 5 mL powder for oral solution Oral > Solid: 15 mg; 20 mg; 30 mg	Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Streptokinase

Section	Indications
Thrombolytic medicines Parenteral > General injections > IV: 1.5 million IU in vial powder for injection	Acute myocardial infarction

Streptomycin (injection)

Section

Antituberculosis medicines

Parenteral > General injections > IV: 1 g powder for injection (as sulfate) as vial

Indications

Multi-drug resistant Mycobacterium tuberculosis

Succimer

Section

Antidotes and other substances used in poisonings > Specific

Oral > Solid: 100 mg

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sulfacetamide

Section

Ophthalmological preparations > Anti-infective agents

Local > Ophthalmological > Ointment: 10% (sodium salt)

Indications

Other specified conjunctivitis

Sulfadiazine

Section

Antipneumocystosis and antitoxoplasmosis medicines

Oral > Solid: 500 mg

Anti-infective medicines

Oral > Solid: 500 mg

Parenteral > General injections > unspecified: 250 mg in 4 mL ampoule (sodium salt)

Indications

Toxoplasmosis

Bacterial infection of unspecified site

Sulfadimidine

Section

Anti-infective medicines

Oral > Liquid: 500 mg per 5 mL

Oral > Solid: 500 mg

Parenteral > General injections > unspecified: 1 g in 3 mL ampoule (sodium salt)

Indications

Bacterial infection of unspecified site

Sulfadoxine + pyrimethamine

Section	Indications
Antimalarial medicines > For curative treatment Oral > Solid: 500 mg + 25 mg tablet	Malaria due to Plasmodium falciparum Malaria due to Plasmodium vivax
Antimalarial medicines > For chemoprevention Oral > Solid: 500 mg + 25 mg tablet Oral > Solid: 250 mg + 12.5 mg tablet	Malaria due to Plasmodium falciparum Malaria

Sulfamethoxazole + trimethoprim

Section	Indications
Access group antibiotics Parenteral > General injections > IV: 80 mg + 16 mg per mL in 10 mL ampoule; 80 mg + 16 mg per mL in 5 mL ampoule Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet Oral > Solid > dispersible tablet: 100 mg + 20 mg	Premier choix Infectious cystitis Second choix Infectious gastroenteritis or colitis without specification of infectious agent
Antipneumocystosis and antitoxoplasmosis medicines Parenteral > General injections > IV: 80 mg + 16 mg per mL in 5 mL ampoule; 80 mg + 16 mg per mL in 10 mL ampoule Oral > Liquid: 200 mg + 40 mg per 5 mL oral liquid Oral > Solid: 100 mg + 20 mg tablet; 400 mg + 80 mg tablet; 800 mg + 160 mg tablet Oral > Solid > dispersible tablet: 100 mg + 20 mg	Autres indications Bacterial infection of unspecified site Pneumocystosis

Sulfasalazine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) Oral > Solid: 500 mg	Rheumatoid arthritis, serology unspecified
Anti-inflammatory medicines Oral > Solid: 500 mg Local > Rectal > Suppository: 500 mg Local > Rectal > Retention enema:	Crohn disease site Ulcerative colitis

Sumatriptan

Section

Antimigraine medicines > For treatment of acute attack
Oral > Solid: 50 mg

Indications

Migraine

Sunscreen (broad-spectrum)

Section

Dermatological medicines
Topical cream, lotion, gel, ointment, spray with sun protection factor $\geq$ 50.

Indications

Albinism or other specified genetically-determined hypomelanotic disorders

Xeroderma pigmentosum

Suramin sodium

Section

Antifilarials
Parenteral > General injections > IV: 1 g in vial

Medicines for the treatment of 1st stage African trypanosomiasis
Parenteral > General injections > IV: 1 g in vial

Indications

Filariasis

African trypanosomiasis

Surfactant

Section

Medicines administered to the neonate [c]
Respiratory > Suspension: 80 mg per mL for intratracheal instillation; 25 mg per mL for intratracheal instillation

Indications

Respiratory distress syndrome of newborn

Suxamethonium

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors
Parenteral > General injections > IV: 50 mg per mL in 2 mL ampoule (suxamethonium chloride); powder for injection (suxamethonium chloride) in vial

Indications

Muscle relaxants

Tacalcitol

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Indications

Therapeutic equivalent to calcipotriol pour

Psoriasis of unspecified type

Tacrolimus

Section	Indications
Immunomodulators for non-malignant disease Parenteral > General injections > IV: 5 mg per mL in 1 mL vial Oral > Liquid: 0.2 mg granules for oral suspension; 1 mg granules for oral suspension Oral > Solid: 0.5 mg (immediate-release); 0.75 mg (immediate-release); 1 mg (immediate-release); 2 mg (immediate-release); 5 mg (immediate-release)	Failure or rejection of transplanted organs or tissues

Tamoxifen

Section	Indications
Hormones and antihormones Oral > Solid: 10 mg (as citrate); 20 mg (as citrate)	Other specified malignant neoplasms of breast Malignant neoplasms of breast

Tedizolid

Section	Indications
Reserve group antibiotics	Therapeutic equivalent to linezolid pour Methicillin resistant Staphylococcus aureus Therapeutic equivalent to linezolid pour Vancomycin resistant Staphylococcus aureus Therapeutic equivalent to linezolid pour Vancomycin resistant Enterococcus

Telmisartan + amlodipine

Section	Indications
Antihypertensive medicines Oral > Solid: 40 mg + 5 mg; 80 mg + 5 mg; 80 mg + 10 mg	Essential hypertension

Telmisartan + hydrochlorothiazide

Section	Indications
Antihypertensive medicines Oral > Solid: 40 mg + 12.5 mg; 80 mg + 12.5 mg; 80 mg + 25 mg	Essential hypertension

Tenofovir alafenamide

Section

Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors
Oral > Solid: 25 mg

Indications

Chronic hepatitis B

Tenofovir disoproxil fumarate

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors
Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Contact with or exposure to human immunodeficiency virus

Medicines for hepatitis B > Nucleoside/Nucleotide reverse transcriptase inhibitors

Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)

Chronic hepatitis B

Terbinafine

Section

Dermatological medicines > Antifungal medicines
Local > Topical > Cream: 1% terbinafine hydrochloride
Local > Topical > Ointment: 1% terbinafine hydrochloride

Indications

Fungal infection of the skin

Terbutaline

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Indications

Therapeutic equivalent to salbutamol pour

Chronic obstructive pulmonary disease

Therapeutic equivalent to salbutamol pour

Asthma

Terizidone

Section

Antituberculosis medicines

Indications

Therapeutic equivalent to cycloserine pour

Multi-drug resistant Mycobacterium tuberculosis

Testosterone

Section	Indications
Androgens Parenteral > General injections > IM: 200 mg per mL in 1 mL ampoule (enantate)	Testicular dysfunction or testosterone-related disorders

Tetanus antitoxin

Section	Indications
Immunologicals > Sera, immunoglobulins and monoclonal antibodies Parenteral > General injections > IM: 50000 IU in vial	Tetanus

Tetanus vaccine

Section	Indications
Recommendations for all immunization programmes All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against tetanus alone

Tetracaine

Section	Indications
Ophthalmological preparations > Local anaesthetics Local > Ophthalmological > Solution (eye drops): 0.5% (hydrochloride)	Local anaesthetics

Tetracycline

Section	Indications
Access group antibiotics Oral > Solid: 250 mg (hydrochloride)	Bacterial infection of unspecified site
Ophthalmological preparations > Anti-infective agents Local > Ophthalmological > Ointment: 1% (tetracycline hydrochloride)	Neonatal conjunctivitis or dacryocystitis
	Other specified conjunctivitis
	Infectious keratitis
	Infectious blepharitis
	Trachoma

Thalidomide

Section	Indications
Immunomodulators Oral > Solid: 50 mg	Plasma cell myeloma

Thiamine

Section

Vitamins and minerals

Oral > Solid: 50 mg (hydrochloride)

Indications

Vitamin B1 deficiency

Thiopental

Section

General anaesthetics and oxygen > Injectable medicines

Indications

Therapeutic equivalent to propofol pour

Anaesthetics and therapeutic gases

Ticagrelor

Section

Anti-platelet medicines

Oral > Solid > tablet: 60 mg; 90 mg

Indications

Acute ischaemic heart disease

Presence of coronary angioplasty implant or graft

Tick-borne encephalitis vaccine

Section

Recommendations for certain regions

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against arthropod-borne viral encephalitis

Tigecycline

Section

Reserve group antibiotics

Parenteral > General injections > IV: 50 mg in vial powder for injection

Indications

Other specified bacterial diseases

Timolol

Section

Ophthalmological preparations > Miotics and antiglaucoma medicines

Local > Ophthalmological > Solution (eye drops): 0.25% (hydrogen maleate); 0.5% (hydrogen maleate)

Indications

Primary open-angle glaucoma

Ocular hypertension

Tinidazole

Section

Antiamoebic and anti giardiasis medicines

Indications

Therapeutic equivalent to metronidazole pour

Amoebiasis

Tioguanine

Section

Cytotoxic medicines

Oral > Solid: 40 mg

Indications

Lymphoid leukaemia, not elsewhere classified

Tiotropium bromide

Section

Antiasthmatic and medicines for chronic obstructive pulmonary disease

Respiratory > Inhalation > dry powder: 18 µg capsule

Respiratory > Inhalation > solution: 1.25 µg per actuation; 2.5 µg per actuation

Indications

Chronic obstructive pulmonary disease

Tisagenlecleucel

Section

Antineoplastics and supportive medicines

Indications

Therapeutic equivalent to axicabtagene ciloleucel pour

Diffuse large B-cell lymphomas

Tislelizumab

Section

Immunomodulators

Parenteral > General injections > IV: 100 mg per 10 mL

Indications

Malignant neoplasms of urinary tract

Hodgkin lymphoma

Other specified malignant neoplasms of bronchus or lung

Tobramycin

Section

Ophthalmological preparations > Anti-infective agents

Indications

Therapeutic equivalent to gentamicin pour

Other specified conjunctivitis

Therapeutic equivalent to gentamicin pour

Infectious blepharitis

Tocilizumab

Section

Medicines for juvenile joint diseases

Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial

Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe

Indications

Juvenile systemic arthritis

Tolbutamide

Section

Oral hypoglycaemic agents

Oral > Solid: 500 mg

Indications

Type 2 diabetes mellitus

Torsemide

Section

Medicines used in heart failure

Diuretics

Indications

Therapeutic equivalent to furosemide pour

Heart failure

Therapeutic equivalent to furosemide pour

Anuria or oliguria

Therapeutic equivalent to furosemide pour

Oedema

Toripalimab

Section

Immunomodulators

Parenteral > General injections > IV: 240 mg per 6 mL in vial

Indications

Malignant neoplasms of nasopharynx

Malignant neoplasms of oesophagus

Tramadol

Section	Indications
Opioid analgesics Oral > Liquid: 100 mg per mL (hydrochloride) Oral > Solid: 50 mg (hydrochloride) immediate release; 50 mg (hydrochloride) controlled release; 100 mg (hydrochloride) controlled release; 150 mg (hydrochloride) controlled release; 200 mg (hydrochloride) controlled release; 300 mg (hydrochloride) controlled release; 400 mg (hydrochloride) controlled release Parenteral > General injections > unspecified: 50 mg per mL in 2 mL ampoule (hydrochloride)	Chronic cancer pain

Trametinib

Section	Indications
Targeted therapies Oral > Solid: 0.5 mg; 2 mg	Other specified melanoma of skin

Tranexamic acid

Section	Indications
Medicines affecting coagulation Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Haemorrhage, not elsewhere classified
Other medicines administered to the mother Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule	Postpartum haemorrhage

Trastuzumab

Section	Indications
Targeted therapies Parenteral > General injections > IV: 60 mg in vial powder for injection; 150 mg in vial powder for injection; 440 mg in vial powder for injection	Carcinoma of breast, specialised type

Trastuzumab emtansine

Section	Indications
Targeted therapies Parenteral > General injections > IV: 100 mg in vial powder for injection; 160 mg in vial powder for injection	Carcinoma of breast, specialised type

Triamcinolone hexacetonide

Section

Medicines for juvenile joint diseases

Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

Indications

Juvenile idiopathic arthritis

Triclabendazole

Section

Antischistosomes and other antitremitode medicines

Oral > Solid > tablet: 250 mg (scored)

Indications

Fascioliasis

Paragonimiasis

Trihexyphenidyl

Section

Antiseizure medicines

Oral > Solid: 2 mg (hydrochloride); 5 mg (hydrochloride)

Medicines for parkinsonism

Indications

Parkinson disease

Therapeutic equivalent to biperiden pour

Parkinson disease

Trimethoprim

Section

Access group antibiotics

Oral > Liquid: 50 mg per mL

Oral > Solid: 100 mg; 200 mg

Indications

Premier choix

Infectious cystitis

Triptorelin

Section

Hormones and antihormones

Indications

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of breast

Therapeutic equivalent to leuprorelin pour

Malignant neoplasms of prostate

Tropicamide

Section

Diagnostic agents > Ophthalmic medicines

Local > Ophthalmological > Solution (eye drops): 0.5%

Indications

Cycloplegic drug

Tropisetron

Section	Indications
Medicines for other common symptoms in palliative care	Therapeutic equivalent to ondansetron pour Palliative care
Antiemetic medicines	Therapeutic equivalent to ondansetron pour Nausea or vomiting

Tuberculin, purified protein derivative

Section	Indications
Immunologicals > Diagnostic agents Parenteral > Locoregional injections > Intradermal:	Mycobacterial diseases

Tubocurarine

Section	Indications
Muscle relaxants (peripherally-acting) and cholinesterase inhibitors Parenteral > General injections > IV: 10 mg per mL in 1.5 mL ampoule (tubocurarine chloride)	Muscle relaxants

Typhoid vaccine

Section	Indications
Recommendations for some high-risk populations All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against typhoid-paratyphoid alone

Ulipristal

Section	Indications
Oral hormonal contraceptives Oral > Solid: 30 mg tablet (ulipristal acetate)	Contact with health services for postcoital contraception

Umeclidinium

Section	Indications
Antiasthmatic and medicines for chronic obstructive pulmonary disease	Therapeutic equivalent to tiotropium bromide pour Chronic obstructive pulmonary disease

Urea

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Local > Topical > Cream: 10%; 5%

Local > Topical > Ointment: 10%; 5%

Indications

Xerosis cutis or asteatosis

Ustekinumab

Section

Dermatological medicines > Medicines affecting skin differentiation and proliferation

Parenteral > General injections > SC: 45 mg per 0.5 mL in vial; 90 mg per mL in pre-filled syringe

Indications

Psoriasis of unspecified type

Valaciclovir

Section

Antiherpes medicines

Indications

Therapeutic equivalent to [aciclovir](#) pour

Zoster

Therapeutic equivalent to [aciclovir](#) pour

Varicella

Therapeutic equivalent to [aciclovir](#) pour

Herpes simplex infections

Valganciclovir

Section

Other antivirals

Oral > Liquid: 50 mg per mL powder for oral solution

Oral > Solid: 450 mg Tablet

Indications

Cytomegaloviral retinitis

Valproic acid (sodium valproate)

Section

Antiseizure medicines

Oral > Liquid: 200 mg per 5 mL

Oral > Solid: 200 mg tablet (enteric-coated); 500 mg tablet (enteric-coated); 100 mg tablet (crushable)

Parenteral > General injections > IV: 100 mg per mL in 4 mL ampoule; 100 mg per mL in 10 mL ampoule; 100 mg per mL in 3 mL ampoule

Medicines for bipolar disorders

Oral > Solid: 200 mg tablet (enteric-coated); 500 mg tablet (enteric-coated)

Indications

Epilepsy or seizures

Status epilepticus

Bipolar or related disorders

Vancomycin

Section

Watch group antibiotics

Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection

Parenteral > General injections > IV: 250 mg in vial (as hydrochloride) powder for injection; 500 mg in vial (as hydrochloride) powder for injection; 1 g in vial (as hydrochloride) powder for injection

Indications

Premier choix

Necrotising fasciitis

co-prescrire avec ceftazidime

Endophthalmitis

co-prescrire avec ceftriaxone

Endophthalmitis

Second choix

Neutropenia (high-risk)

Autres indications

Methicillin resistant Staphylococcus aureus

Varenicline

Section

Medicines for nicotine use disorders

Oral > Solid: 0.5 mg; 1 mg

Indications

Nicotine dependence

Varicella vaccine

Section

Recommendations for immunization programmes with certain characteristics

All vaccines should comply with the WHO requirements for biological substances.

Indications

Need for immunization against certain single viral diseases

Vecuronium

Section

Muscle relaxants (peripherally-acting) and cholinesterase inhibitors
Parenteral > General injections > IV: 10 mg in vial powder for injection (vecuronium bromide)

Indications

Muscle relaxants

Vemurafenib

Section

Targeted therapies

Indications

Therapeutic equivalent to [dabrafenib](#) pour

Other specified melanoma of skin

Verapamil

Section

Antiarrhythmic medicines

Parenteral > General injections > IV: 2.5 mg per mL in 2 mL ampoule (hydrochloride)

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Antianginal medicines

Oral > Solid: 40 mg (hydrochloride); 80 mg (hydrochloride)

Indications

Supraventricular tachyarrhythmia

Angina pectoris

Vinblastine

Section

Cytotoxic medicines

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate)

Parenteral > General injections > IV: 10 mg in vial (vinblastine sulfate); 10 mg per 10 mL in vial (vinblastine sulfate)

Indications

Unspecified malignant neoplasms of ill-defined or unspecified sites

Other specified gliomas of brain

Kaposi sarcoma of unspecified primary site

Hodgkin lymphoma

Other specified malignant neoplasms of the ovary

Germ cell tumour of testis

Anaplastic large cell lymphoma, ALK-negative

Anaplastic large cell lymphoma, ALK-positive

Langerhans cell histiocytosis

Vincristine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate) Parenteral > General injections > IV: 1 mg in vial (vincristine sulfate); 5 mg in vial (vincristine sulfate); 1 mg per mL in vial (vincristine sulfate); 2 mg per 2 mL in vial (vincristine sulfate)	Unspecified malignant neoplasms of ill-defined or unspecified sites Other specified gliomas of brain Malignant trophoblastic neoplasms of placenta Follicular lymphoma Kaposi sarcoma of unspecified primary site Rhabdomyosarcoma primary site Hodgkin lymphoma Diffuse large B-cell lymphomas Retinoblastoma Burkitt lymphoma including Burkitt leukaemia Malignant neoplasms of kidney, except renal pelvis Ewing sarcoma of bone and articular cartilage of unspecified sites Lymphoid leukaemia, not elsewhere classified Langerhans cell histiocytosis

Vinorelbine

Section	Indications
Cytotoxic medicines Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 50 mg per 5 mL in 5 mL vial Parenteral > General injections > IV: 10 mg per mL in 1 mL vial; 10 mg per mL in 5 mL vial Oral > Solid: 20 mg; 30 mg; 80 mg	Other specified malignant neoplasms of bronchus or lung Other specified malignant neoplasms of breast Rhabdomyosarcoma primary site

Voriconazole

Section	Indications
Antifungal medicines Parenteral > General injections > IV: 200 mg in vial powder for injection Oral > Liquid: 40 mg per mL powder for oral liquid Oral > Solid: 50 mg tablet; 200 mg tablet	Chronic pulmonary aspergillosis Invasive aspergillosis

Warfarin

Section	Indications
Medicines affecting coagulation Oral > Solid: 5 mg (sodium salt); 1 mg (sodium salt); 2 mg (sodium salt); 0.5 mg (sodium salt)	Other specified diseases of arteries or arterioles Cerebral ischaemic stroke due to embolic occlusion Venous thromboembolism Atrial fibrillation

Water for injection

Section	Indications
Solutions correcting water, electrolyte and acid-base disturbances > Miscellaneous Parenteral > General injections > unspecified: 2 mL ampoule; 5 mL ampoule; 10 mL ampoule	Unspecified appropriateness of dosing or administration without injury or harm

Whole blood

Section	Indications
Blood and blood components Parenteral > General injections > IV:	Blood transfusion without reported diagnosis

Xylometazoline

Section	Indications
Ear, nose and throat medicines [c] Local > Nasal > Spray: 0.05% w/v equivalent to 0.5 mg per mL	Nasal congestion

Yellow fever vaccine

Section	Indications
Recommendations for certain regions All vaccines should comply with the WHO requirements for biological substances.	Need for immunization against certain single viral diseases

Zanubrutinib

Section	Indications
Targeted therapies Oral > Solid: 80 mg	Mantle cell lymphoma Chronic lymphocytic leukaemia or small lymphocytic lymphoma

Zidovudine

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors
Parenteral > General injections > IV: 10 mg per mL in 20 mL vial solution for IV infusion injection
Oral > Liquid: 50 mg per 5 mL
Oral > Solid: 250 mg capsule; 300 mg tablet

Indications

Human immunodeficiency virus disease without mention of associated disease or condition, clinical stage unspecified

Zinc sulfate

Section

Medicines for diarrhoea
Oral > Solid: 20 mg

Indications

Diarrhoea

Zoledronic acid

Section

Supportive medicines
Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle
Medicines for endocrine disorders
Parenteral > General injections > IV: 4 mg per 5 mL in 5 mL vial; 4 mg per 100 mL in 100 mL bottle

Indications

Malignant neoplasm metastasis in bone or bone marrow

Osteogenesis imperfecta

Zuclopenthixol decanoate

Section

Medicines for psychotic disorders

Indications

Therapeutic equivalent to fluphenazine pour

Schizophrenia or other primary psychotic disorders