

Abacavir + dolutegravir + lamivudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid > tablet: 60 mg (as sulfate) + 5 mg + 30 mg (dispersible, scored)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Abacavir + lamivudine

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 120 mg (as sulfate) + 60 mg tablet (dispersible, scored)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Abacavir + lamivudine + lopinavir + ritonavir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 30 mg + 15 mg + 40 mg + 10 mg capsule containing oral granules	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Dolutegravir + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 50 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Efavirenz + lamivudine + tenofovir

Section	Indications
Fixed-dose combinations of antiretrovirals Oral > Solid: 400 mg + 300 mg + 300 mg (tenofovir disoproxil fumarate, equivalent to 245 mg tenofovir disoproxil)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified Therapeutic equivalent to efavirenz + emtricitabine + tenofovir for Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid: 150 mg tablet

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine + nevirapine + zidovudine

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 50 mg + 60 mg Tablet; 150 mg + 200 mg + 300 mg Tablet

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine + tenofovir

Section

Fixed-dose combinations of antiretrovirals

Indications

Therapeutic equivalent to emtricitabine + tenofovir for

Contact with or exposure to human immunodeficiency virus

Therapeutic equivalent to emtricitabine + tenofovir for

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine + zidovudine

Section

Fixed-dose combinations of antiretrovirals

- Oral > Solid: 30 mg + 60 mg Tablet; 150 mg + 300 mg Tablet

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified