

Acetylcysteine**Section**

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule
- Oral > Liquid: 10%; 20%

Indications

Acute or subacute hepatic failure

Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Atropine**Section**

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule or vial (sulfate)

Indications

Exposure to or harmful effects of undetermined intent of pesticides

Calcium gluconate**Section**

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule or vial (10%)

Indications

Hypermagnesaemia

Deferoxamine**Section**

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)

Indications

Harmful effects of drugs, medicaments or biological substances, not elsewhere classified

Dimercaprol**Section**

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule; 100 mg per mL in 3 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Fomepizole**Section**

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule or vial (sulfate); 1 g per mL in 1.5 mL ampoule or vial (base)

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Methionine

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: 250 mg (DL)

Indications

Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Methylthioninium chloride

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule

Indications

Acquired methaemoglobinaemia

Naloxone

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)

Indications

Unintentional exposure to or harmful effects of opioids or related analgesics

Penicillamine

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: 250 mg

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Potassium ferric hexacyanoferrate (Prussian blue)

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: Powder for oral administration

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Pralidoxime

Section

Antidotes and other substances used in poisonings > Specific

- .

Indications

Exposure to or harmful effects of undetermined intent of pesticides

Sodium calcium edetate

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium nitrite

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium thiosulfate

Section

Antidotes and other substances used in poisonings > Specific

- Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Succimer

Section

Antidotes and other substances used in poisonings > Specific

- Oral > Solid: 100 mg

Indications

Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified