Model List of Essential Medicines

Acetylcysteine

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 200 mg per mL in 10 mL ampoule ◦ Oral > Liquid: 10%; 20%	Acute or subacute hepatic failure Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Atropine

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > unspecified: 1 mg in 1 mL ampoule or vial (sulfate)	Exposure to or harmful effects of undetermined intent of pesticides

Calcium gluconate

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 100 mg per mL in 10 mL ampoule or vial (10%)	Hypermagnesaemia

Deferoxamine

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > unspecified: 500 mg in vial powder for injection (mesilate)	Harmful effects of drugs, medicaments or biological substances, not elsewhere classified

Dimercaprol

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IM: 50 mg per mL in 2 mL ampoule; 100 mg per mL in 3 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Fomepizole

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 5 mg per mL in 20 mL ampoule or vial (sulfate); 1 g per mL in 1.5 mL ampoule or vial (base)	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Methionine

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Oral > Solid: 250 mg (DL)	Exposure to or harmful effects of undetermined intent of analgesics, antipyretics or nonsteroidal anti-inflammatory drugs

Methylthioninium chloride

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 10 mg per mL in 10 mL ampoule	Acquired methaemoglobinaemia

Naloxone

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > unspecified: 400 µg per 1 mL in ampoule (naloxone hydrochloride)	Unintentional exposure to or harmful effects of opioids or related analgesics

Penicillamine

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Oral > Solid: 250 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Potassium ferric hexacyanoferrate (Prussian blue)

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Oral > Solid: Powder for oral administration	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Pralidoxime

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ .	Exposure to or harmful effects of undetermined intent of pesticides

Sodium calcium edetate

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 200 mg per mL in 5 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium nitrite

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 30 mg per mL in 10 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Sodium thiosulfate

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Parenteral > General injections > IV: 250 mg per mL in 50 mL ampoule	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified

Succimer

Section	Indications
Antidotes and other substances used in poisonings > Specific ◦ Oral > Solid: 100 mg	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified