

Amphotericin B

Section	Indications
Antifungal medicines ◦ Parenteral > General injections > IV: 50 mg in vial powder for injection (liposomal complex); 50 mg in vial powder for injection (as sodium deoxycholate)	Candidosis Blastomycosis Cryptococcosis Coccidioidomycosis Histoplasmosis Sporotrichosis Aspergillosis

Anidulafungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin for Systemic or invasive candidosis

Caspofungin

Section	Indications
Antifungal medicines	Therapeutic equivalent to micafungin for Systemic or invasive candidosis

Clotrimazole

Section	Indications
Antifungal medicines ◦ Local > Topical > Other: 1% vaginal cream; 10% vaginal cream; 100 mg vaginal tablet; 500 mg vaginal tablet	Vulvovaginal candidosis

Fluconazole

Section	Indications
Antifungal medicines ◦ Parenteral > General injections > IV: 2 mg per mL in vial ◦ Oral > Liquid: 50 mg per 5 mL; 50 mg per 5 mL powder for oral liquid ◦ Oral > Solid > capsule: 50 mg	Cryptococcosis Candidosis

Flucytosine

Section

Antifungal medicines

- Parenteral > General injections > IV: 2.5 g per 250 mL infusion
- Oral > Solid: 250 mg capsule

Indications

Cryptococcosis

Griseofulvin

Section

Antifungal medicines

- Oral > Liquid: 125 mg per 5 mL
- Oral > Solid: 125 mg; 250 mg

Indications

Dermatophytosis

Itraconazole

Section

Antifungal medicines

- Oral > Liquid: 10 mg per mL
- Oral > Solid: 100 mg

Indications

Chronic pulmonary aspergillosis

Chromoblastomycosis

Histoplasmosis

Talaromycosis

Sporotrichosis

Paracoccidioidomycosis

Micafungin

Section

Antifungal medicines

- Parenteral > General injections > IV: 50 mg in vial (as sodium) powder for injection; 100 mg in vial (as sodium) powder for injection

Indications

Systemic or invasive candidosis

Nystatin

Section

Antifungal medicines

- Oral > Liquid: 100000 IU per mL
- Local > Topical > Other: 100000 IU pessary
- Oral > Solid > dosage form: 500000 IU
- Oral > Solid > lozenge: 100000 IU

Indications

Candidosis

Potassium iodide

Section

Antifungal medicines

- Local > Topical > Solution: Saturated solution

Indications

Sporotrichosis

Voriconazole

Section

Antifungal medicines

- Parenteral > General injections > IV: 200 mg in vial powder for injection
- Oral > Liquid: 40 mg per mL powder for oral liquid
- Oral > Solid: 50 mg tablet; 200 mg tablet

Indications

Invasive aspergillosis

Chronic pulmonary aspergillosis