

Abacavir

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors Oral > Solid: 300 mg tablet (as sulfate)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Didanosine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors - Oral > Liquid: 100 mg buffered powder for oral solution; 167 mg buffered powder for oral solution; 250 mg buffered powder for oral solution - Oral > Solid: 25 mg buffered chewable dispersible tablet; 50 mg buffered chewable dispersible tablet; 100 mg buffered chewable dispersible tablet; 150 mg buffered chewable dispersible tablet; 200 mg buffered chewable dispersible tablet; 125 mg unbuffered enteric coated capsule; 200 mg unbuffered enteric coated capsule; 250 mg unbuffered enteric coated capsule; 400 mg unbuffered enteric coated capsule	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Emtricitabine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors - Oral > Liquid: 10 mg per mL - Oral > Solid: 200 mg	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lamivudine

Section	Indications
Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors - Oral > Liquid: 50 mg per 5 mL - Oral > Solid: 150 mg tablet	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Stavudine

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Liquid: 5 mg per 5 mL powder for oral solution
- Oral > Solid: 15 mg; 20 mg; 30 mg

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Tenofovir disoproxil fumarate

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Oral > Solid: 300 mg tablet (equivalent to 245 mg tenofovir disoproxil)

Indications

Contact with or exposure to human immunodeficiency virus

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Zidovudine

Section

Antiretrovirals > Nucleoside/Nucleotide reverse transcriptase inhibitors

- Parenteral > General injections > IV: 10 mg per mL in 20 mL vial solution for IV infusion
- Oral > Liquid: 50 mg per 5 mL
- Oral > Solid > tablet: 300 mg
- Oral > Solid > capsule: 250 mg

Indications

Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified