Atazanavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 100 mg (as sulfate); 300 mg (as sulfate)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Atazanavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 300 mg (as sulfate) + 100 mg tablet (heat stable)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Darunavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 75 mg; 400 mg; 600 mg; 800 mg	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Indinavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 400 mg (as sulfate)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Lopinavir + ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid > tablet: 100 mg + 25 mg tablet (heat-stable); 200 mg + 50 mg tablet (heat-stable) ◦ Oral > Solid > dosage form: 40 mg + 10 mg	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Nelfinavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 250 mg (as mesilate) ◦ Oral > Other: 50 mg per g oral powder	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Ritonavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 25 mg tablet (heat stable); 100 mg tablet (heat stable)	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified

Saquinavir

Section	Indications
Antiretrovirals > Protease inhibitors ◦ Oral > Solid: 200 mg; 500 mg	Human immunodeficiency virus disease without mention of tuberculosis or malaria, clinical stage unspecified