

Benznidazole**Section**

American trypanosomiasis

- Oral > Solid > tablet: 100 mg (scored); 50 mg (scored); 12.5 mg

Indications

Chagas disease

Eflornithine**Section**

Medicines for the treatment of 2nd stage African trypanosomiasis

- Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)

Indications

Gambiense trypanosomiasis

Fexinidazole**Section**

Medicines for the treatment of 1st stage African trypanosomiasis

- Oral > Solid > tablet: 600 mg
- Oral > Solid: 600 mg

Medicines for the treatment of 2nd stage African trypanosomiasis

- Oral > Solid > tablet: 600 mg
- Oral > Solid: 600 mg

Indications

Rhodesiense trypanosomiasis

Gambiense trypanosomiasis

Rhodesiense trypanosomiasis

Gambiense trypanosomiasis

Melarsoprol**Section**

Medicines for the treatment of 2nd stage African trypanosomiasis

- Parenteral > General injections > IV: 3.6% in 5 mL ampoule solution (180 mg of active compound)

Indications

Gambiense trypanosomiasis

Rhodesiense trypanosomiasis

Nifurtimox**Section**

American trypanosomiasis

- Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

Medicines for the treatment of 2nd stage African trypanosomiasis

- Oral > Solid > tablet: 30 mg (scored); 120 mg (scored)

Indications

Chagas disease

Gambiense trypanosomiasis

Pentamidine

Section

Medicines for the treatment of 1st stage African trypanosomiasis

- Parenteral > General injections > IM: 300 mg in vial (as isetionate) powder for injection

Indications

Gambiense trypanosomiasis

Suramin sodium

Section

Medicines for the treatment of 1st stage African trypanosomiasis

- Parenteral > General injections > IV: 1 g in vial

Indications

Rhodesiense trypanosomiasis