



Model List of Essential Medicines



Anti-D immunoglobulin [General information](#)

Section

Human immunoglobulins

- Parenteral > General injections > IV: 250 µg in single-dose vial
- Parenteral > General injections > IM: 250 µg in single-dose vial

Indications

[Maternal care for red cell antibodies Rh incompatibility reaction](#)



Anti-rabies immunoglobulin [General information](#)

Section

Human immunoglobulins

- Parenteral > General injections > IM: 150 IU per mL in vial
- Parenteral > Locoregional injections > Other: 150 IU per mL in vial

Indications

[Rabies](#)



Anti-tetanus immunoglobulin [General information](#)

Section

Human immunoglobulins

- Parenteral > General injections > IM: 500 IU in vial

Indications

[Tetanus](#)



C1 esterase inhibitor [General information](#)

Section

Human immunoglobulins

- Parenteral > General injections > IV: 500 IU in vial powder for injection

Indications

[Hereditary angioedema](#)



Coagulation factor IX complex [General information](#)

Section

Coagulation factors

Indications

Therapeutic equivalent to [coagulation factor IX, plasma-derived](#) for [Coagulation defects, purpura or other haemorrhagic or related conditions](#) Therapeutic equivalent to [coagulation factor IX, plasma-derived](#) for [Haemophilia B](#)



Coagulation factor IX, plasma-derived [General information](#)

Section

Coagulation factors

- Parenteral > General injections > IV: 500 IU in vial powder for injection; 1000 IU in vial powder for injection

Indications

[Coagulation defects, purpura or other haemorrhagic or related conditions](#) [Haemophilia B](#)



Coagulation factor IX, recombinant [General information](#)

Section

Coagulation factors

- Parenteral > General injections > IV: 250 IU per mL in vial lyophilized powder for solution for injection; 500 IU in vial lyophilized powder for solution for injection; 1000 IU in vial lyophilized powder for solution for injection; 1500 IU in vial lyophilized powder for solution for injection; 2000 IU in vial lyophilized powder for solution for injection; 3000 IU in vial lyophilized powder for solution for injection; 4000 IU in vial lyophilized powder for solution for injection

Indications

[Haemophilia B](#)



Coagulation factor VIII, plasma-derived [General information](#)

Section

Coagulation factors

- Parenteral > General injections > IV: 250 IU in vial powder for injection; 500 IU in vial powder for injection; 1000 IU in vial powder for injection

Indications

[Haemophilia A](#)



Coagulation factor VIII, recombinant [General information](#)

Section

Coagulation factors

- Parenteral > General injections > IV: 250 IU in vial lyophilized powder for solution for injection; 500 IU in vial lyophilized powder for solution for injection; 1000 IU in vial lyophilized powder for solution for injection; 1500 IU in vial lyophilized powder for solution for injection; 2000 IU in vial lyophilized powder for solution for injection; 3000 IU in vial lyophilized powder for solution for injection; 4000 IU in vial lyophilized powder for solution for injection

Indications

[Haemophilia A](#)



Cryoprecipitate (native) [General information](#)

Section

Blood and blood components

Indications

Therapeutic equivalent to [cryoprecipitate \(pathogen-reduced\)](#) for [Coagulation defects, purpura or other haemorrhagic or related conditions](#) Therapeutic equivalent to [cryoprecipitate \(pathogen-reduced\)](#) for [Obstetric haemorrhage](#) Therapeutic equivalent to [cryoprecipitate \(pathogen-reduced\)](#) for [Haemophilia A](#) Therapeutic equivalent to [cryoprecipitate \(pathogen-reduced\)](#) for [Von Willebrand disease](#)



Cryoprecipitate (pathogen-reduced) [General information](#)

Section

Blood and blood components

- Injection: frozen liquid in bag or lyophilized powder in vial containing:
 - > 50 IU Factor VIII
 - > 100 IU vWF
 - > 140 mg clottable fibrinogen per unit

Indications

[Coagulation defects, purpura or other haemorrhagic or related conditions](#) [Obstetric haemorrhage](#) [Haemophilia A](#) [Von Willebrand disease](#)



Dextran 40 [General information](#)

Section
Plasma substitutes

◦ .

Indications
[Hypovolaemia](#)



Dextran 70 [General information](#)

Section
Plasma substitutes

◦ Parenteral > General injections > IV: 6% injectable solution

Indications
[Hypovolaemia](#)



Fresh-frozen plasma [General information](#)

Section
Blood and blood components

◦ Parenteral > General injections > IV:

Indications
[Blood transfusion without reported diagnosis](#)



Normal immunoglobulin [General information](#)

Section
Human immunoglobulins

- Parenteral > General injections > IV: 5% protein solution; 10% protein solution
- Parenteral > General injections > IM: 16% protein solution
- Parenteral > General injections > SC: 15% protein solution; 16% protein solution

Indications
[Primary immunodeficiencies Mucocutaneous lymph node syndrome Langerhans cell histiocytosis](#)



Platelets [General information](#)

Section
Blood and blood components

◦ Parenteral > General injections > IV:

Indications
[Blood transfusion without reported diagnosis](#)



Polygeline [General information](#)

Section
Plasma substitutes

- Parenteral > General injections > IV: 3.5% injectable solution
- Parenteral > General injections > IV: 3.5% in solution

Indications
Therapeutic equivalent to [dextran 70](#) for [Hypovolaemia Hypovolaemia](#)



Red blood cells [General information](#)

Section
Blood and blood components

◦ Parenteral > General injections > IV:

Indications
[Blood transfusion without reported diagnosis](#)



Whole blood [General information](#)

Section
Blood and blood components

◦ Parenteral > General injections > IV:

Indications
[Blood transfusion without reported diagnosis](#)