



Model List of Essential Medicines

Beinaglutide [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) for [Obesity](#)

Canagliflozin [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [empagliflozin](#) for [Type 2 diabetes mellitus](#)

Dapagliflozin [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [empagliflozin](#) for [Type 2 diabetes mellitus](#)

Dulaglutide [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) for [Obesity](#) Therapeutic equivalent to [semaglutide](#) for [Type 2 diabetes mellitus](#)

Empagliflozin [General information](#)

Section

Hypoglycaemic agents

◦ Oral > Solid: 10 mg; 25 mg

Indications

[Type 2 diabetes mellitus](#)

Exenatide [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) for [Obesity](#)

Glibenclamide [General information](#)

Section

Hypoglycaemic agents

- Oral > Solid: 5 mg; 2.5 mg

Indications

[Type 2 diabetes mellitus](#)

Gliclazide [General information](#)

Section

Hypoglycaemic agents

- Oral > Solid: 30 mg controlled-release; 60 mg controlled-release; 80 mg controlled-release

Indications

[Type 2 diabetes mellitus](#)

Insulin (analogue, long-acting) [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 IU per mL in 3 mL cartridge solution; 100 IU per mL in 3 mL pre-filled pen solution; 100 IU per mL in 10 mL vial solution

Indications

[Type 2 diabetes mellitus](#) [Type 1 diabetes mellitus](#)

Insulin (analogue, rapid-acting) [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 IU per 1 mL in 10 mL vial; 100 IU per 1 mL in 3 mL cartridge; 100 IU per 1 mL in 3 mL pre-filled pen

Indications

[Type 1 diabetes mellitus](#) [Type 2 diabetes mellitus](#)

Insulin (human, intermediate-acting) [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 40 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 10 mL vial as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL cartridge as compound insulin zinc suspension or isophane insulin; 100 IU per mL in 3 mL pre-filled pen as compound insulin zinc suspension or isophane insulin

Indications

[Type 2 diabetes mellitus](#) [Type 1 diabetes mellitus](#)

Insulin (human, short-acting) [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 40 IU per mL in 10 mL vial solution; 100 IU per mL in 10 mL vial solution; 100 IU per mL in 3 mL cartridge solution; 100 IU per mL in 3 mL pre-filled pen solution

Indications

[Type 2 diabetes mellitus](#) [Type 1 diabetes mellitus](#)

Insulin analogues [General information](#)

Section

Insulins

- Not specified

Indications

[Type 2 diabetes mellitus](#) [Type 1 diabetes mellitus](#)

Insulin aspart [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 IU per mL in 110 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen
- Parenteral > General injections > SC: 100 IU per mL in 10 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

Indications

Therapeutic equivalent to [insulin \(analogue, rapid-acting\)](#) for [Type 1 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, rapid-acting\)](#) for [Type 2 diabetes mellitus](#)

Insulin degludec [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 units per mL

Indications

Therapeutic equivalent to [insulin glargine](#) for [Type 1 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, long-acting\)](#) for [Type 2 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, long-acting\)](#) for [Type 1 diabetes mellitus](#)

Insulin detemir [General information](#)

Section

Insulins

Indications

Therapeutic equivalent to [insulin glargine](#) for [Type 1 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, long-acting\)](#) for [Type 2 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, long-acting\)](#) for [Type 1 diabetes mellitus](#)

Insulin glargine [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 units per mL

Indications

[Type 1 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, long-acting\)](#) for [Type 2 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, long-acting\)](#) for [Type 1 diabetes mellitus](#)

Insulin glulisine [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 IU per mL in 10 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

Indications

Therapeutic equivalent to [insulin \(analogue, rapid-acting\)](#) for [Type 1 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, rapid-acting\)](#) for [Type 2 diabetes mellitus](#)

Insulin lispro [General information](#)

Section

Insulins

- Parenteral > General injections > SC: 100 IU per mL in 10 mL vial; 100 IU per mL in 3 mL cartridge; 100 IU per mL in 3 mL pre-filled pen

Indications

Therapeutic equivalent to [insulin \(analogue, rapid-acting\)](#) for [Type 1 diabetes mellitus](#) Therapeutic equivalent to [insulin \(analogue, rapid-acting\)](#) for [Type 2 diabetes mellitus](#)

Liraglutide [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) for [Obesity](#) Therapeutic equivalent to [semaglutide](#) for [Type 2 diabetes mellitus](#)

Metformin [General information](#)

Section

Hypoglycaemic agents

- Oral > Solid: 500 mg (hydrochloride)

Indications

[Type 2 diabetes mellitus](#)

Semaglutide [General information](#)

Section

Hypoglycaemic agents

- Parenteral > General injections > SC: 0.25 mg in pre-filled pen injection solution; 0.5 mg in pre-filled pen injection solution; 1 mg in pre-filled pen injection solution; 2 mg in pre-filled pen injection solution; 4 mg in pre-filled pen injection solution
- Oral > Solid > tablet: 3 mg
- Parenteral > General injections > SC: 0.68 mg per mL; 1.34 mg per mL; 2.68 mg per mL

Indications

[Obesity](#) [Type 2 diabetes mellitus](#)

Tirzepatide [General information](#)

Section

Hypoglycaemic agents

Indications

Therapeutic equivalent to [semaglutide](#) for [Obesity](#) Therapeutic equivalent to [semaglutide](#) for [Type 2 diabetes mellitus](#)

Tolbutamide [General information](#)

Section

Hypoglycaemic agents

- Oral > Solid: 500 mg

Indications

[Type 2 diabetes mellitus](#)