

Acamprosate calcium**Section**

Medicines for alcohol use disorders

- Oral > Solid > tablet: 333 mg

Indications

Disorders due to use of alcohol

Buprenorphine**Section**

Medicines for opioid use disorders

IndicationsTherapeutic equivalent to [methadone](#) for

Opioid dependence

Bupropion**Section**

Medicines for nicotine use disorders

- Oral > Solid: 150 mg sustained-release (hydrochloride)

Indications

Nicotine dependence

Cytisine (cytisinicline)**Section**

Medicines for nicotine use disorders

- Oral > Solid > tablet: 1.5 mg

Indications

Nicotine dependence

Methadone**Section**

Medicines for opioid use disorders

- Oral > Liquid: 5 mg per 5 mL (methadone hydrochloride); 10 mg per 5 mL (methadone hydrochloride); 5 mg per mL concentrate for oral liquid (methadone hydrochloride); 10 mg per mL concentrate for oral liquid (methadone hydrochloride)

Indications

Opioid dependence

Naltrexone**Section**

Medicines for alcohol use disorders

- Parenteral > General injections > IM: 380 mg in vial (extended-release)
- Oral > Solid > tablet: 50 mg

Indications

Disorders due to use of alcohol

Nicotine replacement therapy

Section	Indications
Medicines for nicotine use disorders ◦ Local > Topical > Transdermal patch: 5 to 30 mg per 16 hour; 7 to 21 mg per 24 hour ◦ Oral > Solid > lozenge: 2 mg; 4 mg lozenge ◦ Local > Buccal > Oral spray: 1 mg per actuation ◦ Local > Buccal > Chewing gum: 2 mg; 4 mg	Nicotine dependence

Varenicline

Section	Indications
Medicines for nicotine use disorders ◦ Oral > Solid: 0.5 mg; 1 mg	Nicotine dependence