Model List of Essential Medicines

Acetylsalicylic acid

Section	Indications
Medicines for juvenile joint diseases ◦ Local > Rectal > Suppository: 150 mg; 300 mg ◦ Oral > Solid > dispersible tablet: 75 mg; 300 mg; 500 mg ◦ Oral > Solid > tablet: 75 to 500 mg	Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis Acute rheumatic fever without mention of heart involvement

Adalimumab

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

Allopurinol

Section	Indications
Medicines used to treat gout ◦ Oral > Solid: 100 mg	Gout

Anakinra

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	Juvenile systemic arthritis

Azathioprine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid > tablet: 50 mg (scored)	Psoriatic arthritis Rheumatoid arthritis

Certolizumab pegol

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Chloroquine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid: 100 mg tablet (as phosphate or sulfate); 150 mg tablet (as phosphate or sulfate)	Rheumatoid arthritis

Colchicine

Section	Indications
Medicines used to treat gout ◦ Oral > Solid: 0.5 mg	Gout

Etanercept

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Golimumab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Hydroxychloroquine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid: 200 mg (as sulfate)	Systemic lupus erythematosus Cutaneous lupus erythematosus of unspecified type

Infliximab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Methotrexate

Section	Indications
Medicines for juvenile joint diseases ◦ Oral > Solid > tablet: 2.5 mg (as sodium salt)	Juvenile idiopathic arthritis
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid > tablet: 2.5 mg (as sodium salt)	Rheumatoid arthritis

Penicillamine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid: 250 mg	Rheumatoid arthritis

Probenecid

Section	Indications
Medicines used to treat gout ◦ Oral > Solid: 500 mg	Gout

Sulfasalazine

Section	Indications
Disease-modifying anti-rheumatic drugs (DMARDs) ◦ Oral > Solid: 500 mg	Rheumatoid arthritis

Tocilizumab

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial ◦ Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe	Juvenile systemic arthritis

Triamcinolone hexacetonide

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial	Juvenile idiopathic arthritis