Model List of Essential Medicines

Acetylsalicylic acid

Section	Indications
Medicines for juvenile joint diseases ◦ Local > Rectal > Suppository: 150 mg; 300 mg ◦ Oral > Solid > dispersible tablet: 75 mg; 300 mg; 500 mg ◦ Oral > Solid > tablet: 75 to 500 mg	Mucocutaneous lymph node syndrome Juvenile idiopathic arthritis Acute rheumatic fever without mention of heart involvement

Adalimumab

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL	Juvenile idiopathic arthritis

Anakinra

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe	Juvenile systemic arthritis

Certolizumab pegol

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Etanercept

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Golimumab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Infliximab

Section	Indications
Medicines for juvenile joint diseases	Therapeutic equivalent to adalimumab for Juvenile idiopathic arthritis

Methotrexate

Section	Indications
Medicines for juvenile joint diseases ◦ Oral > Solid > tablet: 2.5 mg (as sodium salt)	Juvenile idiopathic arthritis

Tocilizumab

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial ◦ Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe	Juvenile systemic arthritis

Triamcinolone hexacetonide

Section	Indications
Medicines for juvenile joint diseases ◦ Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial	Juvenile idiopathic arthritis