



## Model List of Essential Medicines



### Acetylsalicylic acid [General information](#)

Section

Medicines for juvenile joint diseases

- Local > Rectal > Suppository: 150 mg; 300 mg
- Oral > Solid > dispersible tablet: 75 mg; 300 mg; 500 mg
- Oral > Solid > tablet: 75 to 500 mg

Indications

[Acute rheumatic fever without mention of heart involvement](#) [Mucocutaneous lymph node syndrome](#) [Juvenile idiopathic arthritis](#)



### Adalimumab [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > General injections > SC: 40 mg per 0.8 mL; 40 mg per 0.4 mL; 10 mg per 0.2 mL; 20 mg per 0.4 mL

Indications

[Juvenile idiopathic arthritis](#)



### Anakinra [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > General injections > SC: 100 mg per 0.67 mL in pre-filled syringe

Indications

[Juvenile systemic arthritis](#)



### Certolizumab pegol [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)



### Etanercept [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)



**Golimumab** [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)

**Infliximab** [General information](#)

Section

Medicines for juvenile joint diseases

Indications

Therapeutic equivalent to [adalimumab](#) for [Juvenile idiopathic arthritis](#)

**Methotrexate** [General information](#)

Section

Medicines for juvenile joint diseases

- Oral > Solid > tablet: 2.5 mg (as sodium salt)

Indications

[Juvenile idiopathic arthritis](#)

**Tocilizumab** [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > General injections > IV: 80 mg per 4 mL in vial; 200 mg per 10 mL in vial; 400 mg per 20 mL in vial
- Parenteral > General injections > SC: 162 mg per 0.9 mL in pre-filled syringe

Indications

[Juvenile systemic arthritis](#)

**Triamcinolone hexacetonide** [General information](#)

Section

Medicines for juvenile joint diseases

- Parenteral > Locoregional injections > Intra-articular: 20 mg per mL in vial

Indications

[Juvenile idiopathic arthritis](#)