Liste modèle des médicaments essentiels

Metronidazole

Section	Indications
Antiamoebic and antigiardiasis medicines ◦ Oral > Liquid: 200 mg per 5 mL (as benzoate) ◦ Parenteral > General injections > unspecified: 500 mg in 100 mL vial ◦ Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg	Amoebiasis

Access group antibiotics

- Oral > Liquid: 200 mg per 5 mL (as benzoate)
- Local > Rectal > Suppository: 500 mg; 1 g
- Parenteral > General injections > unspecified: 500 mg per 100 mL in vial
- Oral > Solid > tablet: 200 mg; 250 mg; 400 mg; 500 mg
- Oral > Solid > tablet: 500 mg; 200 mg; 250 mg; 400 mg
- Oral > Solid: 200 to 500 mg tablet
- Parenteral > General injections > unspecified: 500 mg per 100 mL in vial Injection

Premier choix

Trichomoniasis

Intestinal infections due to Clostridioides difficile

Necrotising fasciitis

co-prescrite avec cefazolin

Other specified prophylactic measures

co-prescrite avec cefotaxime

Peritoneal abscess (mild-moderate)

co-prescrite avec cefotaxime

Peritoneal abscess (severe)

co-prescrite avec cefotaxime

Peritonitis (mild-moderate)

co-prescrite avec cefotaxime

Peritonitis (severe)

co-prescrite avec ceftriaxone

Peritoneal abscess (mild-moderate)

co-prescrite avec ceftriaxone

Peritoneal abscess (severe)

co-prescrite avec ceftriaxone

Peritonitis (mild-moderate)

co-prescrite avec ceftriaxone

Peritonitis (severe)

Second choix

co-prescrite avec ciprofloxacin

Peritoneal abscess (mild-moderate)

co-prescrite avec ciprofloxacin

Peritonitis (mild-moderate)

Bacterial infection of unspecified site